

CONVENTION ON INTERNATIONAL TRADE IN ENDANGERED SPECIES
OF WILD FAUNA AND FLORA



Sixteenth Meeting of the CITES Animals Committee
Shepherdstown (United States of America) 11-15 December 2000

Implementation of Resolution Conf. 8.9 (Rev.)

ACIPENSERIFORMES

1. This document has been prepared by TRAFFIC International in cooperation with IUCN and UNEP-WCMC, under contract to the CITES Secretariat.
2. The Annex consists of a review of 10 species of Acipenseriformes pursuant to Decision 11.95 directing the Animals Committee to include the Acipenseriformes in its Review of Significant Trade. The document deals with the following species of Acipenseriformes:

<i>Acipenser baerii</i>	p3
<i>Acipenser fulvescens</i>	p11
<i>Acipenser gueldenstaedtii</i>	p19
<i>Acipenser nudiiventris</i>	p43
<i>Acipenser ruthenus</i>	p55
<i>Acipenser schrencki</i>	p65
<i>Acipenser stellatus</i>	p71
<i>Huso dauricus</i>	p97
<i>Huso huso</i>	p105
<i>Polyodon spathula</i>	p127

3. The relevant parts of the Annex have been sent to all range States of the species concerned, but comments from the range States have either not yet been received or not yet been incorporated in the document.

Acipenser baerii Brandt, 1869

Siberian Sturgeon
Esturgeon sibérien

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

Siberian Sturgeon *Acipenser baerii* may live for up to sixty years and generally reach sexual maturity between 18 and 28 years of age, except in the Lena River where fish mature sexually around 9-12 years of age. The maximum weight recorded is 210 kg, but *A. baerii* usually weigh approximately 65 kg. Little information is available on the current population status for this species in China and Kazakhstan. However, the limited data available for the Russian Federation show that the spawning area has declined by up to 40% in some areas due to dam construction and increased abnormal oogenesis has been observed, probably due to chemical and nuclear water pollution. The population in China is low and economically insignificant.

Catch volumes in the Russian Federation have generally declined, probably as a result of the above listed anthropogenic disturbances as well as over-fishing. However, the Russian Federation is the only country where significant levels of catches have been reported. The annual catch of *A. baerii* in the Ob'-Irtysk River basin decreased from 1,410 tonnes (t) in the 1930s to 11 t in 1997. Annual catches in the two other major Siberian rivers in the late 1990s were 16 t per year in the Yenisey River and 13-20 t per year in the Lena River. The Lena population of *A. baerii* is most commonly bred in captivity, because it completes its life cycle in freshwater and sexually matures relatively early. Virtually all international trade in *A. baerii* (caviar, fertilised eggs and live fish) for 1998 detailed in CITES annual reports was from captive-bred sources from France (280 kg of caviar and 160,000 live juveniles) and the Russian Federation (probably fertilised eggs). The export quota set for caviar by the Russian Federation declined from 19,000kg for 1998 to 400 kg for 2000. Additionally, for 2000 the Russian Federation has set export quotas of 3,000 kg of meat and fertilised eggs of *A. baerii* as well as of three different hybrids.

The species is recommended under Decision 11.106 for inclusion in category 2.

DISTRIBUTION AND POPULATION

The UNEP-WCMC database of CITES-listed species lists the distribution of *Acipenser baerii* as: China, Kazakhstan? and the Russian Federation (Anon., 2000a).

A. baerii is classified as Vulnerable by IUCN (1996):

VU A2d Russia

A. b. baerii EN A2d Russia [Ob and Taz rivers]

A. b. baikalensis EN A1ace Russia [Lake Baikal]

A. b. stenorrhynchus VU A2d Russia [Yenisei, Lena and east Siberian rivers]

China: There is little information about the distribution of *A. baerii* in China, but it is reported to occur within the Irtysk River system of Xinjiang (Lake Zaysan, Kara, Irtysk and Ertix rivers, tributaries of the Irtysk River) (Qing Jianhua, pers. comm., cited in: CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). Ruban notes that *A. baerii* inhabits the part of the Irtysk River (Tchernyi Irtysk) flowing through China to the Kren River (Ruban, 1997; 1999).

Kazakhstan: *A. baerii* inhabits part of the Irtysk River including Shulbinskoe, Ust-Kmenogorskoe and Bukhtarminskoe reservoirs (Ruban, 1999).

Russian Federation: *A. baerii* inhabits virtually all large rivers in Siberia (Sokolov and Vasil'ev, 1989). Its range extends south from the Lena River basin and Ob' Bay to the Cherniy, Irtysk and Selenga rivers. Longitudinally, it is found from the Ob' to the Kolmya rivers. The populations inhabiting the Ob'-Irtysk basin

and Lake Baikal are included in the Russian Federation Red Book, with stocks having declined throughout the country (Dr. G. Ruban, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Three subspecies are recognised. *A. baerii baerii* is restricted to the Ob' River and its tributaries. This subspecies migrates to Ob' Bay during the winter due to seasonal oxygen deficiency in the Ob' River; numbers are reported to be very low (Ruban, 1996; 1997).

A. baerii stenorrhynchus occurs in the basins of the East Siberian Rivers (Alazeya, Anabar, Indigirka, Khatanga, Lena, Lolya, Olenek, Pyasina, Yana and Yenisei. (Ruban, 1997; Ruban and Akimova, 1991; 1993; Ruban and Panaiotidi, 1994). Two forms of this subspecies are recognised. The migrating form, which swims considerable distances upstream from estuaries and deltas to spawn, is more numerous than the non-migratory form. The constant reduction in fisheries catch statistics reflects declining populations (Dr G. Ruban, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

A. baerii baikalensis is a unique lake form from Lake Baikal, which migrates to the Selenga River to spawn. The majority of the population occurs in the northern part of the lake near to the Selenga River delta and the Barguzinskii and Chivyrkuuskii bays, and is rarely encountered in the south. From the northern areas it migrates into the large tributaries of the lake, up to 1,000 km upstream in the Selenga River, and also enters the tributaries of the Chkoi, Delger-Muren, Orhon and Tula rivers. In the mid-1990s, the subspecies was reported to be extremely rare (Ruban, 1997).

However, recent studies suggest that *A. baerii* may be monotypic and forms continuous populations in large river systems (Ruban, 1999).

Introduced populations:

France: In December 1999 several thousand juvenile and several hundred gravid females (more than 7 kg specimens) of *A. baerii* escaped into the Gironde River (Bordeaux region) during two storms. The survival of the escaped fish and their effect on the wild population of *A. sturio*, are unknown. However, the introduction of new pathological germs, food competition in case of acclimatisation of the exotic specimens, and hybridisation with *A. sturio* must be taken into consideration (Cemagref, *in litt.*, press release, 26 January 2000).

Germany: In the Baltic Sea river estuaries east of Rostock (Germany) and further upstream, hybrid sturgeon, including hybrids with *A. baerii*, have been caught. These probably originate from upstream aquaculture farms, or were released by private aquaria due to their large size (Jörn Gessner, Institute of Freshwater Ecology and Inland Fisheries, pers. comm. to TRAFFIC Europe, 15 March 2000).

Italy: The species has been introduced in captive breeding facilities and hybridised with Adriatic Sturgeon *A. naccarii* in Italy in the 1990s (Azienda Agricola and Agroittica Lombarda, Italian sturgeon farmers, *in litt.* to TRAFFIC Europe-Italy, 1999). *A. baerii* are occasionally found in the wild; fish sporadically escape from rearing plants or angling ponds, or are released when they become too large for private aquaria (Dr P. Bronzi *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). There is no documentation on the potential damage of the introduction of exotic Acipenseriformes and their hybrids on native species. If specimens of *A. baerii* escape to the open waters of the Po River and become an "invasive species", this may threaten the Adriatic Sturgeon *A. naccarii*, a species that is on the brink of extinction.

Russian Federation: *A. baerii* has been introduced to Lakes Ladoga, Pskov-Chud, Seliger and others in the Baltic watershed and to the Gor'kov and Volgograd impoundment along the Volga and in manmade lakes downstream from Moscow (Berdichevskii *et al.*, 1983, cited in Sokolov and Vasil'ev, 1989). However, long-term studies have rarely found introduced *A. baerii* in these water bodies (V. S. Malyutin, pers. comm., cited in Dr G. Ruban, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

HABITAT AND ECOLOGY

A. baerii can reach a maximum length of 2 m and weight of 210 kg. However, it usually does not exceed 65 kg in weight with a maximum age of approximately 60 years (Sokolov and Vasil'ev, 1989). Only the

Lena River population reaches sexual maturity at 9-10 years for males and 10-12 years for females; all other populations reach sexual maturity at 18-24 years for males and 24-28 years for females. The minimum recorded size for spawning is 0.6-0.9 m in length and 0.7 kg in weight (Hochleithner and Gessner, 1999). The spawning season is from May to June. *A. baerii* feeds predominantly on benthic organisms including chironomid larvae and river amphipods, isopods and polychaetes (Sokolov and Vasil'ev, 1989).

The results of *A. baerii* captive breeding suggest that young females (two years after reaching sexual maturity, minimum age ten years), born and bred in captivity, may bear 10% of their weight in caviar, while for 18-year old females the yield is 18-20% (T. Rouault, Cemagref, pers. comm. to TRAFFIC Europe, 27 September 2000). Therefore the average rate most commonly used is 12-13% per female and 6-7% for the whole catch, assuming that 50% of the catch are males (1:1 sex ratio of the population).

THREATS TO SURVIVAL AND DOMESTIC USE

China: The size of the population is relatively small and economically insignificant (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

Russian Federation: Much of the habitat of *A. baerii* has been lost due to the damming of many Siberian rivers since the 1950s. The migratory forms have been particularly affected because they can no longer access their spawning sites. There is little documentation as to the extent of habitat loss.

Forty percent of the spawning habitat of the migrating form of *A. baerii stenorrhynchus* has been lost in the Ob'-Irtys' River due to construction of the Novosibirsk, Ust'-Kamengorsk and Shul'binsk hydroelectric power stations (Ruban, 1996). The Yenisey River and Lena River populations have been similarly affected, losing 500-600 km and 300 km of habitat respectively. The development of oil extraction facilities and the use of pesticide in agriculture in the Yenisey River have polluted Siberian rivers (Ruban, 1997). Pollution has drastic effects on fish physiology and has consequently reduced the already low fecundity of all sturgeon species by causing degenerative oocytes (Akimova and Ruban, 1995). In the 1980s, oil and petroleum products in the Ob' River exceeded permissible levels by a factor of ten (Green, 1993). Many Siberian rivers are contaminated by radioactive substances and eastern Siberian rivers are polluted by the gold mining industry (Akimova and Ruban, 1995).

Law enforcement appears to be lacking and numerous experts as well as high position officers claim that the wild population is highly threatened by increasing illegal fishing and trade in caviar in a large part of the range (Anon., 2000b).

In the 1930s, the annual recorded catch of *A. baerii* in the Ob'-Irtys' rivers basin reached 1,410 t. By 1997, this figure had dropped to 11 t (Ruban, 1999). The ratio between illegal and legal catch in the Ob' River in 1994 was estimated at 300 t to 11 t (Ruban, 1996).

Russian catch of *A. baerii* in three large Siberian rivers (tonnes)

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
Ob'	73	74	48	63	48	57	26	15.9	17	11.5	n/a	6.7	11
Yenisey	23.4	28.1	30.4	20.7	17	40.6	36.9	31.1	42.4	44.1	n/a	33.6	n/a
Lena	8.8	11.2	8.7	15.2	8.2	9.5	14.9	4.9	5.4	17.2	12.9	21.6	9
Total	105.2	113.3	87.1	98.9	73.2	107.1	77.8	51.9	64.8	72.8		61.9	

Source: Ruban, 1999

Current catch in the Yenisey River does not exceed 16 t per year, and the Lena River annual catches are approximately 13-20 t. In other Siberian rivers (Anabar, Indigirka, Khatanga, Kolyma, Olenek and Yana) stocks have always been relatively small. There are no recent catch data available for Yakutian (Republic Sakha-Yakutiya of the Russian Federation) rivers (Anabar, Indigirka, Khatanga, Kolyma, Lena, Olenek and Yana) (Anon., 2000b).

INTERNATIONAL TRADE

The listing of *A. baerii* in CITES Appendix II entered into effect on 1 April 1998. Complete data for this species were therefore limited to nine months of trade (April-December 1998).

Gross exports and the comparative tabulation of trade in *A. baerii* in 1998 are given in the Appendix. CITES Annual Reports for 1998 suggest that the international trade in wild specimens of *A. baerii* was extremely limited in 1998: 283 kg "eggs", 360,950 live fish (including 160,000 of captive-bred sources from France and 200,000 of unknown origin that were seized by the US Customs) and 433 kg of meat of captive-bred source from Uruguay. The export of 283 kg "eggs", comprised 280 kg caviar of captive-bred source from France and 3 kg of eggs (probably fertilised eggs destined for captive breeding facilities, but could be caviar; see *Note* at the end of the section) from the Russian Federation to Greece. Together with *A. transmontanus* White Sturgeon, *A. baerii* is one of the most common sturgeon species bred in captivity for meat. Captive breeding of *A. baerii* for caviar is limited to France, with the majority of caviar being consumed within the European Union (EU). Internal EU trade does not require CITES permits and is therefore not recorded in CITES Annual Reports.

"Live" specimens and fertilised "eggs" were probably destined to supply aquaculture farms and the growing demand for private aquaria. Germany and Hungary are also involved in the international trade of live first generation and captive-bred specimens respectively. "Meat" is the major production of sturgeon farms (see "Captive Breeding").

Exports of *A. baerii* from Russia in 1999 (kilogrammes)

	Quantity	Description	Importer	No of permits
<i>A. baerii</i>	27.50	Live eggs	HU, PL, DE, US, AM, IT, SY	10 Export
	300.00	Live fish	KR	2 Export
Hybrids:				
<i>A. baerii</i> x <i>A. gueldenstaedtii</i>	3.50	Live eggs	PL, DE	2 Export
<i>A. gueldenstaedtii</i> x <i>A. baerii</i>	37.50	Live eggs	CN, SY	3 Export

Source: CITES Management Authority, *in litt.* to TRAFFIC Europe, 18 September 2000

Export quotas set for *A. baerii* and its hybrids (kilogrammes)

	1998	1999	2000	
	Caviar	Caviar	Caviar	Meat Fertilised Eggs
Russian Federation	19,000	2,000	400	3,000 30
Hybrids:				
<i>A. baerii</i> x <i>A. gueldenstaedtii</i>				10
<i>A. gueldenstaedtii</i> x <i>A. baerii</i>				70
<i>A. baerii</i> x <i>A. ruthenus</i>				15
Total	19,000	2,000	400	3,000 125

Source: CITES Notification No. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Illegal trade: A large shipment (200,000 live fish) of *A. baerii* of unknown origin was confiscated by the US Customs. Poaching and smuggling are closely related and have been intensively reported in the media of range States and importing countries (Evtouchenko, 1997; Mc Donald, 2000; Snyder, 2000). However, a great deal of illegally caught sturgeon is destined for the domestic market, particularly meat (Anon., 1998).

Note: The use of the term "eggs" and the unit "kg" or "g" to describe shipments of both caviar and fertilised eggs complicates the analysis of trade data. Similar difficulties arise from the unit used for export quotas of fertilised eggs, which is kilogramme ("kg") instead of "number of specimens". Kilogrammes do not directly correspond to the number of specimens, and hence do not allow the effective evaluation of quotas.

CONSERVATION MEASURES

China: Insufficient studies have been undertaken to determine necessary conservation measures for wild populations of *A. baerii* (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

Russian Federation: The Lake Baikal population was listed in the Red Data Book in 1983 (Ruban, 1997) and the Ob' River population in 1998. In the late 1990s, commercial catch in the Yenisey River was prohibited due to rapidly declining populations and their extremely low fecundity (Anon., 2000b), caused by inaccessible spawning grounds and impact of pollution (Ruban, 1997). Catch for scientific and monitoring purposes, as well as subsistence use is limited to 8 t for *A. baerii* (Anon., 2000b).

Available data suggest that amendments of previous regulations and/or adoption of new legislation are in effect in Russian Federation. However, enforcement measures appear to be lacking and numerous experts as well as high-ranking officials have reported the increasing pressure of illegal fishing practices and criminal activities around caviar trade in a large part of the range (Anon., 2000b).

Russian scientists hold a few specimens of *A. baerii baikalensis* in a hatchery near Moscow for research and conservation (gene bank) purposes (Birstein, 1993). Approximately 300,000 larvae are reintroduced annually to Lake Baikal (V.S. Malyutin, pers. comm. to G. Ruban, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

CAPTIVE BREEDING

A. baerii is one of the most frequently captive-bred species of sturgeon. *A. baerii stenorrhynchus* have been successfully bred in several European countries (Sokolov and Vasil'ev, 1989; Willot *et al.*, 1991).

Caviar is not the main product of sturgeon farms. The economic viability of a sturgeon farm depends on the existence of a meat market, particularly until females have reached sexual maturity, during the first 8 to 10 years. Once the stock of caviar producers is constituted, one tonne of caviar per year is correlated to the production of 20 t of sturgeon meat (C. Raymakers, TRAFFIC Europe, *in litt.* to Pr R. Billard, National Natural History Museum, Paris, May 1999).

Western and Central Europe: M. Hochleithner (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) specifically names the following countries: Austria, Belgium, France, Germany, Hungary, Italy and Spain. In 1997, Bronzi *et al.* (1999) estimated the total annual captive-bred production of Siberian Sturgeon to be approximately 200 t in France, 120 t in Italy (pure species and hybrids), and 10 t in Germany.

Bulgaria: Only one Bulgarian hatchery exists. Established in 1997 in Boliartzi village (Plovdiv region), the hatchery has a total capacity of 60,000 juveniles. All juveniles released in the Bulgarian part of the Danube River from 1998 to 2000 were produced here. In 1998, a total of 200 hybrid fingerlings of *A. gueldenstaedtii* x *A. baerii*, with an average weight of 200 g, were released in the Ovcharitza dam reservoir. Since 1999, attempts have been made to breed sturgeon in artificial water basins near Sofia (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

China: In recent years, more than 200,000 fertilised eggs of *A. baerii* have been imported by China from Germany and the Russian Federation for aquaculture. Although not recorded in the CITES Annual Reports, small quantities of fingerlings and live juvenile *A. baerii* of captive bred sources were exported from China in 1998 (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

France: The production of *A. baerii* caviar from aquaculture exceeded three tonnes in 1999 and is expected to reach five tonnes in 2000 (A. Jones, Manager of a *A. baerii* farm in France, pers. comm. to TRAFFIC Europe, 22 September 2000). Based on estimates made (C. Raymakers, TRAFFIC Europe, *in litt.* to Pr R. Billard, National Natural History Museum, Paris, May 1999), this implies a minimum annual production of 100 t of meat of farmed Siberian Sturgeon.

Italy: *A. baerii* has been bred for human consumption (meat) and sport angling in private closed basins since 1997. These specimens originated from larvae imported from France, Hungary and, since 1998, Italy (Dr P. Bronzi *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Japan: *A. baerii* has been introduced into aquaculture in Japan (Katsumi and Genjiroi, 1977) but there are no production figures available.

Uruguay: According to CITES data, the US imported *A. baerii* meat of captive-bred source from Uruguay.

REFERENCES

- Akimova N.V., and Ruban G.I. (1995). *Reaction of reproductive system of sturgeons on anthropogenic impacts as a factor of population dynamics// Ecology of populations: structure and dynamics*. Ioskow. Part II. Pp. 491-500 (in Russian).
- Anon. (1996). *A Guide to Interpreting Outputs from the WCMC CITES Trade Database*, Produced by the World Conservation Monitoring Centre, Cambridge UK. Pp. 16
- Anon. (1998). *Sturgeon catch and trade in the Russian part of the Caspian Sea*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, 21 pp.
- Anon. (2000a). <http://www.cites.org/eng/dbase/fauna>
- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, 23 pp.
- Berdichevskii L.S. Malyutin V.S., Smolyanov I.I., Sokolov L.I. (1983). *The results of aquaculture and acclimatization researches of the Siberian Sturgeon*. Pages 259-269 In: I.A. Barannikova and L.S. Berdichevskii (eds.) *Biologicheskie osnovy osetrovodstva*. Nauka, Moskva (in Russian).
- Bronzi, P., Rosenthal, H., Arlati, G., and Williot, P. (1999). A brief overview on the status and prospects of sturgeon farming in Western and Central Europe In: *Proceedings of the "3rd International Symposium on Sturgeon"*, *J. Appl. Ichthyol.* 15(4-5): 224-227.
- Friend, T. (1997). DNA analysis busts animal smugglers. *USA Today*, 18 November 1997.
- Green, E. (1993). Poisoned legacy: Environmental Quality in the newly independent States. *Environ. Sc. and Technol.* 27: 590-595.
- Hochleithner, M. and Gessner, J. (1999). *The Sturgeon and Paddlefishes (Acipenseriformes) of the World: Biology and Aquaculture*. AquaTech Publications. Pp. 165.
- IUCN. (1996). *IUCN Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- McDonald, L. (2000). Fish that laid the golden egg faces extinction. *Daily Express* (UK), 26 July 2000.
- Ruban G.I., and Akimova N.V. (1991). [Ecological Characteristics of Siberian Sturgeon, *Acipenser baeri*, from the Indigirka River]. *Voprosy ichthyologii*. Vol. 31. N 4. Pp. 596-605 (in Russian).
- Ruban G.I., and Akimova N.V. (1993). Ecological Characteristics of Siberian Sturgeon, *Acipenser baeri*, from the Kolyma River. *Journal of Ichthyology (formerly Problems of Ichthyology) Scripta Technica*, New York. V. 33. N 5. Pp. 66-80.
- Ruban G.I., and Panaiotidi, A.I. (1994). [Comparative morphological analysis of subspecies of the Siberian Sturgeon, *Acipenser baerii stenorrhynchus* A. Nikolosky and *Acipenser baeri chatys* Drjain (Acipenseriformes, Acipenseridae), in the Yenisey and Lena Rivers]. *J. Ichthyol.* 34:58-71 (in Russian).
- Ruban, G.I. (1996). The Siberian Sturgeon, *Acipenser baerii baerii*, Population Status of the Ob' River. *The Sturgeon Quarterly*. V.4. N1/2. Pp. 8-10.
- Ruban, G.I. (1997). Species Structure, Contemporary Distribution and Status of Siberian Sturgeon, *Acipenser baerii*. In: V. Birstein, J.R. Waldman, and W.E. Bemis (ed). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publishers, Dordrecht. Pp. 221-230.
- Ruban, G.I. (1999). [*The Siberian Sturgeon Acipenser baerii Brandt (Structure and Ecology of the Species)*]. Moscow. GEOS publishers. Pp. 235 (in Russian).
- Snyder, D. (2000). Md. Caviar Company Admits Smuggling. *The Washington Post*, 22 July 2000.
- Sokolov and Vasil'ev, V.P. (1989). *Acipenser baerii* Brandt, 1869. In: Holčík J. (ed). *The Freshwater Fishes of Europe*. Vol. I/II: General Introduction of Fishes. Acipenseriformes. Wiesbaden, AULA-Verlag. Pp. 263-284. In: *Document Doc. 10.89; Prop. 10.65*. (1997). Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Williot, P., Brun, R., Rouault, T. and Rooryck, O. (1991). Management of female spawners of the Siberian sturgeon, *Acipenser baerii* Brandt: first results. In: Williot, P. (ed.) *Acipenser*, Cemagref Publ. Bordeaux. Pp. 365-379.

Gross exports* and export quotas for *Acipenser baerii* 1998

TAXON	TERM	UNIT	Exporter	1998	Export quota 1998 (kgs)
<i>Acipenser baerii</i>	Bodies		FR	24	
<i>Acipenser baerii</i>	Eggs	kg	FR	280	
<i>Acipenser baerii</i>	Eggs	kg	RU	3	19,000
<i>Acipenser baerii</i>	Live		DE	400	
<i>Acipenser baerii</i>	Live		FR	160,000	
<i>Acipenser baerii</i>	Live		GB	550	
<i>Acipenser baerii</i>	Live		XX	200,000	
<i>Acipenser baerii</i>	Meat	kg	FR	83	
<i>Acipenser baerii</i>	Meat	kg	UY	350	
<i>Acipenser baerii</i>	Other		CH	50	

Comparative tabulation of trade in *Acipenser baerii* 1998

Year	Imports reported						Exports reported						
	Imp.	Exp.	Origin	Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
Eggs													
1998	GA	FR							100 kg	eggs	T	C	
1998	MA	FR							50 kg	eggs	T	C	
1998	US	FR		61 Kg	eggs	T	C		130 kg	eggs	T	C	
1998	GR	RU							3 kg	eggs	T	C	
	Total:			61 Kg					283 kg				
Live fish													
1998	CZ	DE		200	live	T	F						
1998	PL	DE		200	live		F		200	live	T	F	
1998	CN	FR							150,000	live	T	C	
1998	HK	FR							5,000	live	T	C	
1998	MA	FR							5,000	live	T	C	
1998	XX	GB	HU						550	live	T	C	
1998	US	XX		200,000	live	T	I						
	Total:			200,400					160,750				
Other specimens													
1998	CH	FR		24	bodies	T	C						
1998	CH	FR							83 kg	meat	T	O	
1998	US	UY		350 kg	meat	T	C						
1998	DE	CH	XX						50	other		O	

Definition: "Gross exports are the sum of all reported exports and re-exports in a particular commodity or species in a particular year or series of years; Gross trade is thus a simple measure of the total number of items recorded in international trade. However, gross trade may be an overestimate of the total number of actual specimens in trade as re-exports are not deducted from the total." (Anon, 1996).

This is particularly applicable to caviar of which shipments are often exported, re-packaged and re-exported. For instance, in the mid-1990s, Germany re-exported as much as 45% of its annual caviar imports (De Meulenaer and Raymakers, 1996).

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

The Lake Sturgeon *Acipenser fulvescens* is one of the few sturgeon species that spends its entire life cycle in freshwater. The species is widely distributed through the eastern United States of America (USA) and Canada occurring in the Great Lakes, Hudson Bay-James Bay and Mississippi watersheds. The age at first spawning ranges from 14-16 years with an average size of 114.3 cm for males and 24-26 years with an average size of 139.7 cm for females. Many populations have been fragmented by the construction of dams, and have been affected by increasing levels of pollution. However, populations in Canada are thought to be healthier than the majority of those in the USA. Commercial fisheries for the species were developed during the early 1900s, but in many cases after high initial yields, the fisheries closed after ten years or so when yields dropped drastically and fisheries were no longer viable.

The catch of *A. fulvescens* in the last quarter of the 19th century exceeded 2,000 tonnes (t) annually. Today, almost the entire catch originates from Canada where in 1997 landings of the commercial catch totalled only 223 t caught in New Brunswick and Quebec, valued at USD1.8 million, with 90% (200 t) caught in the St. Lawrence River. The Canadian recreational *A. fulvescens* fishery is very limited. It totalled 1,100 fish in Saskatchewan and Alberta in 1995. In the Great Lakes, the annual average recorded catch of the species in Lake Erie and Lake Ontario, combined for USA and Canada, dropped from 49 t per year in the 1880s to 0.32 t per year in the 1960s. The 1998 international trade in specimens of *A. fulvescens* amounted 18,167 kg of meat from wild sources exported by Canada to USA, 1,120 live fish exported by USA to Taiwan (Province of China) (50% from the wild and 50% from captive bred sources) and 9 specimens (probably adults) declared as pre-Convention stock and exported by USA to the Russian Federation and Germany. Although successful artificial breeding has been reported, information on the present status of development of commercial farming of *A. fulvescens* is currently lacking. No export quotas have been established by range States since the Appendix II listing entered into effect.

The species is recommended under Decision 11.106 for inclusion in category 2 (ii).

DISTRIBUTION AND POPULATION

The distribution of *Acipenser fulvescens* is listed in the CITES database as: Canada, USA (Anon., 2000).

A. fulvescens is classified as Vulnerable by IUCN (1996):
VU A1ad+ 2d Canada, USA.

A. fulvescens is widespread in eastern Canada and USA occurring in three major watersheds: the Great Lakes, the Hudson-James Bay and the Mississippi River (Houston, 1987). The species still inhabits much of its natural distribution in Manitoba, Ontario and Quebec, but some populations are now much reduced or extirpated e.g. lakes Winnipeg, Ontario and Erie. Some populations are isolated and fragmented and differ morphologically and genetically (Ferguson and Duckworth, 1997). In contrast, their distribution within the USA is fragmented and Wisconsin is the only state where the species is considered common.

Canada: The Canadian distribution includes rivers and lakes in the five provinces of Alberta, Saskatchewan, Manitoba, Ontario and Quebec. It is found as far west as Edmonton on the North Saskatchewan River, as far east as St. Roch de Aulinaires on the St. Lawrence River, as far north as the Seal River and as far south as Lake Erie (Ferguson and Duckworth, 1997; Houston, 1987). *A. fulvescens* occurs in larger rivers and lakes from the St. Lawrence River and Great Lakes north to Hudson Bay and west to the North and South Saskatchewan rivers (CITES Management Authority of Canada, *in litt* to TRAFFIC Europe, August 2000).

United States of America: The species is mainly confined to the Great Lakes, but also occurs in the Mississippi River drainage from the upper Mississippi River and its major tributaries to the northern border

of Arkansas. The species is listed as Endangered, Threatened or Extirpated by several states (Iodd, 1999). Unless otherwise stated, information hereunder has been provided by US Fish and Wildlife Service (USFWS) (CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000).

Alabama: *A. fulvescens* is presumed to be extirpated since there are no recent records of occurrence.

Arkansas: Although occurring in Arkansas historically, *A. fulvescens* has long been extirpated from the state.

Iowa: A small population of *A. fulvescens* is found in the Mississippi River.

Kansas: *A. fulvescens* is currently limited to the Missouri River and possibly the lower Kansas River.

Michigan: The current population is believed to be at 1% of its former size.

Ohio: *A. fulvescens* is limited in distribution and relatively rare.

Vermont: *A. fulvescens* is only found in Lake Champlain.

Wisconsin: The population of *A. fulvescens* is large enough to allow sport fishing.

HABITAT AND ECOLOGY

A. fulvescens is one of the few sturgeon species that is generally confined to freshwater. However, there are a few records of individuals being taken in brackish waters of the St. Lawrence River and the Moose River (Scott and Crossman, 1973). The usual habitat is the highly productive shoal area of larger lakes and rivers (Houston, 1987). They require swift currents and large rough substrates for spawning and embryo incubation (Ferguson and Duckworth, 1997). Diet varies according to food availability but mainly consists of benthic invertebrates and in some areas, small fish. The approximate age and size at which *A. fulvescens* become sexually mature in Lake Winnebago (USA) are 14-16 years and 114.3 cm for males, and 24-26 years and 139.7 cm for females (Priegel and Wirth, 1971, cited in Houston, 1987). In Canadian rivers and lakes, this size ranges from 76.2-97.8 cm for males and from 83.8-116.8 cm for females (Houston, 1987).

Typical size/age relationships of *A. fulvescens*

Age	6 months	5 years	15 years	50 years
Length (cm)	10 – 15	30 – 70	63 – 100	101 – 180
Weight (kg)		0.45 – 2.26	2.27 – 11.34	18.14 – 36.29

Source: MacNeill and Busch, 1994

THREATS TO SURVIVAL AND DOMESTIC USE

Loss of habitat due to dam construction on virtually all larger river systems is the main threat to this species (Houston 1987; Ferguson and Duckworth, 1997). Dams constructed for hydroelectric generation affect water levels and prevent fish from reaching the different habitats required for specific life stages. Fragmentation into isolated populations may lead to a loss of genetic variability. Pollution of almost all river systems within the species' range by industry and agriculture development has affected *A. fulvescens* and continues to do so (Houston, 1987).

A. fulvescens was an important food source for Native Americans, especially during their spring ceremonial feasts conducted near the spawning grounds (Ferguson and Duckworth, 1997). The annual average recorded catch of *A. fulvescens* in lakes Erie and Ontario, combined for USA and Canada, dropped from 49 t per year in the 1880s to 0.32t per year in the 1960s (MacNeill and Busch, 1994).

Canada: Sturgeon harvest in Canada consists of aboriginal, commercial and recreational fisheries (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000). The total annual catch of *A. fulvescens* in the last quarter of the 19th century exceeded 2,000 t (Houston, 1987). In 1995, the

reported recreational *A. fulvescens* fishery totalled 1,100 fish in Saskatchewan and Alberta, mostly caught in the North and South Saskatchewan rivers (Department of Fisheries and Ocean Canada, *in litt.* to the CITES Management Authority of Canada, 9 June 2000). In the late 1990s, the remaining commercial *A. fulvescens* fisheries were in the Provinces of New Brunswick, Ontario and Quebec. In 1997, commercial catches totalled 223 t, with 200 t (90%) caught in the St. Lawrence River (Quebec). This volume was valued at USD1.8 million (Department of Fisheries and Ocean Canada, *in litt.* to TRAFFIC North America-Canada, 21 September 2000).

Unless otherwise stated, information hereunder has been provided by the CITES Management Authority of Canada (*in litt.* to TRAFFIC Europe, August 2000).

Alberta: There is no commercial sturgeon fishery but some recreational fishing.

Manitoba: In a number of rivers, stocks have been depleted through hydroelectric development and/or harvest. In 1999, only aboriginal subsistence fishing remained in most rivers.

Ontario: In 1984 there were two areas in Lake Huron that had commercial *A. fulvescens* fisheries: North Channel and the southern ends of the Main Basin. The level of annual catches in 1997 was 5.5 t in Lake Huron and 0.38 t in Lake St. Clair. The commercial fishery in Lake St. Clair was closed in 1970 due to contaminants and was reopened in 1980. There is no fishery for *A. fulvescens* in the Lake of the Woods and the Rainy River, or commercial fishery for the species in Lake Superior and the Ottawa River.

Annual commercial catch of *A. fulvescens* (kilogrammes)

	1994	1995	1996	1997	1998	1999
Catch	8,329	4,984	5,746	5,990	6,790	6,235

Source: Department of Fisheries and Oceans Canada, *in litt.* to TRAFFIC Europe, 27 September 2000

Quebec: Commercial fishing for *A. fulvescens* in rivers and lakes totalled 208.714 t in 1997 (Department of Fisheries and Ocean Canada, *in litt.* to the CITES Management Authority of Canada, 9 June 2000), with 96% of the catch originating from the St. Lawrence River.

Annual Commercial catch data for earlier years (tonnes)

	1986	1987	1988	1989	1990	1991	1992	1993	1994
Catch	184	246	173	158	222	217	223	225	214

Source: Pierre Dumont, *in litt.* to TRAFFIC Europe, 1995

Sport fishing consists of a few hundred fish per year in the Montreal area.

Saskatchewan: The province has a commercial *A. fulvescens* catch quota of 4,500 kg/year on the lower Saskatchewan River. This quota was sustainable in the 1950s, but following the loss of habitat and over-fishing, commercial harvest declined and a self-imposed moratorium began in 1996.

United States of America: Unless otherwise stated, information hereunder has been provided by USFWS (CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000).

Iowa: *A. fulvescens* is on the state endangered species list.

Kansas: There is no legal commercial season for this species. However, anglers who catch *A. fulvescens* incidentally by legal methods are allowed to keep them. There are no records of illegal harvest.

Michigan: Since the mid-nineteenth century, exploitation and habitat degradation have resulted in substantial population declines. These declines have been caused by: overharvest, barriers to migration, sea lamprey (parasite) proliferation, physical alteration of habitat, changes in water flow and quality. Anglers may take *A. fulvescens* by hook-and-line methods and a tribal fishery on Keweenaw Bay and Portage Lake allows Native American anglers to catch the species (Mosher, 1999).

Minnesota: Reported annual anglers' catch of *A. fulvescens* (kilogrammes)

1996	1997	1998	1999	2000
100	643	3,900	2,260	4,170

Source: USFWS, CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000

Local consumption is based on angling harvest where the fishing season for *A. fulvescens* is open. The trade in *A. fulvescens* has decreased in recent years due to the lack of viable eggs from commercial fish farms in other States. There are records of illegal catch (see "Conservation Measures").

New York: Carcasses of *A. fulvescens* are brought in from other states under license. This trade supports an active processing (meat smoking) industry and market for sturgeon products in the New York City area. About 68,000 kg of *A. fulvescens* were imported into New York smoke-houses in 1999.

Ohio: There are no official records of poaching activities. There are no known aquaculture facilities for *A. fulvescens*.

South Dakota: Harvest of all sturgeon species, by both sport and commercial methods, is prohibited in the state. There are no records of poaching activity.

Vermont: The fishery for *A. fulvescens* was closed in 1967.

Wisconsin: The harvest is limited to two separate sport fishing seasons.

Catch of *A. fulvescens* (number of fish)

	1998	1999	2000
Spearing season	2,051	1,484	2,517
Hook and line season	314	347	

Source: USFWS, CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000

INTERNATIONAL TRADE

A. fulvescens was listed on CITES Appendix II in 1975, delisted in 1983 and relisted in 1997, entering into force on 1 April 1998.

Gross exports and comparative tabulation of trade in *A. fulvescens* for 1998 are given in the Appendix.

According to 1978-1984 CITES data, exports of *A. fulvescens* were restricted to meat with an average 6,416 kg/year exported during this seven year period. For 1998, the most important international trade in wild specimens of *A. fulvescens* is still represented by meat; exports from Canada to USA totalled 18,170 kg. This volume is the largest quantity of sturgeon meat trade reported in 1998 CITES Annual Reports for all Acipenseriformes.

A total of 1,120 live specimens (probably juvenile fish) were exported from USA to Taiwan (Province of China), and nine specimens from a pre-Convention stock (probably adults to be used as broodstock in captive breeding facilities) were exported from USA to Germany and the Russian Federation.

CONSERVATION MEASURES

Tagging studies are carried out widely. Habitat reconstruction has been undertaken in some states of USA and appears to have been successful. The protection of remaining habitat is a priority in Canada (Houston, 1987).

Canada: The species is not listed as a species at risk by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). The species is protected and managed in Canada under the *Federal Fisheries Act* in each province of occurrence. Regulations differ between provinces and are revised annually. *A. fulvescens* is listed under provincial legislation as Threatened in Alberta, Manitoba, New

Brunswick, Newfoundland, Ontario, Quebec and Saskatchewan (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000).

Alberta: A Lake Sturgeon Management Plan has been implemented on the South and North Saskatchewan River since 1997. A zero catch limit is in effect for *A. fulvescens* in the North Saskatchewan River and in the South River, a limit of one sturgeon of 130 cm in total length (through a tag system) is in effect. There is a closed season during the spawning period from 1 April to 15 June. There is no aboriginal fishery (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000).

Manitoba: The possession limit for sturgeon was reduced to zero throughout the province in 1995. The remaining few commercial fisheries closed down during the 1990s, but “catch and release” angling is still allowed in addition to aboriginal subsistence fishing in most rivers. *A. fulvescens* population management has been initiated in a couple of rivers, with the involvement of First Nations communities (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000). In addition, sturgeon cultures to rehabilitate populations in the Nelson, Winnipeg and Saskatchewan rivers have begun (CITES Management Authority of Canada *in litt.* to CITES Secretariat, November 2000).

Ontario: In 1984, catch quotas were allocated based upon annual performance by individual fishers from 1978 to 1982. Quotas have not been adjusted although they have never been achieved and may therefore have been set too high. In Lake St. Clair the total annual catch quota was set at 1,500 kg and remains unchanged since 1985. The current total catch quota for Lake Huron is 13,024 kg, although the total 1997 catch was only 5,471 kg, which has been the approximate level of the annual catch since the early 1980s. (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000). All commercial fishers in Ontario are required to fill out daily catch report, however, in the north many First Nations do not complete these daily catch records (CITES Management Authority of Canada *in litt.* to CITES Secretariat, November 2000). Since 1995, a Lake Sturgeon Assessment programme has been implemented, including population abundance estimation and tagging to identify movements and verify ages. A management plan for the species will be drafted after sufficient data are collected. In the meantime, quotas will not be increased, but may be decreased pending preliminary data analysis (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000).

Catch quotas set for *A. fulvescens* in Lake Huron (kilogrammes)

1991		1992		1993		1994		1995		1996		1997	
Quota	Catch	Quota	Catch	Quota	Catch	Quota	Catch	Quota	Catch	Quota	Catch	Quota	Catch
11,121	5,335	11,667	4,632	11,843	4,656	13,124	4,074	13,165	3,848	13,124	4,370	13,024	5,471

Source: Department of Fisheries and Oceans Canada, *in litt.* to TRAFFIC Europe, 27 September 2000

Quebec: Starting just after the spawning season, fish may be harvested with gill nets (19-20.3 cm mesh) from 14 June to 31 October. The minimum size limit is 45 cm (the legal length is defined as: the distance from the posterior edge of the branchial slit to the posterior joint of the dorsal fin). In sport fishing, only one fish per day can be harvested or kept at any one time. Commercial catch must be tagged as soon as fish are caught and remain tagged until they are processed for human consumption. Issuing a specified number of tags to each commercial fisherman enforces catch quotas (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000).

Saskatchewan: A self-imposed moratorium on commercial catch on the lower Saskatchewan River has been in force since 1996. Angling of the species has not been allowed in Saskatchewan since the beginning of 1999. A multi-agency study of habitat, fish migration and abundance is being carried out (CITES Management Authority of Canada, *in litt.* to TRAFFIC Europe, August 2000).

United States of America: Unless otherwise stated, information hereunder has been provided by USFWS (CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000).

Iowa: *A. fulvescens* has been on the state endangered species list for many years.

Missouri: A total of 164,000 *A. fulvescens* fry and fertilised eggs were released in the Mississippi and Missouri rivers between 1983 and 2000. A few adult and pre-adults have been released occasionally.

Michigan: *A. fulvescens* was listed as a state threatened species in 1994. A rehabilitation strategy has been prepared and regulations restrict the sport fishery in season, number as well as size of fish and fishing gear (Mosher, 1999).

Minnesota: *A. fulvescens* and Shovelnose Sturgeon *Scaphirhynchus platyrhynchus* as well as American Paddlefish *Polyodon spathula* may not be taken, bought, sold, transported or possessed unless authorised by the commissioner. Infractions to the following regulations were recorded: no license (8 events), no license in possession (1), closed area (4), closed season (3), illegal equipment (1), undersized fish of less than 45 inches (2) and oversized fish (1). One warning was given for import/transport of sturgeon hybrid without a permit. Farming of *A. fulvescens* is being developed. A cooperative project for restocking *A. fulvescens* is being undertaken using wild broodstock and artificial breeding, or by the transfer of wild juveniles.

New York: The species is considered a threatened species. All harvest or possession is prohibited except under license or permit. Licenses are available for the import of *A. fulvescens* carcasses and the processing of their products. A recovery plan has been conducted since 1994. *A. fulvescens* fingerlings have been reared and released in a selection of target restoration waters since 1995. Survival of released sturgeon has been high and populations are recovering. Eggs for this program are collected from wild stocks in the St. Lawrence River. No captive broodstock is maintained and there are no recorded captive breeding operations in the State.

North Dakota: Although Minnesota is undertaking an *A. fulvescens* re-introduction programme within the Red River drainage, North Dakota is not an active partner of this programme.

Ohio: The species has been listed as an Ohio endangered species since 1974. Ohio is participating in research and outreach programmes for the species. The state does not have a commercial or sport catch/quota for *A. fulvescens*.

South Dakota: Harvest of any specimen by sport and commercial fishermen and selling or trading flesh and/or body parts of sturgeon (including *A. fulvescens*) by anglers is prohibited.

Vermont: The fishery was closed in 1967 and *A. fulvescens* was placed on the state endangered species list in 1987. It is illegal to take or possess *A. fulvescens* without a permit. The USFWS fish culture facility at Pittsford (Vermont) reared the St. Lawrence River strain of the species for a restoration programme on the St. Lawrence River and its tributaries in New York State. The fry were not released in Vermont.

Wisconsin: Fisheries management measures have been adopted for spear and hook-and-line fisheries. For the hook-and-line fishery, there is 127-178 cm minimum length limit depending on the location. The fishing season starts on the first Saturday in September and runs through to 15 October. The catch limit is one fish per angler per season, and anglers must apply for a free tag.

For the spear fishery, there is a 91cm minimum length catch limit. Spear fishers must purchase a USD10 sturgeon spear tag, with the season starting on the first Saturday in February until the quota is reached. The Lake Winnebago spear fishery is regulated under an annual catch quota system (in 2000, 400 adult females, 400 juvenile females and 2,150 males). Once 80% of the quota is reached, the season is closed at the end of the following day. Since the early 1980s, volunteer sturgeon patrols have observed spawning sites for 24-hour shifts each spring. Commercial breeding of *A. fulvescens* is prohibited in Wisconsin. The hatchery of the Wisconsin Department of Natural Resources propagated sturgeon for restocking purposes. Fish can only be restocked into the same basin that the eggs were collected; no inter-basin transfers are allowed.

CAPTIVE BREEDING

Canada: Aquaculture of *A. fulvescens* has been reported, but no results on the volumes of specimens produced were available from the Department of Fisheries and Oceans Canada (TRAFFIC North America-Canada, *in litt.* to TRAFFIC Europe, 21 September 2000).

United States of America: No records of commercial captive breeding were recorded in Arizona, Delaware, Florida, Louisiana, New York, Ohio, Oklahoma, South Dakota, Tennessee, Utah or Wyoming. Sturgeon are in a preliminary stage of culture in Georgia, Hawaii and Virginia (USFWS, CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000).

Minnesota: Three licensed aquaculture farms have stocks of *A. fulvescens*. The fish were either obtained from artificial breeding of wild specimens or were purchased from commercial operations in Kentucky. None of the facilities have yet produced viable eggs.

Wisconsin: Commercial rearing is prohibited, but State-run hatcheries exist.

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database, but according to information provided by range States, and significant production of *A. fulvescens* in commercial farms has not been reported.

REFERENCES

- Anon. (2000). <http://www.cites.org/eng/dbase/fauna->
- Ferguson, M.M. and Duckworth, G.A. (1997). The status and distribution of Lake Sturgeon, *Acipenser fulvescens*, in the Canadian provinces of Manitoba, Ontario and Quebec: a genetic perspective. In: Birstein, V.J., Waldman, J.R., and Bemis, W.E. Sturgeon biodiversity and conservation. Kluwer Academic Publishers, Dordrecht. *Environment Biology of Fishes*. 48:299-309.
- Houston, J.J. (1987). Status of the Lake Sturgeon, *Acipenser fulvescens*, in Canada. *Canadian Field-Naturalist* 101(2):171-185.
- IUCN. (1996). *The 1996 Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- MacNeill, D. and Busch, W.D. (1994). The Biology, History and Management of the Lake Sturgeon in the Lower Great Lakes. *Sea Grant*, Cornell Cooperative Extension, State University of New York, Sportfishing Fact Sheet, January 1994.
- Mosher, T.D. (1999). Sturgeon and paddlefish sportfishing in North America. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 51-66
- Scott, W.B. and Crossman, E.J. (1973). Freshwater Fishes of Canada. *Bull. Fish. Res. Board Can.* 184:1-966.
- Todd, R.M. (1999). Sturgeon and Paddlefish commercial fishery in North America. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 42-50.

Gross exports of *Acipenser fulvescens* 1998

TAXON	TERM	UNIT	Exporter	1998
<i>Acipenser fulvescens</i>	live		US	1,120
<i>Acipenser fulvescens</i>	meat*		CA	5,950
<i>Acipenser fulvescens</i>	meat	kg	CA	12,220
<i>Acipenser fulvescens</i>	meat	kg	US	279
<i>Acipenser fulvescens</i>	specimens		US	9

Comparative Tabulation of trade in *Acipenser fulvescens* 1975-1983 and 1998

Year	Imp.	Exp.	Origin	Imports reported			Exports reported				
				Quantity	Unit	Term	P	S	Quantity	Unit	Term
Meat											
1998	US	CA		12,220 kg	meat	T	W				
1998 *	US	CA		5,950 kg*	meat	T	W				
1998	RU	US	CA					279 kg	meat	T	W
TOTAL				18,170 kg				279 kg			
Other specimens											
1998	TW	US						560	live	T	C
1998	TW	US						560	live	T	W
1998	DE	US		6 specimens		S	O	6	specimen		O
1998	RU	US						3	specimens		O
Earlier years											
1978	US	CA		270 kg	meat			(Listed from 1975 to 1983)			
1979	US	CA		1,415 kg	meat						
1980	US	CA		3,622 kg	meat						
1981	US	CA		9,705 kg	meat						
1982	US	CA		11,613 kg	meat						
1983	US	CA		15,862 kg	meat						
1984	US	CA		2,424 kg	meat			(COP 4: De-listing of the species)			
Average:				6,416 kg/year							

* The Unit is "kg" (US CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000). The total meat of *A. fulvescens* exported from Canada in 1998 was therefore 18,170 kg.

Acipenser gueldenstaedtii Brandt, 1833 Russian Sturgeon
Ossetra

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

The Russian Sturgeon *Acipenser gueldenstaedtii* may live to 48 years weighing around 100 kg. The species matures sexually between 8-16 years and most female spawners in the wild are aged 13-23 years. Spawning occurs every 2-3 years. *A. gueldenstaedtii* has a wide distribution, occurring in the Caspian Sea, Black Sea and the Sea of Azov, and originally in many of the tributaries of these seas. However the construction of dams on virtually all these rivers has significantly reduced the spawning area. The Ural River is one of the few major spawning rivers that has not been dammed. In addition to loss of habitat, *A. gueldenstaedtii* has been subject to overfishing and egg-production has been disrupted due to environmental pollution. In the early 1990s, around 30% of the population in the Caspian Sea and almost all the stock in the Sea of Azov originated from re-stocking programmes. However, the restocking of the Volga River by Russian hatcheries decreased from 1991-1995 levels by almost one third during 1996-1998. In contrast, the number of fingerlings released by Azerbaijan, Bulgaria and Iran has generally increased since 1995. Only Iran monitors the success of their restocking programme. The species is widely bred in captivity to produce meat and fry for both domestic and international trade. There is no record of caviar production in captive breeding facilities.

When commercial sturgeon catch in the Caspian peaked in 1997, *A. gueldenstaedtii* comprised almost 80% of the total catch and still provides a major portion of the global caviar production. Since then, recorded annual catches of the species have decreased in all range States. In the Caspian Sea they decreased from 21,550 tonnes (t) in 1977 to less than 1,000 t in 1999, and in the Danube River annual catches decreased from 24 t in the 1950s to the early 1980s, to less than 11 t in 1999. Caspian Sea annual catch quotas are set by the Intergovernmental Commission for Caspian Biological Resources, composed of representatives of all Caspian range States. The catch quotas of Azerbaijan, Kazakhstan and the Ukraine have all recently been decreased. Domestic markets of sturgeon products are unknown in the Russian Federation, and Iran estimated its domestic consumption to be 5% of the country's total caviar production and 65% of its sturgeon meat production. In 1998, range State exports of *A. gueldenstaedtii* caviar of wild origin totalled 95.3 t, with the Russian Federation (46 t) and Iran (40 t) being the main exporters. This may represent a minimum catch (males and females) of 1,361-1,588 t, using a standard rate of 6-7% in weight of caviar of the total catch. Iran apparently exceeded its caviar export quota of 10 t (but caviar exports of *A. gueldenstaedtii* and *A. persicus* may have been combined in the Iranian CITES Annual Report for 1998). The total caviar export quota for 2000 has decreased from the 1999 quota from 119.5 t to 95.9 t, but increased significantly for meat, from 53.5 t to 204 t.

The species is recommended under Decision 11.106 for inclusion in category 1 (i).

SPECIES IDENTIFICATION

Vlasenko *et al.* (1989) contend that *A. gueldenstaedtii* is not distinguishable from *A. persicus* Persian Sturgeon, where their ranges overlap in the southern and southeastern Caspian. Indeed, the two species are combined in the catch statistics of Azerbaijan, and are combined for Iran's 2000 caviar export quota. However, Dr M. Pourkazemi (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) notes that *A. gueldenstaedtii* can be distinguished from *A. persicus* in the Iranian coasts on the basis of 22 morphological differences (Nazari Chari, 1993, cited in Dr M. Pourkazemi *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). These two species can also be differentiated on the basis of immuno-biochemical characteristics.

Taxonomic confusion also exists concerning the species to which the subspecies *colchicus* belongs; *A. gueldenstaedtii colchicus* (e.g. Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, 2 November 2000), or *A. persicus colchicus* (e.g. Zarkua and Tsuladze, 1999).

DISTRIBUTION AND POPULATION

The CITES database lists the current distribution of *A. gueldenstaedtii* as: Azerbaijan, Bulgaria, Georgia, Iran (Islamic Republic of), Kazakhstan, Romania, Russian Federation, Turkey, Turkmenistan, Ukraine (Anon., 2000a).

A. gueldenstaedtii is classified as Endangered by IUCN (1996):

EN A2d Azerbaijan, Bulgaria, Hungary, Iran, Kazakhstan, Moldova, Romania, Russia, Turkey?, Turkmenistan, Ukraine, Yugoslavia

Caspian Sea stock: EN A2d Azerbaijan, Iran, Kazakhstan, Russia, Turkmenistan

Sea of Azov stock: EN A1acde+ 2d Russia [Mediterranean and Black Sea]

Black Sea stock: EN A1acde Hungary, Romania, Ukraine, Yugoslavia [Mediterranean and Black Sea].

Two forms of the species exist. The anadromous form of this sturgeon inhabits the Caspian Sea, the Black Sea and the Sea of Azov (Vlasenko *et al.*, 1989). Sturgeon inhabiting the Caspian Sea migrate mainly to the Volga and Ural rivers for spawning. Spawning populations in rivers other than the Volga are now minimal (Khodorevskaya *et al.*, 1997; Levin, 1997; Vlasenko, 1990; Vlasenko *et al.*, 1989). In the Caspian Basin, *A. gueldenstaedtii* is estimated to have lost approximately 70% of its spawning grounds (Barranikova *et al.*, 1995). The Caspian and Black Sea populations contain both a spring and a winter spawning race (Vlasenko *et al.*, 1989). A non-migratory form also exists in the Danube and Volga River basins, and formerly in the Ural River (Hensel and Holèik, 1997; Holèik, 1995; Manea, 1966; Vlasenko *et al.*, 1989), although Birstein (1993) believes the non-migratory form to be extinct.

Azerbaijan: *A. gueldenstaedtii* occurs in the Kura River, up to Vavarin reservoir, the Lenkoranka River (Azerbaijan, Georgia) and the Astara River (Azerbaijan, Iran) (Vlasenko *et al.*, 1989). In waters of the mid-western Caspian Sea, along the Azeri shore, *A. gueldenstaedtii* swims in coastal areas in spring and summer and migrates towards deep waters of the open sea as the temperature drops (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Bulgaria: A portion of the Black Sea population migrates into the Danube River to spawn. The breeding seasons are in spring (early-April when the first catch of the year are recorded) and in autumn (end-September) (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). *A. gueldenstaedtii* also occurs along the Black Sea coast. However, it is considered "rare" (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC, September 2000) due to overfishing and reduced reproductive capacity (Karapetkova *et al.*, 1995).

Georgia: The southeast coast of the Black Sea is an important feeding and wintering area for sturgeon (*Huso huso*, *A. stellatus*, *A. nudiventris*, *A. persicus colchicus* and *A. sturio*) that migrate upstream to a number of rivers for spawning. The main rivers involved are the Supsa, Inguri, Chorokhi and particularly the Rioni. *A. persicus colchicus* is still the most abundant sturgeon species in Georgian waters (Zarkua and Tsuladze, 1999). *A. gueldenstaedtii* has also been recorded from the Lenkoranka River (Azerbaijan, Georgia) (Vlasenko *et al.*, 1989).

Hungary: The species is believed by some to be extinct in Hungary (Pintér, 1991). However, it may still occur in the Tisza River (Tiszafüred) where the last catch was recorded in 1980 (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Dr M. Pourkazemi (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) notes that the Sefidrud River and Gorganrud River on the Iranian coasts of the Caspian Sea are used for spawning (Laluyee, 1996; Ramin, 1998, cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). However, Dr J. Holèik (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) contends that access to the spawning grounds in the Sefidrud has been prevented by the Mangil Dam construction, water extraction and pollution. According to Dr M. Pourkazemi (*in litt.* to IUCN/SSC Wildlife Trade Programme, 25 September 2000), among the Mangil Dam and other dams constructed on the Sefidrud River, there is a remaining distance of 45 km from the dam closest to the estuary, and sturgeon are still observed entering the river to spawn.

Kazakhstan: The species enters the Ural River for spawning, one of the few spawning rivers that is not dammed (Vlasenko *et al.*, 1989).

Moldova: The Moldovan range on the Danube River is 811 m and sturgeon also occur in the Prut River, a tributary of the Danube River (junction at km 137th) (Anon., 2000c).

Romania: *A. gueldenstaedtii* is recorded from the Danube River and the coast of the Black Sea. The population density and catches of *A. gueldenstaedtii* dropped dramatically after the damming of the Danube by Djerdap I and II power stations (Bacalbasa-Dobrovici, 1997). The species was formerly caught in the lower reaches of the Prut, Jiu and Mures rivers (Hensel and Holèik, 1997).

Russian Federation: In the Caspian region, the population entering the Volga River to spawn in 1996 was estimated at 120,000-150,000 individuals (Levin, 1997), a decrease from 500,000–600,000 individuals in previous years. Levin (1997) estimated the age range of the population majority to be between 8-25 years, with individuals older than 28 years being extremely rare.

In the Sea of Azov, some natural spawning grounds are no longer accessible due to the construction of dams on the Don and Kuban rivers (Anon., 2000b). In the mid-1980s, Volovik *et al.* (1993) estimated the total biomass of *A. gueldenstaedtii* stock to be 44.8 t. In 1990, 55,000 sturgeon were found dead on the shores of the Sea of Azov. No population estimates have been undertaken since.

Slovakia: The winter race was present in the Danube until 1967, when it is likely that the last specimen was caught (Holèik, 1995). Three or four individuals of the resident form are caught annually in the Danube River (Dr J. Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). The species' future is threatened and the species is critically endangered (Holèik, 1989).

Turkey: The Coruh River is used for spawning. Unidentified sturgeon species have been reported in small numbers in the Sakarya and Mert rivers (Edwards and Doroshov, 1989). In the late 1980s, local fishermen estimated that only five to six gravid female sturgeon could be taken from the Yesilirmak to supply a hatchery and up to 20 from the Kizilirmak (Edwards and Doroshov, 1989).

Ukraine: From the Sea of Azov, *A. gueldenstaedtii* migrates to the Don River, where it spawns in its lower segment, downstream of the Tsimlyansk Hydropower Station. According to Pavlov (1980), *A. gueldenstaedtii* is also occasionally known from the Dniester, Dnieper and Bug rivers.

Former Yugoslavia: The population density has dramatically decreased in the Danube since the construction of the Djerdap I and II power stations. There are not enough adults to maintain fry production in the Kladow hatchery (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC, September 2000). It has been classified as Endangered by Simonovic and Nikolic (1996).

Introduced populations: In the Baltic Sea river estuaries, east of Rostock (Germany), and further upstream, *A. gueldenstaedtii* and hybrid sturgeon specimens have been caught. They probably originate from upstream aquaculture farms and/or private aquaria (specimens that have grown too large to be kept in the aquarium) (Jörn Gessner, Scientist at the Institute of Freshwater Ecology and Inland Fisheries, pers. comm. to TRAFFIC Europe, 15 March 2000).

HABITAT AND ECOLOGY

A. gueldenstaedtii reach 2.2-2.4 m in length and 65-115 kg in weight. In the past, specimens were known to reach an age of 48 years; life expectancy has now been reduced to 38 years due to overfishing (Vlasenko *et al.*, 1989). The species matures sexually between 8-16 years and most female spawners in the wild are aged 13-23 years. The average size of females at first spawning is 1.2 m and 9 kg and spawning occurs every 2-3 years (Hochleithner and Gessner, 1999). The diet alters according to food availability but mainly consists of benthic molluscs, larvae and in some areas, small fish. Migration to riverine spawning grounds occurs during two periods: from early spring to late autumn, peaking in summer; and during the winter, when fish hibernate in the rivers and reproduce the following year. Juveniles migrate downstream to feeding grounds. In the Caspian Sea the main feeding area is in the

north, but a winter migration to the central and southern areas of the Caspian has been observed (Barannikova *et al.*, 1995).

Based on catches and caviar production in the late 1970s, the rate in weight of caviar/total catch was estimated at 7% for the three main commercial species of the northern Caspian Sea, *H. huso*, *A. gueldenstaedtii* and *A. stellatus* (Doroshov and Binkowski, 1985, cited in Williot and Bourguignon, 1991).

Studies indicate that sexual maturity of *A. gueldenstaedtii* in Iranian waters differ to that given by Levin (1997); an average age of 15 years in females (min. 9 and max. 19 years) and an average age of 14 years in males (min. 8 and max. 17 years) (International Sturgeon Research Institute, 1998, cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Until 1993, the mouth of the Ural River was dredged annually to allow vessels to travel upstream to Atyrau (former Gur'yev) harbour. Since 1993, this annual management of the river channel has not been carried out and is an obstacle to sturgeon migration (Anon., 2000c).

THREATS TO SURVIVAL AND DOMESTIC USE

In the Caspian Sea, construction of hydroelectric power stations and dams on most of the rivers that enter the sea has reduced available spawning habitat by up to 70% of that available in the 1950s. Winter migration has been more severely impacted by this construction than the summer migration because the distance travelled upstream in winter is generally greater (Barannikova *et al.*, 1995). The only river with an unregulated flow is the Ural. Poor environmental conditions during 1986 to 1990 have caused degeneration of the female reproductive system in fish caught in the lower Volga River in 1990 (Shagaeva *et al.*, 1993).

In the Black Sea and Sea of Azov Basins, water flow has been altered by construction and irrigation in virtually all rivers entering these seas. Pollution and eutrophication of coastal waters has caused a mass death of fish. Food sources have been depleted by an introduced ctenophore *Mnemiopsis leydyi*. However, it has recently been reported that a second invasive ctenophore *Beroe ovata* in the Black Sea may prey on the first and hence restore the ecological balance (Schrope, 2000).

High levels of pollution have posed a serious threat to sturgeons in both the Caspian and Black Sea basins. Studies have shown that high levels of toxins, from oil and other industrial waste, have altered hormonal balance, disturbed metabolism and increased the number of hermaphroditic fish. However, since the break up of the Soviet Union, heavy industry and resulting discharges have decreased allowing water quality to improve (Levin, 1997).

Caviar from *A. gueldenstaedtii* is commercially known as "Ossetra," as is caviar from several other species such as Ship Sturgeon *A. nudiiventris* and Amur Sturgeon *A. schrenckii*, and is the second most highly sought-after caviar, after that from Beluga *H. huso*. The commercial sturgeon catch in the Caspian Sea basin peaked at 27,300 t in 1977, with *A. gueldenstaedtii* being the most abundant species (79%, 21,550 t), followed by *A. stellatus* (17%, 4,650 t) (Levin, 1997), with the remaining 4% (0.11 t) being shared between *H. huso* and *A. nudiiventris*. The catch of *A. gueldenstaedtii* dropped to 16,300 t in 1978 (Khodorevskaya *et al.*, 1997) and to approximately 965 t in 1997. In the middle Danube River, the average annual catch of the species from 1958 to 1981 was 24.9 t (Hensel and Holèik, 1997). The recorded landings from the lower Danube River in Romania and Bulgaria did not exceed 11 t in 1999.

In the former Soviet Union and Iran, strict legislation regarding sturgeon fisheries had been in effect for decades until the emergence of three new independent States in the Caspian Sea Basin in 1992 (Azerbaijan, Kazakhstan and Turkmenistan) (De Meulenaer and Raymakers, 1996). Current information suggests that amendments of previous regulations and/or adoption of new legislation are now in effect in most new range States in the Caspian Sea as well as in Black Sea (including the Danube River) and Sea of Azov basins. In the latter, Bulgaria, Romania and Ukraine are the most important sturgeon fishing nations. However, enforcement measures appear to be lacking and numerous experts as well as government officials have reported an increasing pressure of illegal fishing practices and criminal activities surrounding the caviar trade in most of

the range (Anon., 2000b; Dobbs, 1992; Doward, 2000; Evtouchenko, 1997; King, 1998; Ward, 2000a; 2000b)

Azerbaijan: Waters of this region represent important winter feeding grounds, but lakes of deposited oil and pollutants lie on the shore lines of the Caspian and the number of these polluting the sea is expected to increase as the sea level rises (Dumont, 1995). Along the northern and southern coast of Azerbaijan, within the 200-nautical mile Exclusive Economic Zone, *A. gueldenstaedtii* is caught as by-catch in stationary nets and *kilka* (Clupeidae) fishing devices (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000). Young and adult fish are caught by coastal fisheries at the time of their migration, when the water temperatures change. Young fish make up to 17.7% of the by-catch.

Species composition of the catch and by-catch of sturgeon species in the late 1990s

	<i>A. persicus</i> & <i>A. stellatus</i>	<i>H. huso</i>	<i>A. nudiventris</i>
Catch	63.47%	30.77%	5.76%
By-catch	78.50%	16.80%	3.70%

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Annual recorded landings (catch and by-catch) of *A. gueldenstaedtii* and *A. persicus* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<i>A. gueldenstaedtii</i>	26.00	40.20	37.00	72.60	100.30	101.20	51.30	79.20	48.60	55.00
<i>A. persicus</i>										
Total	74.00	88.20	111.00	240.10	201.20	162.50	120.30	128.60	82.40	108.90

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

In the late 1990s, for the first time in the history of *kilka* fishery, the by-catch of *A. gueldenstaedtii* has dropped to almost zero. For instance 364 specimens were caught in 1992, while incidental catches were reduced to few anecdotal specimens in more recent years.

Bulgaria: Wild populations of sturgeon have been threatened by the decrease in number of spawning grounds on the Danube River as a result of human development, particularly hydroelectric facilities. The long lasting unregulated fisheries have disrupted the age composition of the populations due to selective catch of sexually mature fish and to by-catch of juveniles as they migrate to the sea. Water pollution has also influenced the metabolism of the fish (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Annual catch of *A. gueldenstaedtii* (tonnes)

	1995	1996	1997	1998	1999
<i>A. gueldenstaedtii</i>	0.9	2.4	5.4	7.5	6.0
Total sturgeon catch	14.7	32.5	49.8	55.9	50.5
n° of licenses issued	0	0	122	155	169

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Domestic consumption of caviar in Bulgaria is limited to 7-10% of the total caviar production of Bulgaria. Bulgaria's entire sturgeon meat production is consumed within the country.

Georgia: During the fishing seasons, particularly coastal anchovy fisheries, numerous sturgeon juveniles are caught as by-catch. The estimated catch of all species of sturgeon declined from 77,000 individuals in 1977, to 57,000 in 1980 and 27,000 in 1986. During the 1990s, illegal fishing in Georgia expanded rapidly due to poor control (Zarkua and Tsuladze, 1999).

Hungary: Threats to *A. gueldenstaedtii* are not documented due to the general lack of records. The species is protected, and domestic utilisation is prohibited (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Spawning grounds of the Sefidrud River have been lost due to the construction of the Mangil Dam in 1961 (Dr J.Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). For the past 5

years, the Iranian domestic market of caviar and sturgeon meat is estimated at 5% and 65% of the country's production respectively (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Annual catch of *A. gueldenstaedtii* and *A. persicus* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. persicus</i>	538.3	601.1	559.6	415.0	408.9	386.5	488.3	448.2	547.3	439.9	153.6
<i>A. gueldenstaedtii</i>	384.7	422.7	363.1	212.7	142.1	114.3	130.9	86.3	75.3	56.7	11.2
Total	2,296.3	2,315.1	2,058.0	1,462.1	1,380.9	1,216.3	1,310.0	1,043.5	1,127.9	923.3	182.4

* Catch in spring 2000 only.

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

Production of *A. gueldenstaedtii* and *A. persicus* in 1992 and 1999 (tonnes)

	1992				1999			
	Meat	% *	Caviar	% *	Meat	% *	Caviar	% *
<i>A. persicus</i>	451.9	28.2	62.6	23.9	355.5	48.7	50.8	51.2
<i>A. gueldenstaedtii</i>	279.7	17.4	38.6	14.7	43.6	6.1	5.8	6.0
Total	1,604.2		262.3		730.2		99.3	

* Percentages relate to the share of *A. gueldenstaedtii* products of the total Iranian sturgeon fisheries.

Source: Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000

The above tables indicate that the total tonnage of meat and caviar of *A. gueldenstaedtii* produced in Iran comprised 87% of the country's annual catch in weight in 1992 and 1999. This rate was 92% for *A. persicus*.

Note on *A. persicus*: The best results of the restocking programme of five sturgeon species were obtained for *A. persicus*. This species is considered to be endemic to the southern Caspian Sea Basin and rarely migrates to the central and northern parts of the sea. The results of the monitoring of catch and stocks of *A. persicus* suggest that the wild population does not follow the same trends as the other commercially caught sturgeon. The status of the population appears to be improving while the stocks of *A. gueldenstaedtii*, *A. stellatus*, *A. nudiiventris* and *H. huso* seem to be decreasing. In the near future, the status of the stock of *A. persicus* is expected to be larger than that of other species. In the past, *A. persicus* comprised 20% of the annual Iranian sturgeon catch, while at present, it comprises more than 55-60%. Whilst the status of *A. persicus* appears to be improving in Iranian waters and comprises over 50% of the Iranian catch, the status of *A. gueldenstaedtii* continues to decline (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Kazakhstan: Oil fields may affect spawning and feeding grounds in the northern area of the Caspian (Sagers, 1994). The central area is threatened by radioactive contamination from a nuclear reactor (Dumont, 1995).

Annual catch of *A. gueldenstaedtii* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. gueldenstaedtii</i>	123.0	154.0	136.0	115.0	47.0	31.6	42.1	39.5	135.0	60.0	34.6
Total	1,963.0	1,767.0	1,437.3	1,109.0	556.0	573.0	367.8	389.5	490.0	279.0	230.5

* Spring catch

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Romania: Threats include: overfishing, poaching, pollution and damming of the Danube by Djerdap I and Djerdap II (Bacalbasa-Dobrovici, 1997).

Annual catch of *A. gueldenstaedtii* (kilogrammes)

	1991	1992	1993	1994	1995	1996	1997	1998	1999
Catch	748	639	398	2,528	2,820	3,273	4,382	1,732	4,314

Source: Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, 5 September 2000

These data include catches in the Danube River and Delta as well as in the Black Sea. From 1991 to 1999, approximately 17% of the total Romanian sturgeon catch comprised *A. gueldenstaedtii*.

Russian Federation: Caspian region: The Volgograd Dam has reduced spawning grounds in the main spawning river, the Volga, to 430 ha from 3,600 ha. Similarly, available spawning grounds in the Terek and Sulak rivers have been reduced to 132 ha and 202 ha respectively. The flow of the Ural River remains unregulated and provides a natural spawning area of 1,400 ha.

Black Sea and Sea of Azov region: Flow regulation of the Kuban River has led to the loss of approximately 140,000 ha of breeding ground, and the Tsymlyansk Reservoir construction on the Don River in 1952 has removed approximately 68,000 ha of spawning ground (Volovik *et al.*, 1993). Worsening environmental conditions in this area caused the death of 55,000 fish in 1990.

Annual catch of *A. gueldenstaedtii* in the Caspian Sea (tonnes)

	1992	1993	1994	1995	1996	1997
Catch	4,446	2,259	1,490	1,113	510	759

Source: Caspian Fisheries Research Institute (KaspNIRKh), cited in Anon., 2000b

From 1992 to 1997, approximately 50% of the total sturgeon catch of the Russian Federation in the Caspian Sea comprised *A. gueldenstaedtii* (Anon., 2000b).

The level of domestic consumption of sturgeon products in the Russian Federation is unknown, but is said to be the main outlet of illegal sturgeon fisheries (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). Based on the number of sets of illegal fishing equipment recorded by enforcement agencies of the Caspian Sea and the Sea of Azov, the illegal sturgeon catch was evaluated at 6 to 10 times the legal catch (Anon., 2000b). In 1997, the estimated volume of the Moscow sturgeon meat market was three times the annual catch quotas for all sturgeon species (Anon., 1998).

Slovakia: No catches have been reported since the construction of the Gabčíkovo River dam system in 1992 (Dr J. Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, 2000). The species future is threatened and the species is critically endangered (Holèik, 1989).

Turkey: From 1969 to 1979, Turkish annual sturgeon landings ranged from 310 t to 10 t, with an annual average of 112 t (Edwards and Doroshov, 1989). In 1989 in Samsun, a Turkish landing site on the Black Sea, one caviar processor claimed that his business had dropped from 8 t per year of caviar between the 1940s and the 1970s, to 200 kg per year by the end of the 1980s. In 1989, sturgeon were caught as by-catch by trawlers that operated legally along the Turkish coast at a distance greater than three nautical miles from the shore, but also illegally within the three nautical mile limit. This represented the main fishing pressure on sturgeon. In April 1989, *A. stellatus*, *A. gueldenstaedtii*, *H. huso* and *A. sturio* were observed in Samsun. The investigators were told that *A. nudiventris* was also landed in Samsun (Edwards and Doroshov, 1989).

Turkmenistan: Sturgeon fisheries and processing are a state monopoly in Turkmenistan (Anon., 2000c).

Ukraine: No information provided.

Yugoslavia: The Danube River spawning grounds are no longer accessible due to the construction of the Djerdap dams. The catch of *A. gueldenstaedtii* on the Yugoslav part of the Danube River dropped drastically before (588 km) and after (17.8 km of fishing ground from the dam to the border; see "Conservation Measures") the construction of the the Djerdap II dam in 1984. The following catches of *A. gueldenstaedtii* were reported: total annual catch of 14,636 kg in 1983 and 1,636 kg in 1985, with an annual average catch of 10,354 kg for the period 1978-1983 and 1,789 kg for the period 1985-1990 (Jankoviè, 1993 and 1996).

INTERNATIONAL TRADE

The listing of *A. gueldenstaedtii* in CITES Appendix II entered into effect on 1 April 1998. Data for this species were therefore limited to nine months of trade (April-December 1998).

Gross exports and the comparative tabulation of trade in *A. gueldenstaedtii* are given in the Appendix. According to 1998 CITES data, gross exports of *A. gueldenstaedtii* caviar from range States totalled 95,392 kg, the second largest volume of all sturgeon species after *A. stellatus*. Five Parties exported *A. gueldenstaedtii* caviar: the Russian Federation (46 t), Iran (40 t), Kazakhstan (7.5 t), Azerbaijan (0.867 t) and Romania (0.233 t). The largest importers of *A. gueldenstaedtii* caviar were Switzerland (19 t), the USA (17 t), France (16 t) and Germany (15 t).

Exports of other specimens of *A. gueldenstaedtii* reported by CITES Parties for 1998 are: 329,417 live fish, mostly of captive bred or "F" (First Generation) sources from Hungary and Poland, and 686 kg of meat almost entirely from Iran. The trade in live *A. gueldenstaedtii* is by far the highest of all sturgeon species. The main exporters are Central European countries, where freshwater fish hatcheries and farms have been successful since the 1960s.

Re-exports of *A. gueldenstaedtii* from Hungary in 1999 (number of fingerlings)

	Germany	Belgium	UK	France	Italy	Netherlands	Austria
Live fingerlings	60,500	25,000	25,000	15,000	5,000	7,000	4,500

Source: CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000

Exports of *A. gueldenstaedtii* from Russia in 1999 (kilogrammes)

	Quantity (kg)	Description	Importer	No of permits
<i>A. gueldenstaedtii</i>	37.30	Live eggs	GR, DE, SY, AM, HU	12 Export
	28,017.68	Caviar	US, DE, DK, NL, FR, GB, BE	63 Ex. + 9 Re-ex.
	8,696.50	Meat	US	4 Export

Hybrids:

<i>A. baerii</i> x <i>A. gueld.</i>	3.50	Live eggs	PL, DE	2 Export
<i>A. gueld.</i> x <i>A. baerii</i>	37.50	Live eggs	CN, SY	3 Export

Source: CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 September 2000

Export quotas for *A. gueldenstaedtii* (kilogrammes)

		1998		1999		2000		
		Caviar		Caviar	Meat	Caviar	Meat	Fert. eggs
Azerbaijan				5,900	50,000	4,350	43,500	
Iran	<i>A. gueldenstaedtii</i>	10,000		14,000		52,000		
	(<i>A. persicus</i>)	(45,000)		(53,000)		including <i>A. persicus</i> see above		
Kazakhstan				5,500		4,200	49,000	
Romania				1,250	3,500	1,800	20,000	
Rus. Fed.		69,000		40,000		Caspian Azov	25,090 8,500	75,000 17,000
	<i>A. gueldenstaedtii</i> x <i>A. ruthenus</i>							90
	<i>A. gueldenstaedtii</i> x <i>A. baerii</i>							15
	<i>A. baerii</i> x <i>A. gueldenstaedtii</i>							10
Turkmenistan						500		
Annual total		124,000		119,650	53,500	95,940	204,500	175

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Note that Iran has established a total combined export quota of 52,000 kg for caviar of *A. gueldenstaedtii* and *A. persicus* for 2000. This is likely to be a result of the difficulty in differentiating between these two species (see Species Identification).

In 1999, new specimens appeared in the Annual Reports, such as 35 kg of swim bladders exported by Iran. Sturgeon swim bladders are exported for processing into isinglass and best quality fish glue. The glue has a wide range of uses, including antique furniture and painting restoration, and once diluted, to clarify beer, wine and champagne.

Note: The use of the term “eggs” and the unit “kg”, “g” or no unit (see # in Appendix and table on export quotas, above) to describe shipments of both caviar and fertilised eggs complicates the analysis of trade data. Similar difficulties arise from the unit used for export quotas of fertilised eggs, which is kilogrammes (“kg”) instead of “number of specimens”. Kilogrammes do not directly correspond to the number of specimens, and hence does not allow the effective evaluation of quotas. This confusion is also a source of mis-interpretation of data and is an obstacle to the proper monitoring of international trade, e.g. underestimation of live specimens in trade.

Illegal trade: In 1998, USA Customs confiscated *A. gueldenstaedtii* caviar of unknown origin upon import. A total of 1,215 kg was confiscated, comprising of 693 kg re-exported by Turkey and 522 kg re-exported by the United Arab Emirates. Discrepancies between trade statistics of countries importing and exporting caviar can indicate illegal activities. Such discrepancies were found in 1995 and 1996 for instance for Turkey (customs declared caviar exports 21 t and 80 t respectively) and European Union Member States (6 t and 20 t respectively) (Raymakers, 1999). Additionally, Turkish 1996 caviar exports totalled 121 t (Raymakers, 1999), while the country’s sturgeon fisheries has been limited to only few incidental catches since the late 1980s (Edwards and Doroshov, 1989).

Poaching and smuggling are closely related and have been intensively reported in the media of both range States and importing countries (Evtouchenko, 1997; McDonald, 2000; Snyder, 2000). However, much of the illegally caught sturgeon is destined to the domestic market, particularly meat (Anon., 1998).

Several Member States of the European Union are also detailed in CITES data as re-exporters of *A. gueldenstaedtii* caviar of unknown origin. According to CITES Annual Reports, these shipments were not confiscated upon import.

CONSERVATION MEASURES

A. gueldenstaedtii is not fully protected in any of its range States. However, a license is required for fishing in most of the countries and in Iran private sturgeon fisheries are prohibited. There is no specific monitoring programme for this species in any range State. Fish lifts and artificial spawning grounds have been introduced to certain regions of the Caspian.

Regarding juveniles restocking efforts, despite the large-scale aquaculture programmes of the Russian Federation and Iran, *A. gueldenstaedtii* stocks have continued to decline (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Azerbaijan: The Intergovernmental Commission for Caspian Biological Resources, composed of representatives of all range States, has set annual catch quotas (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Annual catch quotas for Azerbaijan (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. guelden.</i>	70	225	225	?	?	?	?	?	?	?	54.8
Total	200	308	510	480	360	182	160.5	160.5	160.5	108.67	108.35

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Three hatcheries were built in the lower Kura River by the authorities of the former Soviet Union in the 1960s. A feasibility study of the World Bank reports that the total capacity of the hatcheries was originally 12 million, but that this capacity had dropped significantly over the years (T. Turner, World Bank consultant, *in litt.* to TRAFFIC Europe, 6 October 2000). World Bank funds have been allocated for the construction of a new sturgeon hatchery and farm that will have the capacity of 15 million fry per annum and should be operational in the next few years. Concerns have been raised over the availability of adult sturgeon to constitute the broodstock of the farm.

Estimated combined number of fry released by three hatcheries (million fingerlings)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. gueld.</i>	13.370	5.330	2.130	1.230	0.750	0.940	3.030	3.770	4.490	10.186	8.869
Total	17.523	9.082	2.980	1.838	1.142	1.242	4.070	6.065	6.220	20.290	16.965

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

In Azerbaijan, the State institution responsible for sturgeon fishery and processing matters is the "State Concern Azerbalyg". Commercial sturgeon fisheries are prohibited in the open sea. No fishing license for sturgeon was required until 2000. The only fishing gear permitted for sturgeon catch in the Kura River is "sweep nets" of 35-55 mm mesh size. The minimum size limit set for *A. gueldenstaedtii* catch is 105 cm (Anon., 2000c).

Bulgaria: Measures at national level as described hereunder are needed, but they must be linked to co-ordinated actions of all countries in the Black Sea region. In Bulgaria, sturgeon fisheries are prohibited for 30 days, from 20 April to 20 May each year, which corresponds with the peak of the breeding season. The authorities may approve a 15-day extension of the closed season pending the natural conditions of the year. Explosives, poison, bottom trawling and dragging, as well as stunning substances and devices are prohibited fishing methods. The minimum size limit for *A. gueldenstaedtii* is 80 cm. Export quotas are set on the basis of catch data, trends and the biological characteristics of the species (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). A system was established for distribution of export quotas among export companies that are obliged to restock the Danube River with sturgeon fry before they are authorised to export sturgeon specimens. For example, for a particular year, for each kilogramme of caviar exported, the company must release 30 juvenile sturgeon (Anon., 2000c). The restocking programme is controlled by relevant government agencies.

Number of fingerlings released through the restocking programme (number of fingerlings)

	1998	sizes	1999	sizes	2000	sizes
<i>A. gueldenstaedtii</i>	1,000	200 g each	27,400	20 to 120 g each	20,400	25 g to 1.8 kg each

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Due to budgetary constraints, Bulgaria has not monitored the wild sturgeon population in recent years, nor the success of its restocking programme (e.g. through tagging of released fingerlings). However, plans have been prepared for these activities. A special prohibition was adopted on the use of "unbaited hooks" for sturgeon catch (Anon., 2000c).

Hungary: *A. gueldenstaedtii* is protected in Hungary. According to *Decree No. 1/1982 (III.15.) OKTH on the Protected and Strictly Protected Species of Flora and Fauna, Value of their Specimens, Determination of the Range of Protected and Strictly Protected Caves and Exemptions from Restrictions and Prohibitions Set for Certain Protected Animal Species*, last amended by the *15/1996 (VII.26.) Decree of the Minister for Environment*. The *Nature Conservation Act No. 53 of 1996, Article 43* orders that for all protected species, prior authorisation from the National Park Directorate is required for the: capture; killing; possession; captive breeding; introduction of specimens from exotic population; artificial exchange of genetic matter between populations; exchange; sale and purchase of *A. gueldenstaedtii* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: At all sturgeon catch stations, biometric measurements are carried out for each species. The age, length, sex and weight of fish landed are recorded. The population structure is estimated on the basis of age groups. As a result of these studies carried out for the past 10 years, the number of catch stations were decreased by 50%, changes were adopted on sturgeon catch methods, and lower export quotas were set for the past five years. Export quotas are established on the basis of results of research programmes and annual monitoring of stocks. These activities include the following parameters and activities: Maximum Sustainable Yield and total catch, rehabilitation of stocks, management and sustainable exploitation through restricting illegal catch and overfishing. The monitoring of fishing efforts and regulation of the time (date) of catch are also carried out on the basis of research results (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Annual release of *A. gueldenstaedtii* and *A. persicus* fry (million fingerlings)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<i>A. persicus</i>	4.06	5.92	2.93	3.57	4.66	8.05	11.02	18.75	22.59	17.30
A. gueldenstaedtii		0.04			0.30	0.52	0.67	0.92	0.42	0.96
Total	4.56	6.60	3.45	4.17	5.91	9.13	12.35	21.63	24.56	19.10

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

A labelling system is in place, where each tin of caviar holds a sign or mark that indicates at least the date and place of catch, the name of the catch station, name of processor and the species.

Iranian fishing regulations specify a minimum catch size limit of 1 m. *A. gueldenstaedtii* is artificially bred and reared. More than 10 million fingerlings of this species have been released into the rivers leading to the sea in recent years. Additional conservation measures have been taken for all sturgeon in the southern shores of the Caspian Sea, for instance banning the use of gillnets has been very effective in the conservation of this species. The autumn spawning population of this species is used for breeding operations in Iran (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Commercial sturgeon catch is only allowed in the tributaries of the Caspian Sea, however scientific catch is permitted in the open sea with an annual quota of 20 t (Anon., 2000c).

Sturgeon fisheries are regulated by law. There are two fishing seasons, one during spring spawning and the other during the winter migration. The exact dates of the fishing seasons vary with the climate and are therefore set annually by authorised scientific organisations. As in most former Soviet Republics, "sweep nets" (bottom nets that are hauled several times a day) with a minimum mesh size and which are marked with the name and address of the owner are the only legal fishing gear for sturgeon. Each fisherman can only set one net at a time. A fishing license is required. Sturgeon fishing grounds (called "tonia") are limited: there are 12 on the Ural River and 8 on the Kigach River. The minimum size limit for *A. gueldenstaedtii* is 105 cm. The ratio of males and females caught is not regulated nor monitored. The purchase price at the landing site is based on a fixed legal rate of caviar weight extracted per female; *A. gueldenstaedtii* is set at 18.6% of the total body weight. The annual commercial catch quotas are allocated to Kazakhstan by the Intergovernmental Commission for Caspian Biological Resources which meets annually in Astrakhan (CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000).

Annual commercial catch quotas allocated to Kazakhstan (tonnes)

	1996	1997	1998	1999	2000
A. gueldenstaedtii			135	80	52
Total	375	480	460	405	371

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Two hatcheries exist in Kazakhstan; Uralo-Atyrau and Atyrau. These have been operational since 1998.

Annual release of *A. gueldenstaedtii* fry (number of fingerlings)

	1998	1999	2000
A. gueldenstaedtii	647,200	1,492,800	1,266,300
Total	1,706,193	5,318,471	5,507,160

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Moldova: *A. gueldenstaedtii* catch is prohibited since the species was listed in the Red Data Book in 1995 (Anon., 2000c).

Romania: The type of sturgeon fishing gear is strictly regulated by Romanian law. The fishing season is set every year by Ministerial Order. A fishing license is required to catch sturgeon in Romanian waters. The minimum size limit for *A. gueldenstaedtii* is 140 cm. The catch quota for *A. gueldenstaedtii* for 1998 and

1999 was set at 12.5 t/year. Catch quotas are calculated by independent scientific institutes, such as the Danube Delta Institute in Tulcea and the "Centre for Conservation, Fish Farming, Fishing and Fish Industrialisation" in Galati (Anon., 2000c).

Release of *A. gueldenstaedtii* and *A. stellatus* fry in the Danube River (number of fingerlings)

	1993	1994	1995	1996	1997	1998	1999
<i>A. gueldenst.</i>			10,000				8,250
<i>A. stellatus</i>	1,000	4,000		10,000	11,600	10,000	6,750

Source: Centre for Conservation, Fish Farming, Fishing and Processing, Galati, cited in Anon., 2000c

Russian Federation: For Caspian Sea populations the former Soviet Republics set annual catch quotas in agreement with Iran. Sturgeon catch quotas have been established by the Intergovernmental Commission for Caspian Biological Resources for many years. The annual share of sturgeon catches of each former Soviet Union Caspian range State as adopted by the Commission since 1993 are as follows: the Russian Federation, 70%, Kazakhstan 17.6%, Azerbaijan 6.1% and Turkmenistan 6.3%. There are two fishing seasons in the Caspian Sea waters: from ice melting to June 5 and from 1 September until the time of ice formation. A license is required to catch sturgeon in Russian waters. The minimum size limit for *A. gueldenstaedtii* caught in Caspian Russian waters is 105 cm along the coast and 86 cm in the northern Caspian Sea. Currently, Russian catch quotas are declared by the Federal Government following the recommendations of the State Fisheries Committee, made in consensus with an independent council of experts (Anon., 2000b).

Release of *A. gueldenstaedtii* in the Volga River and the Sea of Azov (number of fry)

	1979-1980	1981-1985	1986-1990	1991-1995	1996-1998
Volga River	25,000,000	35,000,000	40,800,000	42,000,000	28,000,000
Sea of Azov	22,050,000	15,360,000	67,300,000	13,800,000	18,233,333

Source: Azov and Caspian Fisheries Research Institute (Az- and KaspNIRKh), cited in Anon., 2000b

From 1979 to 1998, approximately 54.2% 45.4% of all fry released by Russian hatcheries in the Volga River and the Sea of Azov respectively comprised *A. gueldenstaedtii* (Anon., 2000b).

Export quotas established by the CITES Management Authority of the Russian Federation, consist of 90% of the total caviar production, which is based on the catch quotas set for each species of sturgeon (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). The annual catch quotas or Total Allowable Catches (TACs) allocated by the State Fisheries Committee, is based on the recommendations of scientific agencies such as the Russian Federal Research Institute of Fisheries and Oceanography ("VNIRO") and the Pacific Research Institute of Fisheries and Oceanography ("TINRO"). The TAC includes the commercial catch, the scientific catch, and the catch of mature males and females destined for artificial reproduction.

Recent data on the monitoring of catch per species, including the size composition and sex ratio of fish caught, are not available. The domestic consumption of sturgeon products in the Russian Federation, mostly meat and caviar, is unknown (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000).

Turkey: In 1979, with the exception of *H. huso* of more than 140 cm long, the fishery for sturgeon was prohibited in Turkish waters (Edwards and Doroshov, 1989). Circular 34/1, effective from 2000-2002, prohibits the collection of Acipenseriformes (CITES Management Authority of Turkey, *in litt.* to TRAFFIC Europe, 20 September 2000).

Turkmenistan: Within the country's coastal commercial waters, sturgeon fishery is allowed for scientific research only, with an annual catch limit of 20 t. The Turkmenistan State Committee of Fisheries issues an annual license for sturgeon commercial catches to the State enterprise, Balkanbalyk. No other company, Turkmen or foreign, has the right to engage in commercial sturgeon fisheries in Turkmen inland waters. An annual catch quota for *A. gueldenstaedtii* in Turkmenistan is set by the Intergovernmental Commission for Caspian Biological Resources. There is no sex ratio set for sturgeon landings. Information on any existing

minimum size limits per species or data on the annual catch quotas set for Turkmenistan in the 1990s are unknown. (Anon., 2000b.)

Ukraine: Sturgeon fisheries in the Danube River are permitted all year around except from 16 March to 15 April. The by-catch of sturgeon is regulated. For example, for sprat trawling a tolerance of two sturgeon may be caught per tonne of sprat. The mesh size of nets used to catch sturgeon should be a minimum of 120 mm and a maximum of 150 mm. Only fishermen who were allocated an annual quota for sturgeon by the State Committee for Fishery are allowed to catch sturgeon. The minimum size limits for *A. gueldenstaedtii* are 110 cm for those caught in the Black Sea and the Danube River, and 90 cm for those caught in the Sea of Azov (Anon., 2000b.).

Annual catch quotas for *A. gueldenstaedtii* from 1997 to 1999 (tonnes)

	1996	1997	1998	1999
Total	505	424	210	225
<i>A. gueldenstaedtii</i> Sea of Azov and adjacent waters		275	135	180
Black Sea and Danube River		24	20	12

Source: State Committee for Fishery, cited in Anon., 2000b

There is currently no sturgeon fry restocking programme in the Ukraine.

Yugoslavia: The closed fishing season set in the Danube River for *H. huso*, *A. gueldenstaedtii* and *A. stellatus*, is from 15 May to 15 June. Certain types of fishing gear are prohibited to catch sturgeon and an agreement was signed between Romania and Yugoslavia on the prohibition of gear that causes high mortality of untargeted fish. Any fishing activity, including all commercial and sport fisheries, requires a license in Yugoslav waters. The licenses are issued on a per district basis and the sturgeon fishing grounds are limited to 17.8 km downstream of the Djerdap II dam and the mouth of the Timok River. The minimum size limit for *A. gueldenstaedtii* is 80 cm and the catch quota for the species from 1996 to 2000 was set at 3,000 kg per annum. A restocking programme is being carried out by the Centre for Fish Reproduction of the "Djerdap Fishing Company" in Malla Vrbica town (Anon., 2000b).

CAPTIVE BREEDING

The reduction in the area of natural spawning grounds due to river flow regulation has caused populations to decline. Fish reared in hatcheries have therefore been introduced to maintain the natural stock. In the early 1990s approximately 30% of the Caspian stock reportedly originated from hatcheries (Barannikova, 1995), whilst almost all (80-95%) generations of *A. gueldenstaedtii* in the Sea of Azov reportedly derived from aquaculture (Volovik *et al.*, 1993). The current status of re-stocking in the Sea of Azov is unclear.

The species has been successfully bred in aquaculture facilities in Austria, Belgium, Germany, Hungary, the Netherlands, Poland and Romania. It has been hybridised with various species such as Siberian Sturgeon *A. baerii* and Sterlet *A. ruthenus*.

Azerbaijan: Three hatcheries were built in the 1960s and have contributed to the former Soviet Union sturgeon restocking programme ever since. Their combined annual production was 12 million fry, but this has dropped significantly due to rising level of the Caspian Sea and the lack of funds allocated to hatchery maintenance (T. Turner, World Bank consultant, *in litt.* to TRAFFIC Europe, 6 October 2000).

Bulgaria: A fish farm was established in 1997 in Boliartzi village (Plovdiv region). It is the only existing hatchery. All juveniles released in the Danube River in Bulgaria from 1998 to 2000 were produced there. The hatchery has a total capacity of 60,000 juveniles and the farm is currently stocked with 3.5 year old *H. huso* and *A. gueldenstaedtii* that weigh 11-14 kg. In 1998, fingerlings were released in to the Ovcharitza dam reservoir: 200 juvenile *A. gueldenstaedtii* of 200 g average weight and 200 hybrids of *A. gueldenstaedtii* and *A. baerii* of 200 g average weight. Since 1999, attempts have been made to breed sturgeon in artificial water basins near Sofia (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Hungary: There is no broodstock of the species in Hungary. Fertilized eggs are imported from Russia, and fingerlings are re-exported. There is no caviar production in Hungarian farms (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: More than 500 *A. gueldenstaedtii* between the ages of 9-11 years are currently held in captive breeding in earthen ponds in Iran (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). Hatcheries have been operational for 20 years.

Italy: *A. gueldenstaedtii* larvae were imported six or seven years ago and are present in rearing plants for the production of juvenile fish and hybridisation. For 6-7 year old *A. gueldenstaedtii* females, 1 kg eggs corresponds to an average 45,000 eggs. The 1998 production of one of the aquaculture facilities was 3,000 fingerlings (from 60,000 eggs and 20,000 hatchlings) of *A. gueldenstaedtii*. This indicates a 5% hatching-survival rate (Azienda Agricola and Agroittica Lombarda, Italian sturgeon farmers, *in litt.* to TRAFFIC Europe-Italy, 1999).

Kazakhstan: Two hatcheries exist in Kazakhstan, Uralo-Atyrau and Atyrau, which have been operational since 1998 (CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, September 2000).

Poland: The presence of fish farms and hatcheries of sturgeon in Poland was confirmed by the CITES Management Authority of Poland (Dr Z. Krzeminski, Deputy Director, *in litt.* to TRAFFIC Europe, 31 August 2000). The following sturgeon species are bred in captivity: *Acipenser baerii*, *A. gueldenstaedtii*, *A. ruthenus*, *A. stellatus*, *H. huso*, *Polyodon spathula* and hybrids of some species. In most cases, hybrids are F1 generation live specimens (fertilised eggs) imported by Poland from range States, particularly from the Russian Federation. Some farms keep F2 generation specimens that were bred in Poland and are used as broodstock, partly for the production of juveniles to be sold on domestic and international markets. Various hybridisations of *A. gueldenstaedtii* have been performed and appear commercially promising.

Russia: As detailed above in Conservation Measures, aquaculture has been ongoing since the 1960s.

United States of America: Sturgeon are in a preliminary stage of culture in Hawaii (CITES Management Authority of the US, *in litt.* to TRAFFIC Europe, September 2000).

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database, but according to trade data there is currently a significant production of *A. gueldenstaedtii* juveniles in commercial farms (see Appendix).

REFERENCES

- Anon. (1996). *A Guide to Interpreting Outputs from the WCMC CITES Trade Database*, Produced by the World Conservation Monitoring Centre, Cambridge UK. Pp. 16.
- Anon. (1998). *Sturgeon catch and trade in the Russian part of the Caspian Sea*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, Pp. 21.
- Anon. (2000a). [http://www.cites.org/eng/dbase/fauna-](http://www.cites.org/eng/dbase/fauna)
- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, Pp. 23.
- Anon. (2000c). *Sturgeon fisheries management and trade control measures in the Caspian Sea and Black Sea/ Sea of Azov range States*. TRAFFIC Europe field investigations, December 1999-January 2000. Unpublished.
- Bacalbasa-Dobrovici, N. (1997). Endangered migratory sturgeons of the lower Danube River and its delta. *Environmental Biology of Fishes* 48:201-207.
- Barannikova, I.A., Burtsev, I.A., Vlasenko, A.D., Gershanovich, A.D., Markarov, E.V. and Chebanov, M.S. (1995). Sturgeon Fisheries in Russia. *Proceedings of the Second International Symposium on Sturgeons, September 6-11, 1993*. Moscow-Kostroma-Moscow. VNIRO Publications. Pp. 124-130.
- Birstein, V.J. (1993). Sturgeon and Paddlefishes: threatened fish in need of conservation. *Cons. Biol.* 7(4):773-787.

- De Meulenaer, I. and Raymakers, C. (1996). *Sturgeons of the Caspian Sea and the international trade in caviar*. TRAFFIC International, iv+ 71 Pp.
- Dobbs, M. (1992). The Coming Caviar Crisis: Overfishing, Pollution Endanger Sturgeon. *The Washington Post*, 30 May 1992.
- Doward, J. (2000). Roe rage as piracy rules. *The Guardian*, 21 May 2000.
- Dumont, H. (1995). Ecocide in the Caspian Sea. *Nature* 377:673-674.
- Edwards, D. and Doroshov, S. (1989). Appraisal of the Sturgeon and Seatrout Fisheries and Proposals for a Rehabilitation Programme. FAO Field Document FI.TCP/TUR/8853. Pp. 38;
- Evtouchenko, A. (1997). Au Dagestan une explosion fait 54 morts. Accusés: les gangs du caviar. *Courrier International*, N° 323 (9-15 January 1997)[in French].
- Hensel, K. and Holèik, J. (1997). Past and current status of sturgeon in the upper and middle Danube. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 185-200.
- Hochleithner, M. and Gessner, J. (1999). *The Sturgeon and Paddlefishes (Acipenseriformes) of the World: Biology and Aquaculture*. AquaTech Publications, Kitzbühl. Pp. 165.
- Holèik, J. (1989). Návrh èerveného zoznamu ohrozených kruhoustych a rýb Slovenska. *Pamiatky a príroda* 20:26-28
- Holèik, J. (1995). 4.druh Acipenser (Acipenser) gueldenstaedti Brandt, 1833. In: V.Baruš and O.Oliva (eds.) *Mihulovci Petromyzontes a ryby Osteichthyes*. 1. Fauna ÈR a SR 28/1. Academia, Praha. Pp. 391-397.
- IUCN. (1996). 1996. *Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- Jankoviè, D. (1993). Populations of Acipenseridae prior and after the construction of the HEPS Djerdap I and II. *Acta Biologica Iugoslavica, Seria E. Ichthyologia* 25:29-34
- Jankoviè, D. (1996). Ichthyofauna of the Danube in the Djerdap area after the construction of the Iron Gate I Hydroelectric Power Station. *Acta Universitatis Carolinae, Biologica* 40 (1996):123-131.
- Karapetkova, M., Zhivkov, M. and Pchelarov, T. (1995). *Ribite v B'Igariya*. Geya Libris, Sofia. 247 Pp.
- Khodorevskaya, R.P., Dovgopol, G.F., Zhuraleva, O.L., and Vlasenko, A.D. (1997). Present status of commercial stocks of sturgeons in the Caspian Sea Basin. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 209-219.
- King, R. (1998). [Por un puñado de caviar. *El Mundo*, 13 December 1998.] (in Spanish)
- Laluyee, F. (1996). Migration studies of sturgeons in Sefidrud, Tajen and Gorganrud River. Iranian Fisheries Research and Training Organisation Publications.
- Levin, A.V. (1997). The Distribution and Migration of Sturgeon in the Caspian Sea. In: Birstein, V.J., Bauer, A. Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 13-19.
- McDonald, L. (2000). Fish that laid the golden egg faces extinction. *Daily Express* (UK), 26 July 2000.
- Manea, G.H., (1966). Contributii in studiul sturionilor din apele Romiiniei si al reproducerii lor in legatura cu constructiile hidroenergetice pe DFunarea inferiara. 1. Une aspecte ale biologici sturionilor. *Buletinul Institutului de Cercetari Piscicole* 25:62-86.
- Nasri Chari, A. (1993). Comparative study on morphobiological parameters in Russian and Persian sturgeon in the southern shores of Caspian Sea in order to declare Persian sturgeon as an independent species. Tehran University Publications.
- Pavlov, P.I., (1980). Ribi. Fauna Ukrainy. 8/1. Naukova Dumka, Kiiv. Pp 336.
- Pintèr, K. (1991). Sturgeon in Hungary, past and present situation. In: P. Williot (Ed.) *Actes du premier colloque international sur l'esturgeon. Bordeaux, 3-6 October 1989*. Pp. 173-178.
- Ramin, M. (1998). Migratory studies in sturgeons and their limiting factors in the Sefidrud River. Gilan Fisheries Research Centre Publications.
- Raymakers, C. (1999). Trade in Sturgeons from the Caspian Sea. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 149-161.
- Sagers, M.J. (1994). The oil industry in the southern-tier former soviet republics. *Post-Soviet Geography*, 35(5):267-298.
- Shagaeva, V.G., Nikolskaya, M.P., Akimova, N.V. Markov, K.P., and Nikolskaya, N.G. (1993). A study of the early ontogeny of Volga sturgeon (Acipenseridae) subjected to human activity. *J. Ichthyol.* 33(6):23-41.

- Schrope, M. (2000). Alien Invader: A hungry stowaway is saving the Black Sea. *New Scientist*, 2255:10.
- Simonovic, P.D., and Nikolic, V.P. (1996). Freshwater fish of Serbia: an annotated check list with some faunistic and zoogeographical considerations. *Bios (Macedonia, Greece)*, 4:137-156.
- Snyder, D. (2000). Md. Caviar Company Admits Smuggling. *The Washington Post*, 22 July 2000.
- Vlasenko, A.D. (1990). Sturgeon Population Size in the Caspian Sea. *Rybnoe Khozyaistvo*, 7:53-56.
- Vlasenko, A.D., Pavlov, A.V., Sokolov, L.I. and Vasil'ev, V.P. (1989). *Acipenser gueldenstaedti* Brandt, 1833. In: Holèik J. (ed). The Freshwater Fishes of Europe. Vol. I/II: General Introduction of Fishes. Acipenseriformes. Wiesbaden, AULA-Verlag. In: *Document Doc. 10.89; Prop. 10.65. 1997. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America*. Pp. 295-344.
- Volovik, S.P., Dubinina, V.A., and Semenova, A.D. (1993). Hydrobiology and dynamics of fisheries in the Asov Sea. Studies and reviews. *General Fisheries Council for the Mediterranean*. No. 64. FAO, Rome. Pp. 1-58.
- Ward, P. (2000a). Azov Sturgeons Can't be Touched. *Crimean Time*, 18 August 2000.
- Ward, P. (2000b). Makhachkala: the Scale of Poaching in the Caspian has Reached Enormous Proportions. *Regions.Ru/Russia Regions*, 15 August 2000.
- Williot, P. and Bourguignon, G. (1991). Sturgeon and caviar production, current status and perspectives. In: Williot, P. (ed.), *Acipenser*, Cemagref Publications, Pp. 509-513.
- Zarkua, Z. and Tsuladze, V. (1999). Initiative needed to boost sturgeon numbers. *Eurofish* 3/99:40-41.

Gross exports* and export quotas for *Acipenser gueldenstaedtii* 1998

TAXON	TERM	UNIT	Exporter	1998	Export Quotas 1998
<i>Acipenser gueldenstaedtii</i>	eggs		AE	184	
<i>Acipenser gueldenstaedtii</i>	eggs		CH	12	
<i>Acipenser gueldenstaedtii</i>	eggs		DE	118	
<i>Acipenser gueldenstaedtii</i>	eggs		FR	70	
<i>Acipenser gueldenstaedtii</i>	eggs		GB	330	
<i>Acipenser gueldenstaedtii</i>	eggs		IR	1,500	
<i>Acipenser gueldenstaedtii</i>	eggs		LT	933	
<i>Acipenser gueldenstaedtii</i>	eggs		RO	57	
<i>Acipenser gueldenstaedtii</i>	eggs		RU	89	
<i>Acipenser gueldenstaedtii</i>	eggs		SG	1	
<i>Acipenser gueldenstaedtii</i>	eggs		US	57	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	AE	5,124	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	AZ	867	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	BE	203	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	CH	8,627	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	DE	7,205	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	DK	1,011	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	ES	21	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	FI	958	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	FR	609	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	GB	1,079	
<i>Acipenser gueldenstaedtii</i>	eggs	kg	GR	42	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	IR	40,390	10,000 (<i>A. gueldenstaedtii</i>) 45,000 (<i>A. persicus</i>)
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	KZ	7,543	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	NL	1	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	NZ	2	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	PL	6,530	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	RO	233	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	RU	46,359	69,000 (incl. AZ & KZ)
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	SG	81	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	TR	4,655	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	US	1,667	
<i>Acipenser gueldenstaedtii</i>	Eggs	kg	XX	4	
<i>Acipenser gueldenstaedtii</i>	Live		CZ	1,500	
<i>Acipenser gueldenstaedtii</i>	Live		GB	335	
<i>Acipenser gueldenstaedtii</i>	Live		HU	169,500	
<i>Acipenser gueldenstaedtii</i>	Live		PL	155,000	
<i>Acipenser gueldenstaedtii</i>	Live		US	3,000	
<i>Acipenser gueldenstaedtii</i>	Live		XX	11	
<i>Acipenser gueldenstaedtii</i>	Live	kg	DE	21	
<i>Acipenser gueldenstaedtii</i>	Live	kg	RU	50	
<i>Acipenser gueldenstaedtii</i>	Meat	kg	IR	669	
<i>Acipenser gueldenstaedtii</i>	Meat	kg	RU	15	
<i>Acipenser gueldenstaedtii</i>	Meat	kg	US	2	
<i>Acipenser gueldenstaedtii</i>	Skins		CH	8	
<i>Acipenser gueldenstaedtii</i>	Skins		IR	8	
<i>Acipenser gueldenstaedtii</i>	Unspec- ified		RU	1	

* **Definition:** "Gross exports are the sum of all reported exports and re-exports in a particular commodity or species in a particular year or series of years. Gross trade is thus a simple measure of the total number of

items recorded in international trade. However, gross trade may be an overestimate of the total number of actual specimens in trade as re-exports are not deducted from the total." (Anon., 1996). This is particularly applicable to caviar of which shipments are often exported, re-packaged and re-exported. For instance, in the mid-1990s, Germany re-exported as much as 45% of its annual caviar imports (De Meulenaer and Raymakers, 1996).

Comparative Tabulation of trade in *Acipenser gueldenstaedtii* 1998

Year	Imp.	Exp.	Origin	Imports reported			Exports reported							
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S	
1998	CH	AE	IR		64 kg			O		64 kg	eggs		T	O
1998	CH	AE	RU							369 kg	eggs		T	W
1998	CH	AE	RU		63 kg	eggs		T	O					
1998	DK	AE	RU		10 kg	eggs		T	O					
1998	FR	AE	RU							644 kg	eggs		T	W
1998	FR	AE	RU		487 kg	eggs		T	O					
1998	GB	AE	RU							61 kg	eggs		T	W
1998	JP	AE	RU							65 kg	eggs		T	O
1998	US	AE			522 kg	eggs		T	I					
1998	US	AE	RU		22 kg	eggs		T	I					
1998	US	AE	RU		956 kg	eggs		T	W	255 kg	eggs		T	W
1998	US	AE	RU		714 kg	eggs		T	O	3,497 kg	eggs		T	O
1998	US	AE	RU							159 kg	eggs		P	O
1998	US	AE	RU		104	eggs		T	O					
1998	US	AE	RU		80	eggs		T	W					
1998	US	AE	RU		20 kg	eggs		T	I					
1998	US	AE	XX		159 kg	eggs		T	O					
1998	CA	BE	RU							38 kg	eggs		T	O
1998	CA	BE	TR							11 kg	eggs		T	O
1998	TR	BE	RU							360 g	eggs		T	O
1998	US	BE	RU							153 kg	eggs		T	O
1998	AN	CH	IR							11 kg	eggs			W
1998	AT	CH	IR							8 kg	eggs			O
1998	BB	CH	IR							2 kg	eggs			O
1998	BE	CH	IR							4 kg	eggs			W
1998	BE	CH	IR		4 kg	eggs		T	W					
1998	BE	CH	KZ		605 kg	eggs		T	W	605 kg	eggs			W
1998	CH	CH	IR							77 kg	eggs			O
1998	CH	CH	RU							8 kg	eggs			O
1998	CY	CH	IR							1 kg	eggs			O
1998	DE	CH	IR		2,197 kg	eggs		T	W	1,470 kg	eggs			W
1998	DE	CH	IR							23 kg	eggs			O
1998	DE	CH	KZ		2,488 kg	eggs		T	W					
1998	DE	CH	RU							2,490 kg	eggs			W
1998	DK	CH	IR		252 kg	eggs		T	O	252 kg	eggs			O
1998	ES	CH	IR							1 kg	eggs			O
1998	FR	CH	IR		520 kg	eggs		T	W	1,039 kg	eggs			W
1998	FR	CH	IR		21 kg	eggs		T	O	294 kg	eggs			O
1998	FR	CH	IR		3 kg	eggs		P	W					
1998	FR	CH	KZ		780 kg	eggs		T	W	780 kg	eggs			W
1998	GB	CH	IR							21 kg	eggs			O
1998	GB	CH	KZ		12	eggs		T	W					
1998	GB	CH	RU							259 kg	eggs			W
1998	GB	CH	RU							201 kg	eggs			O
1998	HK	CH	IR							35 kg	eggs			O
1998	HK	CH	IR							27 kg	eggs			W
1998	IT	CH	IR							4 kg	eggs			O

Year	Imp.	Exp.	Origin	Imports reported			Exports reported							
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S	
1998	IT	CH	IR						8	skins			W	
1998	JP	CH	IR						82 kg	eggs			O	
1998	JP	CH	RU						19 kg	eggs			O	
1998	KW	CH	IR						20 kg	eggs			O	
1998	KW	CH	IR						11 kg	eggs			W	
1998	LU	CH	IR						2 kg	eggs			O	
1998	MA	CH	RU						60 kg	eggs			O	
1998	NL	CH	IR						42 kg	eggs			W	
1998	PA	CH	IR						2 kg	eggs			O	
1998	SA	CH	IR						1 kg	eggs			W	
1998	SG	CH	IR		44 kg	eggs		T	O	44 kg	eggs		O	
1998	SG	CH	IR		8 kg	eggs		T	W	8 kg	eggs		W	
1998	SG	CH	RU		1 kg	eggs		T	O	1 kg	eggs		O	
1998	US	CH	IR						13 kg	eggs			O	
1998	US	CH	IR						3 kg	eggs			W	
1998	US	CH	RU		1.5 kg	eggs		T	W					
1998	US	CH	RU						2 kg	eggs			O	
1998	ZA	CH	IR						3 kg	eggs			O	
1998	DE	CZ	RU		1,500	live		T	C	1,500	live		T	C
1998	AR	DE	KZ						2 kg	eggs			T	O
1998	AR	DE	RU						220 kg	eggs			T	O
1998	AU	DE	RU						60 kg	eggs			T	O
1998	BB	DE	RU						63 kg	eggs			T	O
1998	BR	DE	RU						160 kg	eggs			T	O
1998	CA	DE	KZ						20 kg	eggs			T	O
1998	CA	DE	KZ						16 kg	eggs			T	W
1998	CA	DE	RU						14 kg	eggs			T	O
1998	CH	DE	IR		993 kg	eggs		T	W	786 kg	eggs		T	W
1998	CH	DE	IR		904 kg	eggs		T	O	391 kg	eggs		T	O
1998	CH	DE	KZ						70 kg	eggs			T	W
1998	CH	DE	RU		78 kg	eggs		T	W	8 kg	eggs		T	W
1998	CH	DE	RU		57 kg	eggs		T	O					
1998	CL	DE	RU						60 kg	eggs			T	O
1998	CZ	DE	IR						300 g	eggs			T	O
1998	CZ	DE	IR		0 kg	eggs		T	W					
1998	CZ	DE	RU						682 g	eggs			T	W
1998	CZ	DE	RU						6 kg	eggs			T	O
1998	CZ	DE	RU		5 kg	eggs		T	W					
1998	ET	DE	RU						104 kg	eggs			T	O
1998	ET	DE	RU						37 kg	eggs			T	W
1998	HK	DE	AZ						51 kg	eggs			T	O
1998	HK	DE	IR						42 kg	eggs			T	O
1998	HK	DE	KZ						23 kg	eggs			T	O
1998	HK	DE	RU						66 kg	eggs			T	O
1998	JP	DE	AZ						15 kg	eggs			T	O
1998	JP	DE	KZ						25 kg	eggs			T	O
1998	JP	DE	RU						374 kg	eggs			T	O
1998	KR	DE	KG		21 kg	eggs		T	W					
1998	KR	DE	KZ						3 kg	eggs			T	W
1998	KR	DE	RU						25 kg	eggs			T	O
1998	KR	DE	RU		5 kg	eggs		T	W					
1998	MC	DE	RU						28 kg	eggs			T	O
1998	MT	DE	RU						336 g	eggs			T	O
1998	NO	DE	IR						5 kg	eggs			T	O

Year	Imp.	Exp.	Origin	Imports reported			Exports reported						
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	NO	DE	RU						4 kg	eggs		T	O
1998	NZ	DE	RU						60 kg	eggs		T	O
1998	PF	DE	RU						168 kg	eggs		T	O
1998	PL	DE	RU						184 kg	eggs		T	O
1998	SA	DE	RU						63 kg	eggs		T	O
1998	SG	DE	KZ		2 kg	eggs			5 kg	eggs		T	O
1998	SG	DE	RU		5 kg	eggs			5 kg	eggs		T	O
1998	TR	DE	RU						10 kg	eggs		T	O
1998	TW	DE	RU						40 kg	eggs		T	O
1998	US	DE	KZ		329 kg	eggs			829 kg	eggs		T	W
1998	US	DE	KZ		173 kg	eggs			261 kg	eggs		T	O
1998	US	DE	KZ		87	eggs							
1998	US	DE	RU		1,588 kg	eggs			2,114 kg	eggs		T	O
1998	US	DE	RU		210 kg	eggs							
1998	US	DE	RU		100 kg	eggs							
1998	US	DE	RU		40 kg	eggs							
1998	US	DE	RU		21	eggs							
1998	US	DE	RU		10	eggs							
1998	US	DE	XX		67 kg	eggs							
1998	US	DE	RU		21 kg	live							
1998	ZA	DE	RU						8 kg	eggs		T	O
1998	BR	DK	IR						200 kg	eggs		T	W
1998	BR	DK	IR						46 kg	eggs		T	O
1998	BR	DK	XX						268 kg	eggs		T	W
1998	CH	DK	IR		427 kg	eggs			303 kg	eggs		T	W
1998	CL	DK	RU						50 kg	eggs		T	W
1998	SA	DK	IR						11 kg	eggs		T	O
1998	US	DK	RU						10 kg	eggs		T	O
1998	US	DK	RU		10 kg	eggs							
1998	US	ES	RU		21 kg	eggs							
1998	RU	FI	RU						958 kg	eggs		T	W
1998	AD	FR	IR						10 kg	eggs		T	W
1998	AD	FR	IR						1 kg	eggs		T	O
1998	BR	FR	IR						2 kg	eggs		T	W
1998	CA	FR	AZ						6 kg	eggs		T	O
1998	CA	FR	KZ						10 kg	eggs		T	O
1998	CA	FR	KZ						7 kg	eggs		T	W
1998	CA	FR	RU						4 kg	eggs		T	O
1998	CH	FR	IR		249 kg	eggs			83 kg	eggs		T	W
1998	CH	FR	IR		153 kg	eggs			153 kg	eggs		T	O
1998	CI	FR	IR						4 kg	eggs		T	W
1998	CM	FR	KZ						2 kg	eggs		T	O
1998	HK	FR	KZ						2 kg	eggs		T	W
1998	MX	FR	KZ						1 kg	eggs		T	W
1998	SA	FR	IR						3 kg	eggs		T	O
1998	US	FR	KZ						155 kg	eggs		T	W
1998	US	FR	KZ		70	eggs							
1998	US	FR	RU		84 kg	eggs							
1998	US	FR	XX		42 kg	eggs							
1998	BH	GB	IR						150 kg	eggs		T	W
1998	BR	GB	IR						1 kg	eggs		T	W
1998	CH	GB	RU		201 kg	eggs							
1998	EG	GB	IR						1 kg	eggs		T	W
1998	HK	GB	IR						1 kg	eggs		T	W

Year	Imp.	Exp.	Origin	Imports reported			Exports reported						
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	JP	GB	IR						60 kg	eggs		T	W
1998	JP	GB	XX						14 kg	eggs		T	W
1998	NG	GB	IR						1 kg	eggs		T	W
1998	PH	GB	RU						1 kg	eggs		T	W
1998	SG	GB	IR						1 kg	eggs		T	W
1998	US	GB	IR						1 kg	eggs		T	W
1998	XX	GB	IR						50 kg	eggs		T	W
1998	XX	GB	KZ						50 kg	eggs		T	W
1998	XX	GB	HU						335	live		T	C
1998	ZA	GB	IR						243 kg	eggs		T	W
1998	ZA	GB	IR						80	eggs		T	W
1998	ZA	GB	KZ						75 kg	eggs		T	W
1998	ZA	GB	RU						77 kg	eggs		T	W
1998	ZA	GB	XX						250	eggs		T	W
1998	ZA	GB	XX						150 kg	eggs		T	W
1998	US	GR	RU		42 kg	eggs						T	O
1998	CH	HK	IR		50 g	eggs						T	W
1998	AT	HU	RU						2000	live		T	C
1998	BE	HU	RU						32,000	live		T	C
1998	BE	HU	RU						4,000	live		T	F
1998	DE	HU	RU						17,500	live		T	F
1998	DE	HU	RU		4,185	live			50,000	live		T	C
1998	FR	HU	RU						8,000	live		T	C
1998	FR	HU	RU						8,000	live		T	F
1998	GB	HU	RU						20,000	live		T	C
1998	NL	HU			5,000	live						T	C
1998	NL	HU	RU						20,000	live		T	C
1998	NL	HU	RU						8,000	live		T	F
1998	BE	IR			300 kg	eggs						T	W
1998	BE	IR			2 kg	meat						T	W
1998	CH	IR			2,885 kg	eggs						T	O
1998	CH	IR			1,570 kg	eggs						T	W
1998	CH	IR	RU		5,180 kg	eggs						T	W
1998	CH	IR			520 kg	meat						T	W
1998	CH	IR	RU		8	skins						T	W
1998	DE	IR			3,117 kg	eggs						T	O
1998	DE	IR			650 kg	eggs						T	W
1998	DK	IR			800 kg	eggs						T	W
1998	ES	IR			8,320 kg	eggs						T	W
1998	ES	IR			500	eggs						T	W
1998	FR	IR			15,418 kg	eggs						T	W
1998	FR	IR			100 kg	meat						T	W
1998	FR	IR			47 kg	meat						E	W
1998	GB	IR			2,149 kg	eggs						T	W
1998	US	IR			1,000	eggs							W
1998	US	IR			500 g	eggs						P	I
1998	US	IR	XX		400 g	eggs						T	I
1998	US	IR	XX		200 g	eggs							W
1998	US	LT	XX		933	eggs						T	I
1998	BN	NL	IR						1 kg	eggs		T	W
1998	US	NZ	RU		2 kg	eggs						P	I
1998	DE	PL	RU		2,400 kg	eggs						T	O
1998	DE	PL	RU						1,500 kg	eggs			W
1998	DE	PL	RU						900 kg	eggs			O

Year	Imp.	Exp.	Origin	Imports reported			Exports reported						
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	DE	PL							155,000	live			C
1998	DE	PL		75,000	live		T	C					
1998	GB	PL	RU						200 kg	eggs			W
1998	US	PL	RU	244.4 kg	eggs		T	I					
1998	US	PL	RU						3,930 kg	eggs			W
1998	US	PL	RU	760 kg	eggs		T	W					
1998	US	PL	RU	514 kg	eggs		T	O					
1998	US	PL	RU	428 kg	eggs		T	I					
1998	GR	RO		4 kg	eggs		T	W					
1998	US	RO		51.2 kg	eggs		T	W					
1998	US	RO		178 kg	eggs		T	W					
1998	US	RO		44	eggs		T	W					
1998	US	RO		13	eggs		T	O					
1998	AE	RU							207 kg	eggs		T	W
1998	AE	RU	KZ						89 kg	eggs		T	W
1998	AZ	RU							1,458 kg	eggs		T	W
1998	BE	RU		1,044 kg	eggs		T	W	1,128 kg	eggs		T	W
1998	CH	RU		4,391 kg	eggs		T	W	2,919 kg	eggs		T	W
1998	CH	RU		13,30 kg	eggs		T	O					
1998	CH	RU		10 kg	eggs		T	U					
1998	CH	RU	AZ						795 kg	eggs		T	W
1998	CH	RU	KZ						4,391 kg	eggs		T	W
1998	CN	RU							4 kg	eggs		T	F
1998	CZ	RU		2 kg	eggs		T	C	2 kg	eggs		T	C
1998	DE	RU		2,644 kg	eggs		T	W	4,819 kg	eggs		T	W
1998	DE	RU		1,265 kg	eggs		T	O	6,307 kg	eggs		T	O
1998	DK	RU		373 kg	eggs		T	W	373 kg	eggs		T	W
1998	FI	RU		450 kg	eggs		T	W	958 kg	eggs		T	W
1998	FR	RU		1,196 kg	eggs		T	W	1,204 kg	eggs		T	W
1998	FR	RU	KZ	120 kg	eggs		T	W	120 kg	eggs		T	W
1998	HU	RU							8 kg	eggs		T	F
1998	HU	RU							2 kg	eggs		T	C
1998	JP	RU							1,141 kg	eggs		T	W
1998	KR	RU							50 kg	live		S	F
1998	NL	RU		691 kg	eggs		T	W	2,749 kg	eggs		T	W
1998	US	RU		285 kg	eggs		T	I					
1998	US	RU		62.93 kg	eggs		T	W					
1998	US	RU		5,999 kg	eggs		T	W	13,443 kg	eggs		T	W
1998	US	RU		660 kg	eggs		T	O	4,244 kg	eggs		T	O
1998	US	RU		590 kg	eggs		T	I					
1998	US	RU		89	eggs		T	W					
1998	US	RU		15 kg	meat		T	W					
1998	US	RU		1	unspecified		I						
1998	CH	SG	IR	13 kg	eggs		T	O	13 kg	eggs		T	O
1998	US	SG	RU	68 kg	eggs		T	W					
1998	US	SG	RU	1	eggs		T	W					
1998	CH	TR	RU	660 kg	eggs		T	O	663 kg	eggs		T	O
1998	JP	TR	XX						25 kg	eggs		T	O
1998	US	TR		672 kg	eggs		T	I					
1998	US	TR	RU	920 kg	eggs		T	W					
1998	US	TR	RU	838 kg	eggs		T	O	1,080 kg	eggs		T	O
1998	US	TR	RU	7 kg	eggs		T	I					
1998	US	TR	RU	2 g	eggs		T	I					
1998	US	TR	RU	2 g	eggs		T	W					

Year	Imp.	Exp.	Origin	Imports reported			Exports reported						
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	US	TR	XX						2,888	kg	eggs	T	O
1998	US	TR	XX		21	kg	eggs					T	I
1998	AR	US	RU						7	kg	eggs	T	O
1998	AR	US	RU						2	kg	eggs	T	W
1998	CA	US	RU						57	kg	eggs	T	W
1998	CA	US	RU						3	kg	eggs	T	O
1998	CA	US	RU						1		eggs	T	O
1998	CA	US	RU						1		eggs	T	W
1998	CL	US	RU						53	kg	eggs	T	O
1998	CL	US	RU						25		eggs	T	W
1998	DE	US	KZ						1,800	g	eggs	T	W
1998	DE	US	RU						1,360	g	eggs	T	O
1998	DE	US	RU		579	kg	eggs		95	kg	eggs	T	O
1998	DE	US	RU						539	kg	eggs	T	W
1998	DE	US	XX		84	kg	eggs					T	O
1998	DK	US	RU						102	kg	eggs	T	O
1998	DK	US	RU						21	kg	eggs	T	W
1998	ES	US	RU		21	kg	eggs					T	O
1998	FR	US	RU						21	kg	eggs	T	W
1998	FR	US	RU						21	kg	eggs	T	W
1998	GR	US	RU						105	kg	eggs	T	W
1998	GR	US	RU						89	kg	eggs	T	O
1998	IT	US	RU						10.5	kg	eggs	T	O
1998	JP	US	RU						1	kg	eggs	T	W
1998	KR	US	CO		3,000		live					T	C
1998	MC	US	RU						61	kg	eggs	T	O
1998	MC	US	RU						57	kg	eggs	T	W
1998	MX	US	RU						4	kg	eggs	T	W
1998	MX	US	RU						2	kg	meat	T	W
1998	NL	US	RU						21	kg	eggs	T	W
1998	NO	US	RU						2	kg	eggs	T	O
1998	PT	US	RU						76	kg	eggs	T	O
1998	PT	US	RU						43	kg	eggs	T	W
1998	SE	US	RU						63	kg	eggs	T	O
1998	SG	US	RU						24.5	kg	eggs	T	W
1998	TR	US	RU						44	kg	eggs	T	O
1998	TR	US	RU						34	kg	eggs	T	W
1998	TR	US	RU						21		eggs	T	W
1998	XX	US	DE						6	g	eggs	P	O
1998	XX	US	RE						3	g	eggs	P	O
1998	XX	US	RU						166	g	eggs	P	O
1998	XX	US	RU						6		eggs	T	O
1998	XX	US	RU						3		eggs	P	O
1998	XX	US	RU						3	kg	eggs	P	O
1998	XX	US	RU						2	g	eggs	T	O
1998	ZA	US	RU						54	kg	eggs	T	O
1998	ZA	US	RU						3	kg	eggs	T	W
1998	US	XX	RU		3,500	g	eggs					P	I
1998	US	XX	RU			15	g	eggs				P	O
1998	US	XX				11	live					T	I
Total:				103,320.5	kg	eggs			82,008	kg	eggs		
				88,696		live			326,335		live		

Acipenser nudiiventris Lovetzky, 1828

Ship Sturgeon
Esturgeon à barbillons frangés
Esturión barba de flecos

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

The Ship Sturgeon *Acipenser nudiiventris* has always been rare in the Black Sea and the Sea of Azov and their tributaries. There were originally two isolated populations in the northern Caspian Sea, but only the Ural River population is thought to exist today. In Iran (southern Caspian Sea), the species is known to migrate in several rivers. Its former range in the Danube River included Austria and Slovakia. Most range States reported that their wild populations are currently low. Restocking programmes are carried out in Azerbaijan, Iran, and Kazakhstan, but although the species has been in captivity for 60 years, broodstock is currently difficult to obtain. The species is rarely raised in captivity and there is no recorded international trade in products from aquaculture. Of all range States, only Iran, Kazakhstan and the Russian Federation authorise commercial catch of the species. During the past decade, annual catches of *A. nudiiventris* have increased in Iran (from 1.9 tonnes (t) in 1990 to 3.5 t in 2000) and in Kazakhstan (from 12 t in 1990 to 23.8 t in 2000). International trade in 1998 totalled 1,156 kg of *A. nudiiventris* caviar imports comprising: 1,004 kg from Iran, 87 kg from Kazakhstan, and 65 kg from the Russian Federation, 42% of which was declared as pre-Convention stock. No export quotas were established for specimens of *A. nudiiventris* for 1998. Caviar export quotas set by range States increased for Kazakhstan from 1,500 kg for 1999 to 2,200 kg for 2000. The Russian Federation *A. nudiiventris* caviar exports amounted 3,000 kg in 1999, that were probably re-exports from Kazakhstan, exceeding the 1999 export quota (1,500 kg) by 100%. Turkmenistan established their first export quota at 400 kg of caviar for 2000. The latter contradicts the fact that commercial catch of the species is prohibited in Turkmenistan waters and that *A. nudiiventris* has been included in the country's Red Data Book.

The species is recommended under Decision 11.106 for inclusion in category 1 (i).

DISTRIBUTION AND POPULATION

The CITES database lists the distribution of *Acipenser nudiiventris* as Armenia, Azerbaijan, Bulgaria, Georgia?, Hungary (ex), Iran (Islamic Republic of), Kazakhstan, Moldova, Republic of, Romania, Russian Federation, Slovakia (ex), Turkey, Turkmenistan?, Ukraine, Uzbekistan (ex) (Anon., 2000a).

A. nudiiventris is classified as Endangered by IUCN (1996):

EN A1acde+2d Azerbaijan, Bulgaria, Hungary, Iran, Kazakhstan (ex), Moldova, Romania, Russian Federation, Ukraine, Uzbekistan (ex) [River Danube]

Black Sea stock: EN A1acde+2d Russia, Ukraine [Mediterranean and Black Sea]

Caspian Sea stock: EN A1acde+2d Azerbaijan, Iran, Kazakhstan

Aral Sea Stock: EX Kazakhstan, Uzbekistan [Mediterranean and Black Sea]

River Danube population: CR A1cd+2cd Hungary, Romania.

Two morphs of the species have been identified, one anadromous and one sedentary. The anadromous morph has been recorded from the Sea of Azov and the Caspian, Black and Aral Seas and enters associated river systems to spawn. *A. nudiiventris* has always been less abundant than other sturgeon species and historically contributed to approximately 1% of the total sturgeon catch from the Caspian Sea. The species has always been rare in the Black Sea and the Sea of Azov (Sokolov and Vasil'ev, 1989). There were originally two isolated migratory populations in the Caspian Sea basin which ascended the Ural and Volga rivers respectively, for spawning (Makarov *et al.*, 1991a; Sokolov and Vasil'ev, 1989), however only the Ural River population is thought to remain in the Caspian Sea Basin (Avetissov, 1992). The non-migratory morph occurs in the Danube and some Ukrainian rivers (Banarescu, 1964; Holèik, 1995; Manea, 1966; Pavlov, 1980). There are no estimates of the total population size for this species, but Avetissov (1992) believes that current stocks in all rivers, except the Ural River, are on the verge of extinction. The historic range of the species included Austria (Fitzinger and Heckel, 1863) and Slovakia (Hensel and Holèik, 1997; Holèik, 1995).

Armenia: No current distribution or status information available.

Azerbaijan: In the past, the species entered the Kura River for spawning (Makarov *et al.*, 1991b; Sokolov and Vasil'ev, 1989). This spawning population used to be the largest for this particular species, however, during 1983-1987, the migrating population was estimated to number only 66-112 individuals and by the end of the 1980s, only one to four individuals passed through the spawning sites (Makarov *et al.*, 1991b). The entire existing stock is maintained by aquaculture, but this has failed to restore the former abundance (Makarov *et al.*, 1991b). In addition, the Lenkoranka and Astara River systems are used by a small number of spawning individuals (Makarov *et al.*, 1991b; Sokolov and Vasil'ev, 1989).

Bulgaria: *A. nudiventris* occurs in the Danube River and was listed in the Red Data Book of Bulgaria in 1985 as Critically Endangered (Karapetkova *et al.*, 1995).

Former Czechoslovakia: The former range included Slovakia (Hensel and Holèik, 1997; Holèik, 1995).

Georgia: The Rioni River provided the main spawning sites in Georgia, although the species was rarely found in this area. In addition, the south east coast of the Black Sea is an important feeding and wintering area for several species of sturgeon (*Huso huso*, *Acipenser stellatus*, *A. nudiventris*, *A. persicus colchicus* and *A. sturio*) that migrate upstream in a number of rivers to spawning grounds. The main rivers involved are the Supsa, Inguri, Chorokhi and particularly the Rioni (Zarkua and Tsuladze, 1999).

Hungary: *A. nudiventris* is considered to be very rare in the Danube and Tisza rivers (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 10 September 2000), with the last specimen in the Drava River recorded in 1989 (Pintér, 1991).

Iran: *A. nudiventris* migrates to the Sefidrud, Tajen and Babulrud Rivers on the Iranian coasts of the Caspian Sea. Sturgeon hatcheries in Iran apparently obtain breeders of *A. nudiventris* and other species from these rivers (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). However, Dr J. Holèik (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) believes the species no longer enters the Sefidrud rivers to spawn due to construction of the Mangil Dam in 1961. According to Dr M. Pourkazemi (*in litt.* to IUCN/SSC Wildlife Trade Programme, 25 September 2000) among the Mangil Dam and other dams constructed on the Sefidrud River, there is a remaining distance of 45 km from the dam closest to the estuary, and sturgeon breeders are still observed entering the river to spawn.

Kazakhstan: Historically, *A. nudiventris* entered the Ural River to spawn (Makarov *et al.*, 1991b; Sokolov and Vasil'ev, 1989) and from 1978-1990, the number of spawning individuals was estimated at 1,500 –18,600 (Avetissov, 1992). A winter migrating race of the species occurred in the Aral Sea Basin but is believed to have been extinct since the 1970s (Zholdasova, 1997). A population introduced from the Aral Sea occurred in the Ili River, which flows into Lake Balkash (Sokolov and Vasil'ev, 1989). However, even this population is now believed to be extinct because the lake is drying out (Bond *et al.*, 1992).

Moldova: The Moldovan range on the Danube River is 811 m and sturgeon also occur in the Prut River, a tributary of the Danube River (junction at km 137th) (Anon., 2000b). There have been no records to confirm the presence of *A. nudiventris* for 15 years (Dr V. Lobchenko, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Romania: Previously, only the sedentary morph occurred in the Danube and its tributaries. *A. nudiventris* is thought to never have been abundant in the Danube, and is now very rare and considered Critically Endangered in the Danube Basin (Hensel and Holèik, 1997).

Russian Federation: In the past, *A. nudiventris* entered the Volga to spawn (Makarov *et al.*, 1991b; Sokolov and Vasil'ev, 1989). Only the Ural River spawning population is thought to remain.

Turkey: No current distribution information available (see Threats to Survival and Domestic Use).

Turkmenistan: *A. nudiventris* is close to extinction in the Amu-Darya River and is very rare in the Caspian Sea (Dr Salinkov, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). It is listed in Category 1 of the 1999 Red Data Book of Turkmenistan, which means it is classified as a “Vanishing species that cannot be saved without special measures” (Salnikov, 1999).

Ukraine: Bacalbasca-Dobrovici reports that *A. nudiventris* is critically endangered or extinct in the Ukraine (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). It has been recorded in the Danube and Karadag regions of the Black Sea coast, very rarely near the mouth of the Dnieper River, and is not known in the main Dnieper, Dniester or Bug rivers. It has always been very rare and is not of commercial value in this area (Pavlov, 1980).

Uzbekistan: *A. nudiventris* is apparently extinct in Uzbekistan, following the drying of the Aral Sea (Birstein, 1993).

Former Yugoslavia: *A. nudiventris* is thought to be critically endangered or extinct in the Danube and the Iron Gate II reservoir (Bacalbasa-Dobrovici, 1997) and is classified as Very Rare and Endangered upstream the Djerdap I dam (Jankoviè, 1994).

HABITAT AND ECOLOGY

The maximum weight of *A. nudiventris* is around 120 kg. Male sexual maturity is reached at 6-12 years and in females at 12-18 years. Males then spawn at 1-2 year intervals and females at 2-3 year intervals (Hochleithner and Gessner, 1999). *A. nudiventris* may reach a total length of approximately 1.70-2.03 m, and its maximum recorded age is 36 years (Sokolov and Vasil'ev, 1989).

A. nudiventris was most abundant in river mouths, as its preferred habitat is shallow fresh water with muddy bottoms. It is a benthic feeder with a diet of fish, molluscs, insect larvae, crustaceans and occasionally the eggs of other sturgeon species (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000; based on reports from Shahid Marjani hatchery). According to studies conducted in Iran, sexual maturity of this species in the southern shores of the Caspian Sea is attained at average ages of 17 years (from 13 to 22) for females and 13 years (from 11 to 15) for males (International Sturgeon Research Institute, 1998; cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Based on catches and caviar production in the late 1970s, the rate of caviar/catch was estimated at 7% in weight for the three main commercial species of the northern Caspian Sea, *H. huso*, *Acipenser gueldenstaedtii* and *A. stellatus* (Doroshov and Binkowski, 1985; cited in Williot and Bourguignon, 1991).

Kazakhstan: Until 1993, the mouth of the Ural River was dredged annually to allow vessels to travel upstream to Atyrau (former Gur'yev) harbour. Since 1993, this annual management of the river channel has not been carried out and is an obstacle to sturgeon migration (Anon., 2000b).

THREATS TO SURVIVAL AND DOMESTIC USE

Since the 1950s, almost all spawning rivers for this species, with the exception of the Ural River have been dammed for hydroelectric power production and almost all the spawning grounds of the Sea of Azov, the Caspian and the Black Sea region have been lost. Consequently, the species' range has been drastically reduced. In addition, pollution has been rapidly increasing since 1979, the main sources being oil, industrial waste and agricultural chemicals and almost all spawning rivers are now heavily polluted (Anon., 1997).

In the former Soviet Union and Iran, strict sturgeon fishery legislation had been in effect for decades until the emergence of the three new independent States in the Caspian Sea basin in 1992 (Azerbaijan, Kazakhstan and Turkmenistan) (De Meulenaer and Raymakers, 1996). Current information suggests that amendments of previous regulations and/or adoption of new legislation are now in effect in most new range States in the Caspian Sea, as well as in the Black Sea (including the Danube River) and the Sea of Azov basins. In the latter, Bulgaria, Romania and Ukraine are the most important sturgeon fishing nations. However, enforcement

measures appear to be lacking and numerous experts as well as government officials have reported an increasing pressure of illegal fishing practices and criminal activities surrounding the caviar trade in much of the range (Anon., 2000b; Dobbs, 1992; Doward, 2000; Evtouchenko, 1997; King, 1998; Ward, 2000a and 2000b).

Azerbaijan: The Mingechaur Dam constructed on the Kura River prevents sturgeon from accessing spawning grounds higher up the river (Birstein, 1993). *A. nudiiventris*, particularly the population of the Kura River, has always been less abundant than other sturgeon species. The share of *A. nudiiventris* has never exceeded 2% of the total sturgeon catch in Azerbaijan (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Bulgaria: Wild sturgeon populations have been threatened by a decrease in the number of Danube River spawning grounds as a result of human development, particularly hydroelectric facilities. The long lasting unregulated fisheries have disrupted the age composition of the populations due to the selective catch of sexually mature fish and to the incidental catch of young sturgeon in nets set to catch other species when the juveniles migrate to the sea. Water pollution has also influenced the metabolism of the Ship Sturgeon (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Georgia: During the fishing seasons, particularly in the coastal anchovy fisheries, numerous sturgeon fry were caught as by-catch and have perished. The estimated number of sturgeon (no species-specific data) declined from 77,000 individuals in 1977, to 57,000 in 1980 and to 27,000 in 1986. During the 1990s, illegal fishing in Georgia expanded rapidly as a result of poor control (Zarkua and Tsaladze, 1999).

Hungary: Threats to *A. nudiiventris* are unknown because of the rarity of the species and very few recent records. However, the species is protected, with domestic use and trade being prohibited (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Commercial catch of *A. nudiiventris* is allowed in Iranian waters.

For the past 5 years, the Iranian domestic market of caviar and sturgeon meat is estimated to be 5% and 65% of the country's total production respectively (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Annual catch of *A. nudiiventris* (in tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. nudiiventris</i>	1.9	22.4	19.0	17.5	17.3	15.7	16.6	13.5	19.4	21.0	3.5
Total	2,296	2,315	2,058	1,462	1,380	1,216	1,310	1,043	1,127	923	182.
	.3	.1	.0	.1	.9	.3	.0	.5	.9	3	4

* Catch in spring 2000 only.

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000.

Sturgeon meat and caviar production of *A. nudiiventris* (tonnes)

	1992				1999			
	Meat	% *	Caviar	% *	Meat	% *	Caviar	% *
<i>A. nudiiventris</i>	15.2	0.9	1.5	0.6	16.8	2.3	2.0	2.0
Total	1,604.2		262.3		730.2		99.3	

* Percentages relate to the share of *A. nudiiventris* products of the total Iranian sturgeon fisheries.

Source: Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000

Kazakhstan: The Ural River is susceptible to oil pollution from two oil fields (Sagers, 1994) Both tributaries of the Aral Sea Basin have been altered for irrigation, consequently, the Aral Sea has decreased to 60-70% of its former volume (Ellis, 1990).

Annual commercial catch of *A. nudiiventris* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. nudiiventris</i>	12.0	12.0	19.3	58.0	26.0	22.0	26.0	27.1	40.0	26.0	23.8
Total	1,963.0	1,767.0	1,437.3	1,109.0	556.0	573.0	367.8	389.5	490.0	279.0	230.5

* Spring catch

Source: CITES Management Authority, *in litt.* to TRAFFIC Europe, 21 September 2000

Moldova: The main threats are dam construction and poaching (Vedrashko *et al.*, 1998).

Russian Federation: The level of domestic consumption of sturgeon products in the Russian Federation is unknown, but is said to be the main outlet of illegal sturgeon fisheries (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). Based on the number of sets of illegal fishing equipment as recorded by enforcement agencies of the Caspian Sea and the Sea of Azov, the illegal sturgeon catch was estimated at six to ten times the legal catch (Anon., 2000b). In 1997, the estimated volume of the Moscow sturgeon meat market was three times the annual catch quotas for sturgeon species (Anon., 1998).

Turkey: The Yesilirmak River has changed dramatically due to the construction of two dams in 1979, the lowest of which limits upstream migration. The Kizilirmak River was dammed in 1998 and suffers significant amounts of pollution. Spawning ground has been drastically reduced (Edwards and Doroshov, 1989). From 1969 to 1979, Turkish annual sturgeon landings ranged from 10-310 t, with a yearly average of 112 t (Edwards and Doroshov, 1989). In 1989 in Samsun, a Turkish landing site on the Black Sea, a caviar processor claimed that his business had dropped from 8 tonnes/year of caviar of all sturgeon species between the 1940s and the 1970s to 200 kg/year at the end of the 1980s. In 1989, sturgeon were caught as by-catch by trawlers that operated legally along the Turkish coast at a distance greater than three nautical miles from the shore, but also illegally within the three nautical mile limit. This represented the main fishing pressure on sturgeon. In April 1989, *A. stellatus*, *A. gueldenstaedtii*, *H. huso* and *A. sturio* were observed in Samsun. The investigators were told that *A. nudiiventris* was also landed in Samsun (Edwards and Doroshov, 1989).

Ukraine: Catch of *A. nudiiventris* in Ukrainian waters is prohibited since the species was listed in the Red Data Book of Ukraine in 1992 (Anon., 2000b).

INTERNATIONAL TRADE

The listing of *A. nudiiventris* in CITES Appendix II entered into effect on 1 April 1998. Data for this species were therefore limited to nine months of trade (April-December 1998).

Gross exports and the comparative tabulation of trade in *A. nudiiventris* are given in the Appendix. According to 1998 CITES Annual Reports, imports of *A. nudiiventris* amounted to 1,156 kg of caviar that originated from three range States: Iran, Kazakhstan and the Russian Federation. Of this, 42% was declared of pre-Convention stock.

Exports of *A. nudiiventris* from Russia in 1999 (kilogrammes)

	Quantity (kg)	Description	Importer	No of permits
<i>A. nudiiventris</i>	3,000.00	Caviar	US, FR, CH	3 Re-export

Source: CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 Sept. 2000

For *A. nudiiventris*, range States have established export quotas for caviar only.

Caviar export quotas for 1998, 1999 and 2000 (kilogrammes)

	1998	1999	2000	
	Caviar	Caviar	Caviar	Meat
Kazakhstan		1,500	2,200	34,000
Turkmenistan			400	
Total		1,500	2,600	34,000

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Illegal trade: Confiscation of *A. nudiventris* shipments were not reported in 1998, however poaching and smuggling are closely related and have been intensively reported in the media of range States and importing countries (Mc Donald, 2000; Snyder, 2000). However, a great deal of illegally caught sturgeon is destined to the domestic market, particularly meat (Anon., 1998).

CONSERVATION MEASURES

A. nudiventris is not fully protected by legislation in the Caspian region.

Bulgaria, Romania, Russian Federation: Catch is prohibited by the Convention Concerning Fishing In The Black Sea. Bacalbasa-Dobrovici believes that measures which have been put in place are too late to aid species recovery for this species (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Azerbaijan: The State institution responsible for sturgeon fishery and processing matters is the "State Concern Azerbalyg". Commercial catch of the species is prohibited (Anon., 2000b). The species was listed in the Red Data Book of Azerbaijan in 1994. Three hatcheries were built in the Lower Kura River by the authorities of the former Soviet Union. The total annual capacity was 1-2 million fingerlings which were to be released into the river. World Bank funds have been allocated for the construction of a new sturgeon hatchery that will have the capacity of 15 million fry/year and should be operational in the next couple of years. Concerns have been raised on the availability of adult sturgeon to constitute the broodstock of the farm (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Estimated number of fry released (million fish)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. nudiventris</i>	0	0.738	0	0	0	0	0	0.123	0.410	1.120	0.870
Total	17.52	9.082	2.980	1.838	1.142	1.242	4.070	6.065	6.220	20.29	16.96
		3								0	5

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

The Intergovernmental Commission for Caspian Biological Resources set a zero catch quota for the species in Azeri waters (Anon., 2000b).

Bulgaria: The fishing of *A. nudiventris* has been banned since 1985, when the species was included in the country's Red Data Book (Anon., 2000b). Conservation measures at the national level are needed, but they must be linked to the co-ordinated activities of all countries in the Black Sea region (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Hungary: The species is protected according to Decree No. 1/1982 (III.15.) OKTH on the Protected and Strictly Protected Species of Flora and Fauna, Value of their Specimens, Determination of the Range of Protected and Strictly Protected Caves and Exemptions from Restrictions and Prohibitions Set for Certain Protected Animal Species, last amended by the 15/1996 (VII.26.) Decree of the Minister for Environment. The Nature Conservation Act No. 53 of 1996, Article 43 orders that for all protected species, prior authorisation from the National Park Directorate is required for the: capture, killing, possession, captive breeding, introduction of specimens from exotic population, artificial exchange of genetic matter between populations, exchange, sale and purchase of *A. nudiventris* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Biometric measurements for all species have been carried out at all sturgeon catch stations for the past 10 years. The age, length, sex and weight of fish landed are recorded. The population structure is estimated on the basis of age groups. As a result of these studies, the number of catch stations were decreased by half, changes were adopted on sturgeon catch methods, and lower export quotas were set for the past five years.

Annual release of *A. nudiventris* fry (in million fish)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<i>A. nudiventris</i>	0	0	0	0	0	0	0.10	0.23	0.68	0.30
Total	4.56	6.60	3.45	4.17	5.91	9.13	12.35	21.63	24.56	19.1

0

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

Catch of *A. nudiventris* has been restricted in Iranian waters and enforcement of the legislation has been strengthened. Use of gillnets to catch *Rutilus* spp. Roach has been prohibited as sturgeon are caught as bycatch. There has been a forced and drastic reduction in fishing effort for *Rutilus* spp. Fishermen are encouraged to offer incidental catches of all sturgeon species to the Iranian Fisheries authorities for use in their artificial breeding programmes. A restocking programme has produced over one million fingerlings that were released in the rivers leading to the Caspian Sea during 1998-1999. A broodstock shoal of about 3,000 breeders and a gene bank have been established. Egg removal by caesarean has also been practised (Dr. M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Commercial sturgeon catch is only allowed in the tributaries of the Caspian Sea, but scientific catch is permitted in the open sea with an annual quota of 20 t.

Sturgeon fisheries are regulated by law. There are two fishing seasons, one during spring spawning and one during migration for hibernation. The dates of the fishing seasons vary with the climate and are therefore set annually by authorised scientific-research organisations. As with most former Soviet Republics, "sweep net" (bottom nets that are hauled several times a day) with a minimum mesh size and being marked with the name and address of the owner, is the only legal fishing gear for sturgeon. Each fisherman can only set one net at a time. A fishing license is required. Sturgeon fishing grounds (called "tonia") are limited; there are twelve on the Ural River and eight on the Kigach River. The minimum catch size for *A. nudiventris* is 105 cm. The ratio of males and females caught is not regulated or monitored. The purchase price at the landing site is based on a fixed legal rate of caviar weight extracted per female. *A. nudiventris* is set at 18.6% of the total body weight. The annual commercial catch quotas allocated to Kazakhstan by the Intergovernmental Commission for Caspian Biological Resources that meets annually in Astrakhan.

Annual commercial catch quotas allocated to Kazakhstan (tonnes)

	1996	1997	1998	1999	2000
<i>A. nudiventris</i>			40	50	35
Total	375	480	460	405	371

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 Sept. 2000

Two hatcheries exist in Kazakhstan; Uralo-Atyrau and Atyrau. These have been operational since 1998.

Annual release of *A. nudiventris* fry (number of fingerlings)

	1998	1999	2000
<i>A. nudiventris</i>		729,000	277,800
Total	1,706,193	5,318,471	5,507,160

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 Sept. 2000

Russian Federation: For Caspian Sea populations, the former Soviet Republics set annual catch quotas in agreement with Iran. Sturgeon catch quotas have been established by the Intergovernmental Commission for Caspian Biological Resources for many years. The annual share of sturgeon catches of each former USSR Caspian range State as adopted by the Commission since 1993 are as follows: the Russian Federation 70%, Kazakhstan 17.6%, Azerbaijan 6.1% and Turkmenistan 6.3%. There are two fishing

seasons in the Caspian Sea waters: from ice melting to 5 June, and from 1 September until the time of ice formation. A license is required to catch sturgeon in Russian waters. The minimum size limit for Ship Sturgeon caught in Russian waters is 105 cm. Russian catch quotas are declared by the Federal Government following the recommendations given by the State Fisheries Committee in consensus with an independent council of experts.

Export quotas established by the CITES Management Authority consist of 90% of the caviar production, which in turn depends on the catch quotas set for each species of sturgeon (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). The annual catch quotas or Total Allowable Catches (TACs) allocated by the State Fisheries Committee, are based on the recommendations of scientific agencies such as the Russian Federal Research Institute of Fisheries and Oceanography ("VNIRO") and the Pacific Research Institute of Fisheries and Oceanography ("TINRO"). The TAC includes the commercial catch, the scientific catch, and the catch of mature males and females destined to captive breeding.

There has been no release of *A. nudiventris* fry by Russian hatcheries in the Volga River. Recent data on the monitoring of catch per species, including the size composition and sex ratio of fish caught, are not available. The domestic consumption of sturgeon products in the Russian Federation, mostly meat and caviar, is unknown (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000).

Moldova: All sturgeon fisheries are prohibited by the Law of the Republic of Moldova on Fund of Natural Areas Protected by State (1998) (Anon. 1999).

Turkey: In 1979, with the exception of *H. huso* Beluga of more than 140 cm long, the fishery for sturgeon was prohibited in Turkish waters (Edwards and Doroshov, 1989). Circular 34/1, effective from 2000-2002, prohibits the collection of Acipenseriformes (CITES Management Authority of Turkey, *in litt.* to TRAFFIC Europe, 20 September 2000).

Turkmenistan: *A. nudiventris* was included in the country's Red Data Book in 1999 (Anon., 2000b). Commercial catch is prohibited but there are no surveys or management plans for this species (Dr. P. Salinkov, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Ukraine: Catch of *A. nudiventris* in Ukrainian waters is prohibited since the species was listed in the Red Data Book of Ukraine in 1992 (Anon., 2000b).

CAPTIVE BREEDING

In the former Soviet Union *A. nudiventris* has been bred in captivity since the 1960s in order to maintain the spawning stocks for commercial harvests of the Sea of Azov and the Caspian and Black seas. An average of 0.8 million fry were released annually into the southern Caspian Sea at the end of the 1980s. Most hatcheries of Azerbaijan, Kazakhstan and the Russian Federation, stopped being operational in the early- to mid-1990s following shortage of government funds (De Meulenaer and Raymakers, 1996).

Captive breeding is not occurring at present as there are no mature individuals available to create a broodstock. M. Hochleithner (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) notes that restocking projects for the non-migratory form of the species are planned.

Georgia: Up to the early 1990s, artificial breeding of sturgeon was carried out in the Varistkhi Sturgeon Plant built in 1983 on the Rioni River with mixed results (Zarkua and Tsuladze, 1999).

Hungary: There are no breeding facilities in Hungary (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Kazakhstan: The two hatcheries, Uralo-Atyrau and Atyyrau, have been operational since 1998, and restocking of fry has restarted (CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000).

REFERENCES

- Anon. (1997). Document Doc. 10.89; Prop. 10.65. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Anon. (1998). *Sturgeon catch and trade in the Russian part of the Caspian Sea*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, Pp. 21.
- Anon. (1999). *The Law of the Republic of Moldova on Fund of Natural Areas Protected by State (1998)*. In: "Ekologicheskoe Zakonodatel'stvo Respubliki Moldova (1996-1998)". Chisinau, BIOTICA Publ., 1999. Pp. 85-209.
- Anon. (2000a). <http://www.cites.org/eng/dbase/fauna>
- Anon. (2000b). *Sturgeon fisheries management and trade control measures in the Caspian Sea and Black Sea/ Sea of Azov range States*. TRAFFIC Europe field investigations, December 1999-January 2000. Unpublished.
- Avetissov, K.B. (1992) The present status of the Ship Sturgeon *Acipenser nudiventris* within its distribution area. In: A.P. Ivanov (ed). *Reproduction of Acipenserids, Salmonids and some less valuable fish*. VNIRO, Moscow. Pp. 3-15.
- Bacalbasa-Dobrovici, N. (1997). Endangered migratory sturgeons of the lower Danube River and its delta. *Environmental Biology of Fishes* 48: 201-207.
- Banarescu, P. (1964). Pisces-Osteichthyes. Fauna Republicii Populare Romine. Vol. 13.
- Birstein, V.J. (1993). Sturgeon and Paddlefishes: threatened fish in need of conservation. *Con. Biol.* 7(4):773-787.
- Bond, A.R., Micklin, P.P. and Sager, J. (1992). Lake Balkhash dwindling, becoming increasingly saline. *Post-Soviet Geography*, 33(2):131-134.
- De Meulenaer, T. and Raymakers, C. (1996). *Sturgeons of the Caspian Sea and the international trade in caviar*. TRAFFIC International, iv+ 71 Pp.
- Dobbs, M. (1992). The Coming Caviar Crisis: Overfishing, Pollution Endanger Sturgeon. *The Washington Post*, 30 May 1992.
- Doward, J. (2000). Roe rage as piracy rules. *The Guardian*, 21 May 2000.
- Edwards, D. and Doroshov, S. (1989). *Appraisal of the Sturgeon and Seatrout Fisheries and Proposals for a Rehabilitation Programme*. FAO Field Document FI.TCP/TUR/8853. Pp. 38.
- Ellis, W.S. (1990). The Aral: A soviet sea lies dying. *National Geographic* 177(2):71-92.
- Evtouchenko, A. (1997). [Au Dagestan une explosion fait 54 morts. Accusés: les gangs du caviar. *Courrier International*, N° 323 (9-15 January 1997)] (in French)
- Fitzinger, L.J., and Heckel, J. (1863). Monographische Darstellung der Gattung Acipenser. *Ann. Wiener Mus.Nat.* 1:269-326.
- Hensel, K. and Holèik, J. (1997). Past and current status of sturgeon in the upper and middle Danube.: In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 185-200.
- Hochleithner, M. and Gessner, J. (1999). *The Sturgeon and Paddlefishes (Acipenseriformes) of the World: Biology and Aquaculture*. AquaTech Publications, Kitzbühl. Pp. 165.
- Holèik, J. (1995). [1. Druh Acipenser (*Lioniscus*) *nudiventris* Lovetzky, 1828. Pp. 379-381. In: V.Baruš and O.Oliva (eds.) *Mihulovci Petromyzontes a ryby Osteichthyes*. 1. Fauna ÈR a SR 28/1. Academia, Praha.] (in Czechoslovakian).
- IUCN. (1996). 1996 *Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- Jankoviè, D. (1994). Chapter V, Ichthyofauna. In: *The Danube in Yugoslavia*. Eds. Jankoviè, D. and Yovicic M. Belgrade, Pp. 131-148.
- Karapetkova, M., Zhivkov, M. and Pchelarov, T. (1995). *Ribite v B'lgariya*. Geya Libris, Sofia. Pp. 247
- King, R. (1998). [Por un puñado de caviar. *El Mundo*, 13 December 1998.] (in Spanish)
- McDonald, L. (2000). Fish that laid the golden egg faces extinction. *Daily Express* (UK), 26 July 2000.
- Makarov, E.V., Rekov, Y.I. and Balandina L.G. (1991a). Current state of populations of Azov sea sturgeons. In: Williot, P. (ed.), *Acipenser*, Cemagref Publications. Pp. 69.
- Marakov, I.A., Alekperov, A.P. and Zarbalina, T.S. (1991b). Present status of the spawning run of Ship Sturgeon *Acipenser nudiventris*, in the Kura River. *J. Ichthyol.* 31(5):17-22.
- Manea, Gh. (1966). Contributii in studiul sturionilor din apele Romiiniei si al reproducerii lor in legatura cu constructiile hidroenergetice pe DFunarea inferiara. 1.Une aspecte ale biologici sturionilor. *Buletinul Institutului de Cercetari Piscicole* 25: 62-86.

- Pavlov, P.I. (1980). *Ribi*. Fauna Ukrainy. 8/1. Naukova Dumka, Kiiv. Pp. 336.
- Pintér, K. (1991). Sturgeon in Hungary, past and present situation. In: P. Williot (Ed.) *Acipenser*. Actes du premier colloque international sur l'esturgeon. Bordeaux, 3-6 October 1989. Pp. 173-178.
- Sagers, M.J. (1994). The oil industry in the southern-tier former soviet republics. *Post-Soviet Geography*, 35(5):267-298.
- Salnikov V. B. 1999. Ship *Acipenser nudiiventris* (Kessler, 1877). *The Red Data Book of Turkmenistan*. Volume 1. Invertebrate and Vertebrate Animals. Second edition, revised and updated. Ashgabat, Turkmenistan.
- Snyder, D. (2000). Md. Caviar Company Admits Smuggling. *The Washington Post*, 22 July 2000.
- Sokolov and Vasil'ev V.P. (1989). *Acipenser nudiiventris* Lovetsky, 1828. In: Holèik, J. (ed). The Freshwater Fishes of Europe. Vol. I/II: General Introduction of Fishes. Acipenseriformes. Wiesbaden, AULA-Verlag. In: Document Doc. 10.89; Prop. 10.65. 1997. Proposal to list all Acipenseriformes in Appendix II. Pp. 206-225.
- Vedrashko, A., Lobchenko, V. and Pirtsu, I. (1998). [The Problem of Conservation of the Acipenserids in the Lower Dniester. In: Problemele Conservarii Biodiversitatii din Cursul Medial si Inferior al Fluviului Nistru. *Abstr. Int. Conf. Chisinau*, 6-7 November 1998. Chisinau, BIOTICA Publ. Pp. 35-36.] (In Russian)
- Ward, P. (2000a). Azov Sturgeons Can't be Touched. *Crimean Time*, 18 August 2000.
- Ward, P. (2000b). Makhachkala: the Scale of Poaching in the Caspian has Reached Enormous Proportions. *Regions.Ru/Russia Regions*, 15 August 2000.
- Williot, P. and Bourguignon, G. (1991). Sturgeon and caviar production, current status and perspectives. In: Williot, P. (ed.), *Acipenser*, Cemagref Publications, Pp. 509-513.
- Zarkua, Z. and Tsuladze, V. (1999). Initiative needed to boost sturgeon numbers. *Eurofish* 3/99:40-41.
- Zholdasova, I. (1997). Sturgeons and the Aral Sea ecological catastrophe. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 373-380.

Gross exports of *Acipenser nudiiventris* 1998

TAXON	TERM	UNI	Exporte	1998
		T	r	
<i>Acipenser nudiiventris</i>	eggs		DE	44
<i>Acipenser nudiiventris</i>	eggs	kg	AE	330
<i>Acipenser nudiiventris</i>	eggs	kg	CH	21
<i>Acipenser nudiiventris</i>	eggs	kg	DE	601
<i>Acipenser nudiiventris</i>	eggs	kg	FR	154
<i>Acipenser nudiiventris</i>	eggs	kg	IR	510

Comparative tabulation of trade in *Acipenser nudiiventris* 1998

Year	Imp.	Exp.	Origin	Imports reported				Exports reported					
				Quantit	Unit	Term	P	S	Quantit	Unit	Term	P	S
				y				y					
1998	CH	AE	IR	330	Kg	eggs	T	W					
1998	FR	CH	IR	2	Kg	eggs	T	O	2	kg	eggs		O
1998	HK	CH	IR						9	kg	eggs		O
1998	MC	CH	IR	7	Kg	eggs	T	O	7	kg	eggs		O
1998	MX	CH	IR						2	kg	eggs		O
1998	CH	DE	RU	22	Kg	eggs	T	O	22	kg	eggs	T	O
1998	JP	DE	KZ						108	kg	eggs	T	O
1998	NO	DE	RU						6	kg	eggs	T	O
1998	US	DE	KZ	44		eggs	T	O					
1998	US	DE	KZ	43	Kg	eggs	T	O	465	kg	eggs	T	O
1998	US	DE	RU	43	Kg	eggs	T	O					
1998	CH	FR	IR	154	Kg	eggs	T	W					
1998	CH	IR		500	Kg	eggs	T	O					
1998	CH	IR		11	Kg	eggs	T	W					
Total:				1,156	Kg								

SUMMARY

The Sterlet *Acipenser ruthenus* is a freshwater species and is the smallest species of sturgeon. It reaches sexual maturity in just 3-8 years with an average life span of 15 years. It originally inhabited the rivers of Eurasia, being widely distributed in rivers flowing into the Caspian, Black, Baltic, White, Barents and Kara Seas and the Sea of Azov. In the Russian Federation, populations are now thought to be virtually extirpated from three Siberian rivers, Ob', Irtysh and Yenisey, being currently found mainly in the Volga River. The Danube is a stronghold for the species and in recent years, catches have been reported as far upstream as Austria.

Historically, the highest catches were recorded in the Danube River basin (63.5 tonnes (t) average annual catch from 1958 to 1981), and in the Siberian rivers (average annual catch in the 1990s of 69 t), while the Caspian Sea has yielded only 1-2 t annually since 1997. Nowadays, the main commercial catches are reported from the Danube River basin range States: Bulgaria, Romania and the former Yugoslavia. Recorded catches in the Russian Federation decreased by around 50% between 1990 and 1996, and commercial catch of the species has been prohibited in the Yenisey River. Re-stocking of the species is reported in Bulgaria, Hungary and the former Yugoslavia, but none in the Russian Federation. The majority of international trade is composed of live juveniles produced in aquaculture. In 1998, Hungary exported 90,000 and the Russian Federation 2,000 juveniles raised in aquaculture. Additionally, the Russian Federation exported 4 kg of eggs (probably fertilised eggs) in 1998. No export quotas were in place for live juvenile sturgeon in 1998, 1999 and 2000. Only the Russian Federation has established export quotas for *A. ruthenus* in 2000: 300 kg of caviar, 2 t of meat and a total of 125 kg of fertilised eggs for the species and three hybrids.

The species is recommended under Decision 11.106 for inclusion in category 2 (ii).

DISTRIBUTION AND POPULATION

The CITES database gives the distribution of *Acipenser ruthenus* as: Austria, Bosnia and Herzegovina, Bulgaria, Czech Republic, Georgia, Germany, Hungary, Kazakhstan, Latvia, Lithuania, Moldova, Republic of, Romania, Russian Federation, Slovakia, Slovenia, Switzerland, Turkey, Ukraine, Yugoslavia (Anon., 2000a).

A. ruthenus is classified as Vulnerable by IUCN (1996):

VU A1c Austria, Bosnia and Herzegovina, Czech Republic, Germany, Hungary, Kazakhstan, Latvia, Lithuania, Romania, Russian Federation, Slovakia, Switzerland, Ukraine, Yugoslavia

Caspian and Black Sea drainage stock: VU A1cd Hungary, Romania, Russian Federation, Ukraine, Yugoslavia [Mediterranean and Black Sea]

Russian stock: VU A1c Russia [Irtysh, Ob and Yenisei rivers]

Austria: *A. ruthenus* has a limited distribution in the Danube and Drava rivers (Honsig-Erlenburg and Schultz, 1989; Jungwirth, 1975; Schiemer and Spindler 1989, cited in Dr J. Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). It is recorded as Endangered (Anon., 1997).

Bulgaria: *A. ruthenus* occurs in the Danube River and previously, in the lower courses of its tributaries. However, population density and catch have decreased in recent years (Karapetkova *et al.*, 1995, cited in Dr J. Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Czech Republic: *A. ruthenus* occurs in the lower course of the Morava River (the western tributary of the Danube River), and its tributary, the Dyje River (Dr J. Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). The status of the species has not been evaluated (Lusk and Hanel, 1996b, cited in Dr J. Holèik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Germany: *A. ruthenus* has been extirpated (Dr J. Holcik, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Hungary: *A. ruthenus* is considered to be common (Pintér, 1991; A. Ronyai, *in litt.* to IUCN/SSC Wildlife Trade Programme, 8 August 2000). No information is available on population size, other than that it is quite common and occurs in all major rivers: Berettyó, Danube, Dráva, Körös, Maros, Mura, Rába, Szamos and Tisza (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Moldova: The Moldovan range on the Danube River is 811 m and sturgeon also occur in the Pruth River, a tributary of the Danube River (junction at km 137th) (Anon., 2000c). The species also occurs in the Dniester (or Nistru) River (Sokolov and Vasil'ev, 1989).

Romania: The range of *A. ruthenus* in the Danube River increased in the 1980s due to improved water quality, but has now diminished due to overfishing (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). The range is fragmented (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Russian Federation: In the Caspian Sea region, *A. ruthenus* occur throughout the course of the Volga River including reservoirs and in many of its major tributaries. The most recently published information concentrates on the Vyatka River population, which was reported to be abundant in the 1990s (Kuznetsov *et al.*, 1995). The species is rarely encountered in the Ural River. In the Sea of Azov basin, *A. ruthenus* was recorded from the Don and Kuban Rivers, but there are no current records.

In the White and Barents Sea drainage, *A. ruthenus* inhabited the Northern Dvina River basin, though there is no current information on its status. *A. ruthenus* also once inhabited the Ob', Irtysh and Yenisey Rivers in Siberia, but is now thought to be virtually extirpated from these (Birstein, 1993). Introduced populations occur in the Pechora and Western Dvina rivers (Anon., 1997).

Slovakia: Classified as Vulnerable, *A. ruthenus* occurs in the Danube River and its main tributaries, the Morava, Váh, Hron, Ipec and Tisa rivers. The species had a limited distribution until 1970 due to pollution, but improved water quality has coincided with increased distribution and catch (Holcik, 1995).

Ukraine: Listed in the Red Data Book of Ukraine (Gringevsky, 1995), *A. ruthenus* occurs in the lower Danube and the Dniester rivers where its distribution is limited since the construction of the Dubosary dam. On rare occasions it is found in the Dnieper and Bug River systems and some of their tributaries. Population density has decreased since the construction of the dam (Pavlov, 1980).

Former Yugoslavia: No information is available for Croatia. In Serbia, *A. ruthenus* occurs in the Danube and the lower parts of the Sava and Tisa rivers (Janković, 1993). In Slovenia, the species is considered Vulnerable and in decline (Povž and Sket, 1990).

Introduced populations:

Germany: In the Baltic Sea river estuaries, east of Rostock (Germany), and further upstream, specimens of non-indigenous sturgeon species and probably hybrids have been caught. They probably originate from upstream aquaculture farms and/or aquaria ("specimens that had grown too large to be kept in the private aquaria") (Jörn Gessner, Scientist at the Institute of Freshwater Ecology and Inland Fisheries, pers. comm. to TRAFFIC Europe, 15 March 2000).

Italy: *A. ruthenus* has been introduced in captive breeding facilities in Italy in 1990s (Agroittica Lombarda and Azienda Agricola, Italian sturgeon farmers, *in litt.* to TRAFFIC Europe-Italy, 1999). There is no documentation on the potential damage of the introduction of exotic Acipenseriformes on local species. If specimens of *A. ruthenus* escape into the Po River, they may threaten the Adriatic Sturgeon *Acipenser naccarii*, which is on the brink of extinction.

HABITAT AND ECOLOGY

A. ruthenus is the smallest of all sturgeon species, usually reaching a total length of 1 m and a weight of up to 6.5 kg (Anon., 1997). The maximum age is reported to be 27 years, with an average life span of 15 years. Sexual maturity is reached at 3-5 years in males and 5-8 years in females (Hochleithner and Gessner, 1999). It occurs in freshwater and prefers deep depressions in rivers with a stony, gravelly or sandy bed. It is a benthic feeder with a diet of insect larvae, small molluscs and invertebrates. It does not undertake long migrations. During the spawning season, *A. ruthenus* feed on sturgeon eggs of other species.

THREATS TO SURVIVAL AND DOMESTIC USE

The damming of virtually all rivers in the range of the species began in the 1930s and led to a sharp decline in spawning habitat. Spawning success is related to the water level which in turn is controlled by hydroelectric power stations; lowered water levels in rivers can cause mass death of fingerlings (Jankoviè, 1993). Reduced flow velocity has also led to degeneration in the reproductive system (Lukin *et al.*, 1981, cited in Anon., 1997). Increasing pollution (oil products, phenols, PCBs, mercury etc.) in the Volga River System and Siberian rivers in recent years has threatened the survival of *A. ruthenus* (Romanov and Altuf'ev, 1993). Studies of sympatric species have shown degeneration of the reproductive system, liver and muscle tissue as a result of pollution. Lukin *et al.*, (1981, cited in Anon., 1997) found that specimens in large water reservoirs of the Volga River grew unusually quickly but did not mature. The newly constructed Novodnestrovskaya hydro power station has lowered the spring and summer water temperatures to 5-6°C and has caused reproductive problems for *A. ruthenus* and other sturgeon in the Middle Dniester (Sarapanovskaia, 1999).

In the Danube River, from 1958 to 1981, catches of *A. ruthenus* ranged from 117 t (in 1963) to 36 t (in 1979) and averaged 63.5 t per annum for this 24-year period. The average share of these annual catches were distributed amongst the Danube countries as follows: former Yugoslavia averaging 57.5% of the total catch from 1958 to 1981, Bulgaria 21.8%, Romania 10.5%, Hungary 3.5% and the former Czechoslovakia 0.5% (Hensel and Holèik, 1997). Little data are available on recent and current catch data of *A. ruthenus* in the Danube River basin.

Bulgaria: Wild populations of sturgeon have been threatened by the decrease in the number of spawning grounds on the Danube River as a result of human development, particularly hydroelectric facilities. The long lasting unregulated fisheries have disrupted the age composition of the populations due to selective catch of sexually mature fish, and incidental catch of young sturgeon in nets set to catch other species when the juveniles migrate to the sea. Water pollution has also influenced the metabolism of the fish (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Annual catch of *A. ruthenus* (tonnes)

	1995	1996	1997	1998	1999
<i>A. ruthenus</i>	0.1	0.8	0.8	1.2	1.5
Total sturgeon catch	14.7	32.5	49.8	55.9	50.5
n° of licenses issued	0	0	122	155	169

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Domestic consumption of caviar in Bulgaria is limited to 7-10% of the total Bulgarian caviar production. All sturgeon meat production is consumed in Bulgaria.

Hungary: *A. ruthenus* is the only species of sturgeon commonly found in Hungary. Estimated annual catches range from 3-20 t (Pintér 1991; A. Ronyai, *in litt.* to IUCN/SSC Wildlife Trade Programme, 8 August 2000). Hungary is not involved in caviar production. From 1997 to 1999, the following catches were recorded: 1997- 14.2 t, 1998 - 9.5 t, 1999 - 34.9 t (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Romania: There are many small populations of *A. ruthenus* in the Danube River, but no records of national catch levels (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, 5 Sept. 2000).

Russian Federation: *A. ruthenus* catch has always been relatively small on the territory of the former Soviet Union, compared with that of major commercial sturgeon species (*Huso huso*, *A. gueldenstaedtii* and *A. stellatus*). This may be partly explained by the small size of *A. ruthenus*, but also because Sterlet caviar is not as valuable as caviar of other species. *A. ruthenus* has been primarily targeted by fishermen to supply the domestic demand (Anon., 2000b).

Catch of *A. ruthenus* (tonnes)

	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997
Caspian Sea	?	?	?	?	1	1	1	1	2	2
Water reservoirs of Russia	40.5	57.8	30.4	31.4	29.7	12.3	13.1	24.1	16.8	?
Ob' River	?	?	15.0	15.0	15.0	15.0	15.0	15.0	15.0	15.0
Yenisey River	39.2	28.9	63.3	57.2	56.0	64.4	61.9	58.2	42.0	25.9
Northern Dvina River	?	?	7.7	8.0	9.5	7.0	5.0	8.0	4.8	8.9
Total			116.	111.	111.	99.7	96.0	106.	80.6	
			4	6	2			3		

Source: Federal Research Institute of Fisheries and Oceanography of the Russian Federation ("VNIRO"), cited in Anon., 2000a

The level of domestic consumption of sturgeon products in the Russian Federation is unknown, but is said to be the main outlet of illegal sturgeon fisheries (Mr R.V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). Based on the number of sets of illegal fishing equipment recorded by enforcement agencies of the Caspian Sea and the Sea of Azov, the illegal sturgeon catch for all species was evaluated at six to ten times the legal catch (Anon., 2000b). In 1997, the estimated volume of the Moscow sturgeon meat market was three times the annual catch quotas for sturgeon species (Anon., 1998).

Ukraine: Catch of *A. ruthenus* in Ukrainian waters has been prohibited since the species was listed in the Red Data Book of Ukraine in 1992 (Anon., 2000c).

INTERNATIONAL TRADE

The listing of *A. ruthenus* in CITES Appendix II entered into effect on 1 April 1998. Data for this species are therefore limited to nine months of trade (April-December 1998).

Gross exports and the comparative tabulation of trade in *A. ruthenus* are given in the Appendix. CITES Annual Reports for 1998 suggest that live juvenile fish represented the majority of international trade in *A. ruthenus*. Hungary is the main exporter, exporting more than 95% of the total 92,000 live individuals in trade declared as "F" – First Generation). The European Union (EU) is the largest importer (Germany, the Netherlands, and the United Kingdom (UK)). International trade in live juvenile *A. ruthenus* supplies the sturgeon aquaculture industry and the ornamental fish (aquarium) industry. In Italy, for instance, *A. ruthenus* is imported for ornamental purposes (Dr P. Bronzi *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). According to 1998 Annual Reports most importing countries do not report live juvenile imports (see Appendix). Possible reasons for this omission are: reporting errors; non-implementation of the CITES requirement to record imports of "live" fish and fertilised "eggs" in CITES Annual Reports; and lack of awareness of the Wildlife Regulation (EC) 338/97 among importers in the EU and therefore lack of applications for import permits for live juvenile sturgeon.

The small size of the species makes it attractive to the apparently increasing meat market (see export quotas for 2000) and because restaurants prefer smaller sized fish.

Hungary is not involved in *A. ruthenus* caviar production and export (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Exports of *A. ruthenus* from Hungary in 1999

	China	USA	Germany	Slovenia	United Kingdom	Czech Republic	Bulgaria	Netherlands	Belgium
Live fingerlings		4,000	22,000	15,000	10,000	8,000	6,000	5,000	5,000
Fry		200,000							
Fert. eggs		30,000							

Source: CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000

Exports of *A. ruthenus* from Russia in 1999

	Quantity (kg)	Description	Importer	No of permits
<i>A. ruthenus</i>	16.80	Live eggs	DE, CN, US, HU, GR	7 Export
	200.00	Live fish	KR	2 Export
<i>Huso huso x A. ruth.</i>	62.80	Live eggs	PL, DE, JP, CN, LV, US, HU, IL, GR	16 Export
	300.00	Live fish	KR	1 Export

Source: CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 September 2000

Export quotas for *A. ruthenus* and its hybrids set by Bulgaria and the Russian Federation (kilogrammes)

	1998		1999		2000		Fertilised Eggs
	Caviar	Meat	Caviar	Meat	Caviar	Meat	
Bulgaria			30				
Russian Federation					300	2,000	5
<i>A. baerii x A. ruthenus</i>							15
<i>A. gueldenstaedtii x A. ruthenus</i>							15
<i>H. huso x A. ruthenus</i>							90
Total	0	0	30	0	300	2,000	125

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Note: The use of the term "eggs" and the unit "kg" or "g" (see # in Appendix and table on export quotas, above) to describe shipments of both caviar and fertilised eggs complicates the analysis of trade data. Similar difficulties arise from the unit used for export quotas of fertilised eggs, which is kilogrammes ("kg") instead of "number of specimens". Kilogrammes do not directly correspond to the number of specimens, and hence does not allow the effective evaluation of quotas. This confusion is also a source of mis-interpretation of data and is an obstacle to the proper monitoring of international trade, e.g. underestimation of live specimens in trade.

CONSERVATION MEASURES

There are no specific monitoring programmes for *A. ruthenus* in most range States.

Bulgaria: Conservation measures at the national level are needed, but they must be linked to the co-ordinated activities of all countries in the Black Sea region (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). In Bulgaria, sturgeon fisheries are prohibited for 30 days, from 20 April to 20 May which corresponds to the peak of the breeding season. The authorities may allow a 15-day extension of the closed season pending the natural conditions of that year. Explosives, poison, bottom trawling and dragging, as well as stunning substances and other destructive fishing devices are prohibited. The minimum size limit for *A. ruthenus* is 33 cm. Export quotas for all sturgeon species are set

on the basis of catch data and trends, as well as the biological characteristics of the species. A system has been established for the distribution of export quotas among export companies. These companies are obliged to restock the Danube River with sturgeon fry before they are authorised to export sturgeon specimens. For example, for a given year, for each kilogramme of caviar exported, the company must release 30 juvenile sturgeon (Anon., 2000c). The restocking programme is controlled by relevant government agencies. It is expected that the first released individuals will return to the Danube River in 2005-2007. Only 300 *A. ruthenus* fingerlings of 15 g have been released in Bulgarian waters (in the Danube River in 1998). There was no restocking for this species in 1999 or 2000 (Anon., 2000c).

Due to budgetary constraints, Bulgaria has not monitored the wild population of any sturgeon species in recent years. However, plans have been prepared for these activities (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Hungary: Mature and immature individuals of *A. ruthenus*, *A. baerii* and *A. gueldenstaedtii* are bred in captivity as broodstock. Fingerlings of *A. ruthenus* are released in Hungarian waters. The catch of the species is regulated by the Act on Fishery and Sport Fishing by length limitation and with closed season (A. Ronyai, *in litt.* to IUCN/SSC Wildlife Trade Programme, 8 August 2000). The closed fishing season operates from 1 March to 31 May. During the open season, fish less than 45 cm in length cannot be caught and the daily catch limit is three specimens per angler. *A. ruthenus* is not protected by law, but the introduction of hybrids into natural waters is prohibited by the *Nature Conservation Act* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Romania: In Romania, the Centre of Research for Fish Farming, Fishing and Fish Processing in Galati is monitoring the status (population sizes/trends) of sturgeon stocks upriver of the Danube Delta (Birstein, 1996). Bacalbasa-Dobrovici (*in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, September 2000) believes that fishing pressure should be reduced in the lower Danube. He reports that that the existing legislation pertaining to the region is neither adequate nor enforced, particularly in Romania.

A fishing license is required in Romania to capture this species (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, 5 September 2000). Sturgeon fishing gear, for instance the type of net and mesh size, is strictly regulated by Romanian law. The fishing season is set each year by Ministerial Order. A fishing license is required to catch any species of sturgeon in Romanian waters. No minimum size limit or catch quota has been established for *A. ruthenus*. The Romanian restocking programme has not released juvenile Sterlet into the Danube River (Anon., 2000c).

Russian Federation: In the late 1990s, sturgeon commercial catch in the Yenisey River (*A. ruthenus* and *A. baerii*) was prohibited due to rapidly declining populations and the extremely low rate of natural reproduction (Anon., 2000b), caused by inaccessible spawning grounds and impact of pollution on the physiology of fish (Ruban, 1997). In 2000, catch of *A. ruthenus* for scientific and monitoring purposes, as well as for local subsistence use, is limited to 5 t in the Yenisey River. In the Ob'-Irtysh rivers basin, the Total Allowable Catch (TAC – as calculated by VNIRO and accepted by the independent Council of experts) should not exceed 10 t (Anon., 2000b).

Export quotas established by the CITES Management Authority for Acipenseriformes of the Russian Federation consist of 90% of the total caviar production in the country. This in turn depends on the catch quotas set for each species of sturgeon (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). The TAC allocated by the State Fisheries Committee is based on the recommendations of scientific agencies such as VNIRO and the Pacific Research Institute of Fisheries and Oceanography (TINRO). The TAC includes the commercial catch, scientific catch and the catch of mature males and females destined for captive breeding.

There has been no release of *A. ruthenus* fry by Russian hatcheries in the Volga River (Anon., 2000b). Recent data on the monitoring of catch per species, including the size composition and sex ratio of fish caught are not available. The domestic consumption of sturgeon products in the Russian Federation, mostly meat and caviar, is unknown (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000).

Ukraine: Catch of *A. ruthenus* in Ukrainian waters is prohibited since the species was listed in the Red Data Book of Ukraine in 1992 (Anon., 2000c; Gringevsky, 1995).

Yugoslavia: It seems that *A. ruthenus* is permitted to be fished throughout the year. Certain types of fishing gear are prohibited to catch sturgeon by Yugoslavia law, and an agreement has been signed between Romania and Yugoslavia on the prohibition of gear that cause high mortality of untargeted fish. Any fishing activity in Yugoslav waters, including all commercial and sport fisheries, requires a license. The licenses are issued on a per district basis. The minimum size limit for *A. ruthenus* is set at 33 cm, but catches of the species have been prohibited from 1996 to 2000. A restocking programme is being carried out by the Centre for fish reproduction of the "Djerdap Fishing Company" in Malla Vrbica village. Data on sturgeon juvenile release have not been provided (Anon., 2000b).

CAPTIVE BREEDING

The species is produced in aquaculture in most of the warm water ponds in the Russian Federation, Ukraine, Hungary and Germany for domestic meat markets. The species has been bred in captivity since 1993 and stocking programmes are in place for the Danube and Drava rivers (M. Hochleithner, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

In the Dniester River, Moldova, limited quantities of *A. ruthenus* were bred in ponds for fishing from 1980 to 1988 (Lobchenko *et.al.*, 1996).

A. ruthenus has been introduced into aquaculture in Belgium (W. Verdonck, sturgeon farmer, pers. comm. to TRAFFIC Europe, July 2000), Italy (Agroittica Lombarda and Azienda Agricola, Italian sturgeon farmers, *in litt.* TRAFFIC Europe-Italy, 1999) and France (Welcomme, 1988). The size of the broodstock is not known. In European countries, a hybrid of *H. huso* and *A. ruthenus* called 'bester' is subject to aquaculture; it is fast growing and produces fertile and early maturing offspring (Burtsev, 1997).

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database, but according to trade data there is currently a significant production of *A. ruthenus* juveniles in commercial farms (see Appendix).

Bulgaria: A fish farm was established in 1997 in Boliartzi village (Plovdiv region). It is the only existing hatchery. All juveniles released in the Danube River in Bulgaria from 1998 to 2000 were produced there. The hatchery has a total capacity of 60,000 juveniles. Since 1999, attempts have been made to breed sturgeon in artificial water basins near Sofia (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Hungary: Through captive breeding and imports of fertilised eggs (from the Russian Federation), tens of thousands of fry and fingerlings are reared annually in Hungary. Most of them are exported, but some are kept until they reach a marketable size for the domestic market (A. Ronyai, *in litt.* to IUCN/SSC Wildlife Trade Programme, 8 August 2000). There are two major exporters of live fingerlings who have broodstock of *A. ruthenus*. These stocks consist of wild collected females and captive bred specimens. Other exporters do not have breeding stocks, but obtain eggs from wild caught females and fertilize them artificially. They export fertilised eggs, yolk sac fry and fingerlings. There are no sturgeon hybrids in captive breeding facilities (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Poland: The presence of fish farms and hatcheries of sturgeon in Poland was confirmed by the CITES Management Authority of Poland. The list of sturgeon species bred in captivity in Poland is as follows: *Acipenser baerii*, *A. gueldenstaedtii*, *A. ruthenus*, *A. stellatus*, *Huso huso*, *Polyodon spathula* and hybrids of some species. In most cases, hybrids are F1 generation live specimens (fertilised eggs) imported into Poland from range States, particularly from the Russian Federation. Some farms keep F2 generation specimens, bred in Poland and used as broodstock, partly for the production of juveniles to be sold on the domestic market and exported (Dr Z. Krzeminski, Deputy Director, *in litt.* to TRAFFIC Europe, 31 August 2000).

United States of America: Sturgeon are in the preliminary stage of captive breeding in Hawaii (USFWS, CITES Management Authority, *in litt.* to TRAFFIC Europe, September 2000), but no detailed information has been provided on the species concerned.

REFERENCES

- Anon. (1996). *A Guide to Interpreting Outputs from the WCMC CITES Trade Database*, Produced by the World Conservation Monitoring Centre, Cambridge UK. Pp. 16
- Anon. (1997). Document Doc. 10.89; Prop. 10.65. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Anon. (1998). *Sturgeon catch and trade in the Russian part of the Caspian Sea*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, Pp. 21.
- Anon. (2000a). <http://www.cites.org/eng/dbase/fauna>
- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia field investigations. Unpublished report, Pp. 23.
- Anon. (2000c). *Sturgeon fisheries management and trade control measures in the Caspian Sea and Black Sea/ Sea of Azov range States*. TRAFFIC Europe field investigations, December 1999-January 2000. Unpublished.
- Birstein, V.J. (1993). Sturgeon and Paddlefishes: threatened fish in need of conservation. *Cons. Biol.* 7(4):773-787.
- Birstein, V.J. (1996). Sturgeons in the lower Danube. *Sturgeon Quarterly* 4 (1/2):10-11.
- Burtsev, I.A. (1997). Bester in aquaculture. In: Birstein, V.J., Bauer, A. and Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 35-43.
- Gringevsky, M.V. (1995). Evaluation of the contemporary status of sturgeons in the Dnieper River basin. *The Sturgeon Quarterly*, 3 (1): 9.
- Hensel, K. and Holèik, J. (1997). Past and current status of sturgeon in the upper and middle Danube. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *The Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 185-200.
- Hochleithner, M. and Gessner, J. (1999). *The Sturgeon and Paddlefishes (Acipenseriformes) of the World: Biology and Aquaculture*. AquaTech Publications. Pp. 165.
- Holèik, J. (1995). 3.druh *Acipenser* (*Acipenser ruthenus* Linnaeus, 1758. In: V.Baruš and O.Oliva (eds.) *Mihulovci Petromyzontes a ryby Osteichthyes* (1). Fauna ĚR a SR 28/1. *ACADEMIA*, Praha, Pp.384-391.
- Honsig-Erlenburg, W. and N.Schultz. (1989). *The fishes of Carinthia*. Verl. Naturwiss. Verein f. Kärnten, Klagenfurt.
- IUCN. (1996). *1996 Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- Jankoviè, D. (1993). Populations of Acipenseridae prior and after the construction of the HEPS Derdap I and II. *Acta Biologica Iugoslavica, Seria E, Ichthologia* 25:29-34.
- Jungwirth, M. (1975). Die Fische in Niederösterreich. *Wiss. Schrift. Niederösterreichisch* 6:1-31.
- Karapetkova, M., M.Zivkov and T.Pchelarov, (1995): *Ribite v B'lgariya*. Geya Libris, Sofia. Pp. 247.
- Kuznetsov, V.A., Grekov, M.L. and Kas yanenko, E.B. (1995). Some ecological and morphological characteristics of the Sterlet, *Acipenser ruthenus*, from the Middle Vyatka River. *J. Ichthyol.* 35(9):8-19.
- Lobchenko V.V., Vedrashko A.I. and Pirtsu I.A. (1996). Conservation of Genofund of Middle Dniester Sterlet. In: *Management Ecologic si Dezvoltare Durabila: Abstr. Int. Conf.* Chisinau, 5-6 iunie, 1996. Chisinau, Pp. 105-106 [In Russian].
- Lusk, S. and Hanel, L. (1996a). Druhová diverzita ichthyofauny Ěeské republiky. *Biodiverzita Ictyofauny ĚR* (I):5-15.
- Lusk, S. and Hanel, L. (1996b). Ěervený seznam mihulí a ryb Ěeské republiky. *Biodiverzita Ichtyofauny ĚR* (I):16-25.
- Pavlov, P.J., (1980). Ribi. Fauna Ukrainy. 8/1. Naukova Dumka, Kiiv. Pp. 336.
- Pintér, K. (1991). Sturgeon in Hungary, past and present situation. In: P. Williot (Ed.) *Acipenser. Actes du premier colloque international sur l'esturgeon*. Bordeaux, 3-6 October 1989. Pp. 173-178.
- Povž, M. and B. Sket, (1990). Naše slatkovodne ribe. Založba Mladinska knjiga, Lubljana. Pp. 374.

- Romanov, A.A. and Aituf'ev, Y.V. (1993). Ectopic Histogenesis of Sexual Cells of Caspian Sea Sturgeon. *J. Ichthyol.* 33 (2):140-150.
- Ruban, G.I. (1997). Species Structure, Contemporary Distribution and Status of Siberian Sturgeon, *Acipenser baerii*. In: V.Birstein, J.R.Waldman, and W.E.Bemis (ed). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publishers, Dordrecht. Pp. 221-230.
- Sarapanovskaia T. (1999). [Ekologicheskie Problemy Srednego Dnestra. Chisinau. Pp.90.] (in Russian)
- Schiemer, F. and Spindler, T. (1989). Endangered fish species of the Danube River in Austria. *Regulated Rivers* 4:397-407.
- Sokolov, L.I. and Vasil'ev, V.P. (1989). *Acipenser ruthenus* Linnaeus, 1758. In: Holèik, J. (ed.) The Freshwater Fishes of Europe, Vol. 1, pt. II, General Introduction to Fishes, Acipenseriformes, AULA – Verlag, Wiesbaden. Pp. 227-262.
- Welcomme R.L. (1988). International introductions of inland aquatic species. *FAO Fisheries Technical Paper*. 294: 318 pp.

Gross exports* of *Acipenser ruthenus* 1998

TAXON	TERM	UNI	Exporter	1998
<i>Acipenser ruthenus</i>	eggs #	kg	RU	4
<i>Acipenser ruthenus</i>	live		CZ	0
<i>Acipenser ruthenus</i>	live		GB	300
<i>Acipenser ruthenus</i>	live		HU	89,700
<i>Acipenser ruthenus</i>	live		RU	2,000
<i>Acipenser ruthenus</i>	live	kg	RU	50
<i>Acipenser ruthenus</i>	specimens		DE	0

Comparative tabulation of *Acipenser ruthenus* 1998

Year	Imp	Exp	Origin	Imports reported				Exports reported					
				Quantit	Unit	Term	P	S	Quantit	Unit	Term	P	S
1998	JP	GB	HU						300		live	T	F
1998	BE		HU						5,000		live	T	F
1998	BG		HU						1,200		live	T	F
1998	DE		HU						40,000		live		
1998	DE		HU						2,000		live	T	F
1998	GB		HU						11,500		live	T	F
1998	NL		HU						20,000		live	T	F
1998	SI		HU						10,000		live	T	F
1998	KR		RU						2,000		live	S	F
Sub-total								92,000 Live fish					
1998													
1998	CN		RU						2 kg #	eggs		T	F
1998	GR		RU						2 kg #	eggs		T	C

* **Definition:** "Gross exports are the sum of all reported exports and re-exports in a particular commodity or species in a particular year or series of years. Gross trade is thus a simple measure of the total number of items recorded in international trade. However, gross trade may be an overestimate of the total number of actual specimens in trade as re-exports are not deducted from the total." (Anon., 1996).

This is particularly applicable to caviar of which shipments are often exported, re-packaged and re-exported. For instance, in the mid-1990s, Germany re-exported as much as 45% of its annual caviar imports (De Meulenaer and Raymakers, 1996).

SUMMARY

The Amur Sturgeon *Acipenser schrencki* is one of two species of Acipenseriformes endemic to the Amur River basin. The population of *A. schrencki* has been declining for more than a century. The two species, *Huso dauricus* Kaluga and *A. schrencki*, are rarely distinguished in harvest statistics making it difficult to assess the impact of harvest on individual species. Females of the most common morph (grey) of *A. schrencki* reach sexual maturity between 10-14 years, at which time their size ranges from 105-125 cm and 6-18.5 kg. The Amur River has not yet been affected by the construction of hydroelectric dams, however plans have been made in the past and their achievement would render numerous spawning grounds inaccessible. The water quality of the river has been increasingly degraded by pollution from oil products, mineral fertilisers from agricultural development, and by-products of gold mining operations. Although the effects of pollution on sturgeon have not been studied. The main threat appears to be over-exploitation.

Official catch statistics of *A. schrencki* in the Russian Federation and the former USSR dropped from 607 t in 1881 to 8 t in 1996. International trade in 1998 totalled 3,897 kg of caviar, with 2,366 kg being exported from China and 1,385 kg exported from the Russian Federation. The annual catch quota set for *A. schrencki* by the Russian Federation for 1999 and 2000 was 15 t. Export quotas of 1,500 kg (1999) and 2,000 kg (2000) were set for caviar, with a meat quota of 4,000 kg (2000). According to current available data it appears that the Russian Federation exceeded the 1999 caviar export quota by 1,476 kg. In China, the 1998 catch quota was 1,888 individuals with a caviar production reaching 3,235 kg. Export quotas of 2,783 kg (1998) and 2,510 kg (1999 and 2000) were set. The domestic consumption of caviar in China is estimated at 20% of the national production.

The species is recommended under Decision 11.106 for inclusion in category 1 (i).

DISTRIBUTION AND POPULATION

The CITES database lists the current distribution of *Acipenser schrencki* as: China, Japan?, Russian Federation (Anon., 2000a).

A. schrencki is classified as Endangered by IUCN (1996):
EN A2d China, Russia [Amur River].

A. schrencki is endemic to the Amur River system which is 4,092 km long if its longest tributary, the Shilka River, is included (Krykhtin and Svirskii, 1997a). The middle reach of the Amur River is bordered by both the Russian Federation and China. It is 975 km long and extends from the city of Blagoveshchensk (Heihe in China) to the mouth of the Ussuri (Wusuli) River between Khabarovsk (Russian Federation) and Fuyan (China), 966 km from the estuary (Krykhtin and Svirskii, 1997a). All populations have declined considerably since the turn of the century (Krykhtin and Svirskii, 1997b). The middle Amur population has been most affected being subject to catch from both the Russian and Chinese sides (Krykhtin and Svirskii, 1997b; Wei *et al.*, 1997).

China: The species occurs in the lower to upper reaches of Amur river basin, but is most common in the middle Amur and the lower reaches of the Songhuajiang and Wusulijiang river basins. The species is included in the China Red Data Book (Yue and Chen, 1998). Although the biomass of the wild populations is unknown, it is clear that stocks of sturgeon in the Amur River are declining dramatically due to over-fishing (Wei *et al.*, 1997).

Russian Federation: *A. schrencki* occurs from the Amur River delta to the upper reaches, including the Argun and Shilka tributaries. There are two morphs, the grey being more abundant than the brown (Krykhtin and Svirskii, 1997a) which occurs in the middle and lower parts of the Amur and has restricted movements to 100 km up or down stream. The grey morph is found in the middle to upper reaches of the Amur River. It

occurs in four fragmented populations: the estuary population which does not migrate to the sea (3,000 individuals); the middle Amur (190, 000 individuals); the lower Amur (95,000 individuals); and the Zeya-Bureya lowlands population, which is thought to be on the verge of extinction (Krykhtin and Svirskii, 1997a; 1997b).

HABITAT AND ECOLOGY

A. schrencki feeds on benthic freshwater molluscs and the larvae of the Arctic Lamprey *Lampetra japonica* (Svirskii, 1971, cited in Krykhtin and Svirskii, 1997a). Spawning occurs from the end of May to early-July (Krykhtin and Svirskii, 1997a). At 12 years of age, mature *A. schrencki* females of the brown morph are 96-117 cm long and weigh 3.5-5.6 kg (Krykhtin and Svirskii, 1997a), while females of the grey morph are 125-142 cm long and weigh 8.3-16.4 kg. Most grey morphs mature at the age of 10-14 years, with a length of 105-125 cm and weighing 6-18.5 kg (Krykhtin and Svirskii, 1997a). The maximum known size of the grey morph is approximately 3 m in length and more than 190 kg at an age exceeding 60 years (Krykhtin and Svirskii, 1997a). According to Wei *et al.* (1997), surveys carried out on the Chinese side of the Amur River suggest that the youngest spawning age for *A. schrencki* males is 7-8 years, at about 103 cm in length and 4 kg in weight. The corresponding age and size for females are 9-10 years, 105 cm and 6 kg. Fecundity ranges from 114,000-1,292,000 with a mean of 385,000 eggs. The average number of eggs per gram of body weight is 44.6 (Wei *et al.*, 1997). However, the survival rates of juveniles is low (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, August 2000).

THREATS TO SURVIVAL AND DOMESTIC USE

Unlike many of the large rivers in the region, the Amur has not been affected by construction of hydroelectric dams. However, it is feared that hydroelectric dams would restrict access to many spawning sites (Birstein, 1993). The river has been polluted by oil products, mineral fertilisers and by-products of gold mining operations. Agricultural pollution downstream from towns is increasing from both the Russian and Chinese sides (Krykhtin and Svirskii, 1997a; Matthieson, 1993). The effects of pollution on sturgeon have not been studied. The main threat appears to be overexploitation, particularly in the middle Amur (Heilong) River, where both Chinese and Russian fisherman are operating.

Current information suggests that amendments of previous regulations and/or adoption of new legislation are in effect in both range States. However, enforcement measures appear to be lacking and numerous experts as well as government officials have reported increasing pressure from illegal fishing practices and criminal activities such as sturgeon poaching and black markets which occur in a large part of the range (Medetsky, 2000; Winchester, 2000).

At the turn of the century the highest volumes of *A. schrencki* were caught in the middle Amur River where both brown and grey morphs of the species occur. Official catch records of *A. schrencki* in the former USSR and the Russian Federation decreased from 607 t in 1881 to 4.2 t in 1948 (Krykhtin and Svirskii, 1997a). In 1996, the catch was 8 t (Anon., 2000b). Official records from China indicate that sturgeon (both Kaluga and Amur Sturgeon) catches on the Chinese side of the Amur River peaked in 1987 (452 t), dropped to 136 t in 1997 and then increased to 149 t in 1998 and 141 t in 1999 (TRAFFIC East Asia *in litt.* to TRAFFIC Europe, September 2000).

Russian catch data for 1996 suggest that the current composition of *A. schrencki* versus *Huso dauricus* in the Amur River is 9% and 91% respectively (Anon., 2000b). However, late 19th century records suggested an equal composition (Krykhtin and Svirskii, 1997a).

In China, all caviar of wild sources is from *A. schrencki* and *Huso dauricus*. Heilongjiang Province is the only province in China where caviar is processed from wild specimens (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000). Most caviar originating from the Amur River in China is destined for export, with domestic consumption representing less than 20% of the total caviar production in the country (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

Domestic consumption of *A. schrencki* caviar in China (kilogrammes)

	1993	1994	1995	1996	1997	Average
<i>A. schrencki</i>	386.0	407.0	403.0	480.0	582.0	451.6

Source: CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000

Aquaculture products (captive breeding sources) of a variety of sturgeon species are sold on the domestic market (including in Hong Kong Special Administrative Region) and the rest is exported (TRAFFIC East Asia *in litt.* to TRAFFIC Europe, September 2000).

INTERNATIONAL TRADE

The listing of *A. schrencki* in CITES Appendix II entered into effect on 1 April 1998. Available complete trade data is therefore limited to nine months of trade (April-December 1998).

Gross exports and the comparative tabulation of trade in *A. schrencki* are given in the Appendix.

According to Annual Report data for 1998, the China and the Russian Federation reported respective exports of 2,366 kg and 1,385 kg of *A. schrencki* caviar. The importing countries were Japan (2,589 kg), Germany (900 kg), USA (247 kg) and South Africa (15 kg). In 1999, Russian exports of *A. schrencki* caviar totalled 2,976 kg. These shipments were imported into Japan and the USA (CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 September 2000).

Caviar export quotas established for *A. schrencki* (kilogrammes)

		1998	pre- Convention	1999	2000
Caviar	RU	1,700		1,500	2,000
	CN	2,783	1,000	2,510	2,510
	Total	4,481		4,010	4,510
Meat	RU	0		0	4,000

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Only trade in caviar is detailed in 1998 CITES data. However, the Russian Federation has established an export quota of 4,000 kg of meat for 2000.

Confiscation of *A. schrencki* shipments was not reported in 1998, however smuggling of caviar on the Sino-Russian border seems to occur frequently during the migration season (Medetsky, 2000).

CONSERVATION MEASURES

The species is not fully protected in the Russian Federation or China.

China: The listing of *A. schrencki* in the State Category of Protected Wildlife is expected to be adopted in the near future (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). In the early 1950s, the Heilongjiang Province authorities adopted protection and management regulations, such as gear restrictions, harvest size, closed seasons and areas, and the requirement of a fishing licence. These were renewed in 1982 through "The Heilongjiang Ordinance on the Protection and Propagation of Fisheries Resources". However, the regulations have not been fully implemented (Wei *et al.*, 1997).

Restriction on fishing size

The Ordinance of 1982 prescribed the minimum catch size limits for *A. schrencki* to be 100 cm or 4 kg (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). Specimens under these size and weight specifications must be released (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Other restrictions

According to the "Protocol on Fisheries Resources Conservation, Regulation and Multiplication in Border Areas of Heilong (Amur) River and Wusuli (Ussuri) River of The People's Republic of China and the Russian Federation" adopted on 27 May 1994, fishing on the Heilong River is prohibited from mid-June to mid-

July. The protocol also established areas where fisheries are prohibited: the mainstream of the Heilong River from Dagangzi, Luobei County, to Saniangkou where the Heilong and Songhua rivers converge. These areas were designated as no-fishing zones by the Government of China. The fishing grounds stretch over 1,270 km, from Fuyan (downstream) to Heihe (or Aihui) (Blagoveshchensk on the Russian side) (upstream). These measures protect *A. schrencki*'s natural breeding sites and hence the survival of juvenile sturgeon.

The catch of *A. schrencki* on the Chinese side of the Heilong (Amur) River requires prior acquisition of a fishing licence. In 1991, 2,248 sturgeon fishing licences were issued, and in 2000 the number had been reduced to 1,850 (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Restocking of juveniles

From 1988 to 1998, a total of 3,780,000 larvae and fingerlings of *A. schrencki* bred artificially in hatcheries in China were released in the Heilong (Amur) and Wusuli rivers (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). Since 1998, the restocking activities of the main hatchery, Qindeli State Farm (Heilongjiang Province), have been jeopardised by financial constraints and an increase in the price of eggs (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

In 2000, the Heilongjiang Province Fisheries Bureau authorised seven stations to undertake research on the captive breeding of *A. schrencki* and *H. dauricus* (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Determination and allocation of quotas by local government

Based on the average annual production for 1986-1988, the average size of fish landed, decreasing caviar production and the hydrological resource and migration conditions, Chinese scientists calculated Amur River sturgeon (*A. schrencki* and *H. dauricus*) annual catch quotas for the period 1989-1997. This quota was revised in 1997 and the Heilongjiang Province Fisheries Bureau adopted a new total catch quota of 105.4 t for 1998. The 1998 quota is composed of 1,888 individuals for *A. schrencki* with an expected caviar production of 3,235 kg and 309 individuals for *H. dauricus* with an expected caviar production of 4,045 kg (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Processing and export company management

Caviar processing companies are required to register with the Heilongjiang Provincial Fisheries Bureau. In 1998, six companies were registered as sturgeon product processors, all established in Heilongjiang Province. Six other companies were registered as caviar exporters: four in Heilongjiang Province, one in Beijing and one in Dalian Free Trade Zone (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000). According to the Ministry of Agriculture's (MOA) "*Regulation on Special License of Aquatic Wildlife Utilization, People's Republic of China*", caviar processing companies must acquire special qualification from the local Fisheries Bureau, including a "special (caviar) purchase license" and a "special (caviar) processing license". Caviar purchase and processing are controlled by the local Fisheries Bureau. The Fisheries Bureau will gradually decrease the number of licenses issued and also monitor those companies with special purchase and production licenses. According to the "*Law of Wildlife Protection of the People's Republic of China*", the MOA only issues caviar trade licenses to a limited number of import and export companies. The MOA also strictly manages those companies issued with this special license (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Russian Federation: A total prohibition on sturgeon catch throughout the USSR was adopted in 1923 and withdrawn in 1930. A ban on the catch of *A. schrencki* and *Huso dauricus* was introduced in 1958. This prohibition is still in effect (Krykhtin and Svirskii, 1997a). However, a tolerance called "controlled catch" for incidental and scientific catches is permitted. These catches are the current source of caviar and sturgeon meat from the Amur River. "Controlled catch" is not well defined and difficult to enforce. Russian catch quotas of 15 t annually for 1999 and 2000 were adopted by the Federal Government following recommendations set by the State Fisheries Committee in consensus with an independent body of experts (Anon., 2000b).

CAPTIVE BREEDING

China: Aquaculture has been successful for 30 years (Dr Zhang Chun-Guang, *in litt.* to IUCN/SSC, Wildlife Trade Programme, September 2000). An aquaculture station was set up on the Chinese side of the Amur River and was in operation from 1988 to 1991. During this period it released 900,000 fry and 168,000 fingerlings into the Amur River. The restocking programme was not able to continue after this time due to lack of funds (Dr Zhang Chun-Guang, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). However, additional hatcheries in China continued to restock the Amur and Wusuli rivers (see above) until 1998 (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

Russian Federation: Sturgeon hatcheries were under construction during 1996–97 (Krykhtin and Svirskii, 1997b). A broodstock also exists in captivity in research stations (Svirskii *et al.*, 1993).

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database and China have not reported the results of their research on sturgeon captive breeding nor the development of commercial farms.

However, *A. schrenckii* is recognised as an important species for China's sturgeon aquaculture industry (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000). In 1992, construction began of a large commercial sturgeon aquaculture farm in Dalian at the tip of the peninsula in Bo Hai Bay, west of Beijing. The expected annual production of the farm was 2,000 t, with 800 marketable fish (average weight of 2.5 kg) and 200 t of fish fillets in 1999 and 50 t of caviar in 2001. Other sturgeon products planned in the marketing strategy of the farm are skin (to be processed into leather), bones, meat, and internal organs (as raw materials for biomedical products and tonics). This large intensive aquaculture facility was developed with the assistance of Russian experts (Anon., 2000c).

REFERENCES

- Anon. (2000a). [http://www.cites.org/eng/dbase/fauna-](http://www.cites.org/eng/dbase/fauna)
- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia field investigations. Unpublished report, Pp. 23.
- Anon. (2000c). Dalian Russian Sturgeon Cultivation Co., Ltd. Promotion material (not dated).
- Birstein, V.J. (1993). Draft Application to CITES: Order Acipenseriformes. Unpublished.
- IUCN. (1996). *IUCN Red List of Threatened Animals*. IUCN, Gland, Switzerland
- Krykhtin, M.L. and Svirskii, V.G.. (1997a) Sturgeon of the Amur River, Kaluga *Huso dauricus* Georgy and Amur sturgeon *Acipenser schrenckii* Brandt. In Birstein, V.J., Waldman, J.R. and Bemis, W.E. (1997). Sturgeon biodiversity and conservation. Kluwer Academic Publishers, Dordrecht. *Environment Biology of Fishes*. 48:231-239.
- Krykhtin, M.L. and Svirskii, V.G. (1997b). Sturgeon catch and the current status of sturgeon stocks in the Amur River. In Birstein, V.J., Bauer, A. and Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 29-34.
- Matthieson, P. (1993). The last cranes of Siberia. *The New Yorker*, May 3 1993:76-86.
- Medetsky, A. (2000). Black caviar feeds thousands on Amur. *Vladivostok News*, 15 September 2000.
- Svirskii, V.G., Nazarov, V.A. and Rachek, E.I. (1993). The Amur river sturgeons and prospects for their culture in the far east of Russia. In *Abstract Bulletin*, Second International Symposium on Sturgeons. VNIRO Publ., Moscow. pp. 67.
- Wei, Qiwei, Ke, F., Zhang, J., Zhuang, P., Luo, J., Zhou, R. and Yang, W. (1997). Biology, fisheries and conservation of Sturgeons and paddlefish in China. In Birstein, V.J., Waldman, J.R. and Bemis, W.E. 1997. Sturgeon biodiversity and conservation. Kluwer Academic Publishers, Dordrecht. *Environment Biology of Fishes*. 48:241-255.
- Winchester, S. (2000). *National Geographic*, February 2000. Pp. 7-14.
- Yue, Pei-qi. and Chen, Y.Y. (1998). *China Red Data Book of Endangered Animals-Pisces*. Science Press, Beijing, Pp. 6-8.

APPENDIX

Gross exports of, and export quotas for *Acipenser schrencki* 1998

TAXON	TERM	UNIT	Exporter	1998	Export quotas 1998 (kg)
<i>Acipenser schrencki</i>	eggs		RU	145	
<i>Acipenser schrencki</i>	eggs	kg	CN	2,352	2,783 + 1,000 pre-Convention
<i>Acipenser schrencki</i>	eggs	kg	DE	15	
<i>Acipenser schrencki</i>	eggs	kg	RU	1,385	1,700

Comparative Tabulation of trade in *Acipenser schrencki* 1998

<i>Year</i>	Imp. Exp.		Imports reported				Exports reported						
			Origin	Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	DE	CN		900 kg	eggs	T	O	900 kg	eggs	T	O		
1998	JP	CN						1,349 kg	eggs	T	W		
1998	JP	RU						1,240 kg	eggs	T	W		
1998	US	CN		102 kg	eggs	T	W	102 kg	eggs	T	W		
1998	US	RU		145	eggs	T	W						
1998	US	RU						145 kg	eggs	T	W		
1998	ZA	DE	CN					15 kg	eggs	T	O		
	TOTAL			1,147 kg				3,751 kg					

Acipenser stellatus Pallas, 1771

Stellate Sturgeon
Esturgeon étoilé
Esturión estrelado

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

Stellate Sturgeon *Acipenser stellatus* has a wide distribution, occurring in the Caspian Sea, Black Sea and Sea of Azov and originally in many of the tributaries of these seas. However, the construction of dams on virtually all these rivers has significantly reduced the spawning area. The Ural River is one of the few major spawning rivers that has not been dammed. In addition to loss of habitat, the species has been subject to overfishing and egg-production has been disrupted due to environmental pollution. In the early 1990s, around 30% of the population in the Caspian Sea and almost all the stock in the Sea of Azov originated from re-stocking programmes. Re-stocking reportedly occurs in Azerbaijan, Iran, Kazakhstan, Romania, the Russian Federation and Yugoslavia but monitoring of the sturgeon restocking is performed only in Iran. In addition, hatcheries have had difficulty in obtaining broodstock.

A. stellatus has always been rare in the Danube River, with only 7.8 tonnes (t) average annual catch in the middle Danube River between 1958 and 1981. However, recorded landings in Romania and Bulgaria increased from 6.2 t in 1997 to 14.5 t in 1999. In contrast, annual catches of the species have generally decreased in other range States, except in the Danube River Basin. In the northern Caspian Sea, reported catches dropped from 4,650 t in 1977 to approximately 4,600 t in 1990 and to an estimated 1,060 t for 1997 for the whole of the Caspian (according to data collected during this review). Range States decreased the catch quotas for *A. stellatus* by around one third for Azerbaijan, Kazakhstan and Ukraine in 2000. The size of the domestic market for sturgeon products in the Russian Federation are unknown. Iran estimated its domestic consumption to be 5% of the country's caviar production and 65% of the sturgeon meat production.

The major product from wild of *A. stellatus* recorded in international trade in 1998 was caviar, with caviar of this species being commercially known as Sevruga caviar. According to CITES data, caviar from *A. stellatus* represents the largest portion of caviar from all Acipenseriformes traded internationally. *A. stellatus* exports from range States in 1998 included a total of 103 t of caviar, mainly from Iran (46 t with an export quota of 50 t) and the Russian Federation (39 t with an export quota of 85 t, including quotas for Azerbaijan and Kazakhstan). For 1999, the Russian Federation set the caviar export quota at 42 t and exported 32.4 t. Assuming a standard rate of 6-7% in weight of caviar of the total catch (males and females), the total volume of caviar exported in 1998 represents a total catch of 1,471-1,717 t of *A. stellatus* caught by range States. Iran also exported 8,794 kg of meat of wild origin (including 4,000 kg declared as "bodies"). The total combined caviar export quotas for 2000 decreased from that of 1999 (from 111.8 t to 102.9 t), but increased significantly for meat (from 42 t to 334.5 t). A total of 25,700 live fish (of captive bred source - 25,000 from Poland imported into Germany) were exported in 1998. However, commercial aquaculture of this species appears to be in its infancy.

The species is recommended under Decision 11.106 for inclusion in category 1 (i).

DISTRIBUTION AND POPULATION

The CITES database lists the current distribution of *Acipenser stellatus* as: Azerbaijan, Bulgaria, Czech Republic, Georgia, Greece?, Hungary, Iran (Islamic Republic of), Italy?, Kazakhstan, Moldova, Republic of, Romania, Russian Federation, Slovakia, Turkey, Turkmenistan, Ukraine, Yugoslavia

A. stellatus is classified as Endangered by IUCN (1996):

EN A2d Azerbaijan, Bulgaria, Georgia, Hungary, Iran, Kazakhstan, Moldova, Romania, Russian Federation, Turkmenistan, Ukraine, Yugoslavia

Caspian Sea Stock: VU A1acde+ 2d Azerbaijan, Iran, Kazakhstan, Russian Federation, Turkmenistan

Sea of Azov Stock: EN A2d Russia [Mediterranean and Black Sea]

Black Sea Stock: EN A1acde+2d Hungary, Romania, Ukraine, Yugoslavia [Mediterranean and Black Sea].

A. stellatus inhabits the Caspian, Azov and Black Seas although isolated catches have also been recorded in the Adriatic Sea (Birstein, 1993a; Shubina *et al.*, 1989). It is anadromous and enters the rivers that flow into these seas to spawn. The largest population is believed to occur in the Volga-Caspian region and produces 80% of the total sturgeon catch in this region (Anon., 1997). The species has always been considered rare in the middle Danube and after the construction of the Djerdap I and II hydropower stations, it is considered extirpated from both the upper and middle Danube (Hensel and Holèik, 1997). Although there are no recent population estimates available, Volovik *et al.* (1993) estimated the total *A. stellatus* biomass in the Sea of Azov during the mid-1980s to be 12,390 t. In 1990, 55,000 sturgeon were found dead on the shore of the Sea of Azov as a result of pollution and no population estimates have been made since. Due to reduced natural reproduction, about 60-98% of each generation in the Sea of Azov consisted of farm grown fish in the early 1990s (Volovik, *et al.*, 1993), while in the Caspian Sea around 30% originated from hatcheries. A decline in the commercial catch of this species reflects a decline in the population (Anon., 1997).

Azerbaijan: There is no recent information on the distribution in the Kura River, but Khodorevskaya *et al.* (1997) note that sturgeon no longer use the Kura and Terek rivers to spawn.

Bulgaria: As part of the Black Sea population, *A. stellatus* spawn in the Danube River. The breeding seasons are in spring, early-April (when the first catch of the year is recorded) and in autumn (end-September) (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). There has been a remarkable decline in *A. stellatus* catches in recent years (Karapetkova *et al.*, 1995), which now amount to only a few hundred kilogrammes per year (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Georgia: The south east coast of the Black Sea is an important feeding and wintering area for sturgeon (*A. stellatus*, *A. nudiventris*, *A. persicus colchicus*, *A. sturio* and *Huso huso*) that migrate upstream in a number of rivers for spawning. The main rivers used are the Supsa, Inguri, Chorokhi and particularly the Rioni (Zarkua and Tsuladze, 1999).

Greece: In the Aegean and Adriatic Seas, only a few isolated specimens have been recorded and it is doubtful if spawning still occurs in Greece (Lelek, 1987).

Hungary: Pinter (1991, cited in CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000) reported that last specimen was caught in 1965 (Mohács, Danube). Pr N. Bacalbasa-Dobrovici (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) considers the species as exceptionally rare whilst A. Rónyai (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) notes that *A. stellatus* is now considered extinct in Hungary.

Iran: *A. stellatus* previously entered the Sefidrud and the Gorgan rivers on the southern coast of the Caspian Sea to spawn (Shubina *et al.*, 1989). According to Dr. J. Holèik (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000), the species does not occur in these rivers. However, Laluyee (1996) and Ramin (1998, both cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September, 2000), maintain that *A. stellatus* enters the Sefidrud, Tajen and Gorganrud rivers in the southern shores of the Caspian Sea to spawn. *A. stellatus* migrates to the Sefidrud and other rivers during spring and autumn, the intensity of spawning being high particularly after heavy rainfall (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Moldova: The Moldovan range on the Danube River is 811 m and sturgeon also occur in the Pruth River, a tributary of the Danube River (junction at km 137th) (Anon., 2000c). There are no recent records of the species occurring in the Dniester River. Pr. Bacalbasa-Dobrovici (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) reports that specimens have been caught in the Pruth, while Dr V. Lobcenco (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) notes that there have been no records of the species in recent years.

Romania: *A. stellatus* enters the Danube from the Black Sea to spawn downstream of Calarash. However, catches have dramatically declined after the damming of the lower Danube by the Djerdap I and Djerdap II hydropower stations (Bacalbasa-Dobrovici, 1997).

Russian Federation: The largest populations of *A. stellatus* occur in the Caspian Sea from where they ascend to the Volga, Ural and Terek-Kuma rivers to spawn. From the Sea of Azov the spawning population ascends to the lower sections of the Don and Kuban rivers (Shubina *et al.*, 1989). Volovik *et al.* (1993) estimated that *A. stellatus* accounts for 12,390 t (21%) of the total sturgeon biomass of the Sea of Azov.

Approximately 60% of the historic spawning sites remain intact in the northern Caspian Sea (Khodorevskaya *et al.*, 1997). Between 1985 and 1992, stocks declined sharply in the Volga and Ural rivers. In 1992, the Ural River population was only one third of the numbers observed in 1986, and Birstein (1993b) reports 90,000 spawning individuals in the Ural. The spawning population of the Volga River has been estimated by Levin (1997) to be approximately 116,000 mature individuals (a decline in recent years from 500,000 individuals). There are no population estimates available for the other rivers in which they spawn, but Vlasenko (1990) and Levin (1995) believe populations to be minimal. In the Sea of Azov, some natural spawning grounds are no longer accessible due to the construction of dams on the Don and Kuban rivers (Anon., 2000b).

Autonomous Republic of Dagestan: No information is available on the current status of populations in the Sulak and Samur rivers.

Serbia: The annual catch of 1.4-2.0 t dropped to 0.184 t in 1971 when Djerdap I Dam was built on the Danube (Hensel and Holèik, 1997).

Slovakia: The last specimen of *A. stellatus* was caught in 1926 (Holèik, 1995).

Turkey: *A. stellatus* has been observed here, possibly in the Kizil-Irmak and Yesil-Irmac rivers, but it is unclear whether they spawn. The Coruh River is a known spawning site and unidentified sturgeon species have been reported in small numbers in the Sakarya and Mert rivers. In the late 1980s, local fishermen estimated that only five to six gravid female sturgeon were caught from the Yesilirmak to supply a hatchery and up to 20 from the Kizilirmak (Edwards and Doroshov, 1989).

Ukraine: There is very little recent information on the distribution and status of *A. stellatus* in the Ukraine. According to Pavlov (1980), *A. stellatus* was formerly considered rare in the Dnieper River, where it was known mainly from the lower river. It also occurred in the Dniester and Bug rivers, but as a result of the construction of the Dubosar and Oleksandriv hydropower stations, its distribution was reduced and catches subsequently declined. The species occurs in the Don and Kuban rivers, where it has been a commercially important fish, however the catch declined significantly after the construction of the Cimlyansk hydropower station.

Former Yugoslavia: There are no records for the species in Croatian rivers (Taler, 1953). However, there was a significant drop in catch of all sturgeon species after 1970 and 1984, when the lower stretch of the Danube was respectively dammed by Djerdap I and II hydropower stations (Jankoviè, 1993; 1996; Simonovic and Nikolic, 1996). Today, the species occurs in a 17.8 km stretch of the Danube River, from the Djerdap II dam to the border, close to the Timok River mouth (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

HABITAT AND ECOLOGY

The species reaches a total length of 2.18 m, a maximum weight of 54 kg and maximum age of 35 years (Shubina *et al.*, 1989). The normal size ranges between 1.0-1.2 m and 6-8 kg. The age of *A. stellatus* spawners ranges between 6-28 years, on average 11-16 years. The average weight of females is 11-12 kg and 6-7 kg for males. Females constitute 40-48% of the spawning population (Levin, 1997).

Unlike other sturgeon species, *A. stellatus* is found in the middle and upper water layers. There are two forms, one of which spawns in winter and the other in spring. The spring form migration peaks in summer

and ends in late autumn. The winter migrating form hibernates in rivers and reproduces the following year. The main feeding grounds of the Caspian are to the north, but a winter migration to the central and southern areas of the Caspian has been observed (Barannikova *et al.*, 1995).

Based on catches and caviar production in the late 1970s, the rate of caviar extracted per fish caught was estimated at 7 kg per 100 kg of the total catch (including males and females) for the three main commercial species of the northern Caspian Sea, *Huso huso*, *A. gueldenstaedtii* and *A. stellatus* (Doroshov and Binkowski, 1985, cited in Williot and Bourguignon, 1991).

Iran: Two forms of the species are found in the Caspian Sea: *A. stellatus* Pallas or north Caspian, and *A. stellatus cyrensis* Berg or south Caspian. The two forms can not be visually differentiated. The south Caspian form can be identified on the basis of later sexual maturity, slower growth and a lower fecundity (Bilialova *et al.*, 1989, cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). Molecular studies on mitochondrial DNA and alloenzymes have reported high genetic diversity in *A. stellatus* in the southern Caspian Sea region (Pourkazemi, 1995). Sexual maturity of *A. s. cyrensis* is attained at an average age of 12 years for females (range of 6-18 years) and 9 years for males (range of 6-15 years) (Int. Sturgeon Res. Inst., 1998, cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Until 1993, the mouth of the Ural River was dredged annually to allow vessels to travel upstream to Atyrau (former Gur'yev) harbour. Since 1993, this annual management of the river channel has not been carried out and is an obstacle to migration (Anon., 2000c).

THREATS TO SURVIVAL AND DOMESTIC USE

Hydroelectric power stations and water reservoirs constructed on the vast majority of rivers have reduced spawning ground availability by approximately 40% in the Caspian basin (Barannikova *et al.*, 1995). In addition, pollution has far exceeded maximum permissible concentrations in the Volga River and the northern Caspian Sea (Romanov and Altuf'yev, 1993). However, since 1991 the intensity of industrial development decreased, leading to an improvement in water-quality. The rise of the sea level of the Caspian now threatens to engulf lakes of pollutants along the shore line.

Overexploitation has posed an increasing threat over the years. In the mid 1990s, open sea fishing in the Caspian and reputedly high levels of illegal catch served to reduce the number of mature sturgeons to such a level that the Volgograd hatcheries in the Russian Federation were unable to collect mature fish to stock the hatcheries.

In the former Soviet Union and Iran, strict legislation regarding sturgeon fisheries had been in effect for decades until the emergence of three new independent States in the Caspian Sea Basin in 1992 (Azerbaijan, Kazakhstan and Turkmenistan) (De Meulenaer and Raymakers, 1996). Current information suggests that amendments of previous regulations and/or adoption of new legislation are now in effect in most new range States in the Caspian Sea as well as in Black Sea (including the Danube River) and Sea of Azov basins. In the latter, Bulgaria, Romania and Ukraine are the most important sturgeon fishing nations. However, enforcement measures appear to be lacking and numerous experts as well as government officials have reported an increasing pressure of illegal fishing practices and criminal activities surrounding the caviar trade in most of the range (Anon., 2000b; Dobbs, 1992; Doward, 2000; Evtouchenko, 1997; King, 1998; Ward, 2000a; 2000b)

The total commercial sturgeon catch in the Caspian Sea Basin peaked at 27,300 t in 1977. *A. stellatus* represented 17% (4,650 t) of this total (Levin, 1997). The catch of *A. stellatus* in the northern Caspian Sea decreased from approximately 4,600 t in 1990 to 1,600 t in 1994 (Khodorevskaya *et al.*, 1997). The Russian Caspian Sea catch of *A. stellatus* was 1,060 t in 1997 (Anon., 2000b). The species was always considered rare in the Danube River, with only an average annual catch of 7.8 t in the middle Danube River between 1958 and 1981 (Hensel and Holèik, 1997). However, landings recorded from the lower Danube River in Romania and Bulgaria increased from 6.2 t in 1997 to 14.5 t in 1999.

Azerbaijan: The range of the species in the Kura River, including spawning grounds, has been reduced to about 160 ha due to the construction of the Vavarin reservoir and dam. Waters of this region represent important winter feeding grounds, however, lakes of deposited oil and pollutants lie on the shore lines of the Caspian and as the sea level rises these 'lakes' will be engulfed (Dumont, 1995).

Along the northern and southern coast of Azerbaijan, within the 200-nautical mile Exclusive Economic Zone, sturgeon species are caught as by-catch in stationary nets and *kilka* (Clupeidae) fishing devices. Young and adult migrating fish are caught by coastal fisheries when the water temperatures change. Young fish make up to 17.7% of the by-catch.

Species composition of recorded sturgeon landings

	<i>A. persicus & A. gueldenstaedtii</i>	<i>A. stellatus</i>	<i>H. huso</i>	<i>A. nudiventris</i>
Catch	63.47%	30.77%		5.76%
By-catch	78.50%	16.80%	3.70%	1%

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Annual recorded landings (catch and by-catch) of *A. stellatus* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. stellatus</i>	20.00	27.00	25.00	92.00	67.60	31.80	44.80	27.10	20.70	33.50	20.76
Total	74.00	88.20	111.00	240.10	201.20	162.50	120.30	128.60	82.40	108.90	63.22

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Bulgaria: Wild sturgeon populations have been threatened by a decrease in the number of spawning grounds on the Danube River as a result of human development, particularly hydroelectric facilities. The long-lasting unregulated fisheries have disrupted the age composition of the populations through the selective catch of sexually mature fish, and to by-catch of young sturgeon as they migrate to the sea. Water pollution has also influenced the metabolism of sturgeon (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). The species is affected by overfishing and pollution (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Annual catch of *A. stellatus* (tonnes)

	1995	1996	1997	1998	1999
<i>A. stellatus</i>	0.1	0.5	1.4	3.7	6.0
Total sturgeon catch	14.7	32.5	49.8	55.9	50.5
n° of licenses issued	0	0	122	155	169

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Domestic consumption of caviar in Bulgaria is limited to 7-10% of the total caviar production of Bulgaria. The entire production of sturgeon meat is consumed within Bulgaria.

Georgia: During the fishing seasons, particularly coastal anchovy fisheries, numerous sturgeon juveniles are caught as by-catch. The estimated catch of all species of sturgeon declined from 77,000 individuals in 1977, to 57,000 in 1980 and 27,000 in 1986. During the 1990s, illegal fishing in Georgia expanded rapidly due to poor control (Zarkua and Tsuladze, 1999).

Iran: For the past five years, the Iranian domestic market for caviar and sturgeon meat has been estimated at 5% and 65% of the country's total production respectively.

Annual catch of *A. stellatus* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>A. stellatus</i>	1,114.0	1,019.5	917.9	679.1	672.5	562.2	507.1	360.1	346.3	288.1	-
Total	2,296.3	2,315.1	2,058.0	1,462.1	1,380.9	1,216.3	1,310.0	1,043.5	1,127.9	923.3	182.4

* Catch in spring 2000 only.

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

Production of sturgeon meat and caviar in Iran in 1992 and 1999 (tonnes)

	1992		1999	
	Meat % *	Caviar % *	Meat % *	Caviar % *
A. stellatus	695.4	43.9	153.4	56.6
Total	1,604.2	262.3	730.2	99.3

* Percentages relate to the share of *A. stellatus* products in the total Iranian sturgeon fisheries.

Source: Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000

The above tables indicate that the total tonnage of meat and caviar of *A. stellatus* produced in Iran comprised 92% and 88% of the country's annual catch in weight in 1992 and 1999, respectively.

Kazakhstan: The Ural River provides 1,400 ha of spawning ground and is the only river with unregulated flow into the Caspian. Oil fields are developing rapidly in the northern Caspian, especially the Tengiz oil field (Sagers, 1994); pollution from this industry affects spawning and feeding grounds. Radioactive contamination from a nuclear reactor also poses a threat (Dumont, 1995).

Annual catch of *A. stellatus* in the 1990s (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
A. stellatus	1,498.0	1,313.0	1,002.0	547.0	274.0	197.9	175.1	227.6	235.0	140.0	117.4
Total	1,963.0	1,767.0	1,437.3	1,109.0	556.0	573.0	367.8	389.5	490.0	279.0	230.5

* Spring catch.

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Moldova: Catch of *A. stellatus* has been prohibited since the species was listed in the Red Data Book in 1995 (Anon., 2000c).

Romania: Pollution emitted between 1970 and 1989 has been the greatest threat to this species. The loss of spawning habitat due to dam construction has posed an important threat, but to a lesser extent than pollution. Overfishing and poaching have increased in recent years (Pr N. Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Annual catch of *A. stellatus* for 1991 to 1999 (kilogrammes)

	1991	1992	1993	1994	1995	1996	1997	1998	1999
	2,184	2,910	2,405	2,852	6,506	4,941	4,788	4,003	8,556

Source: Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, 5 September 2000

Note: These data include catches in the Danube River and Delta as well as in the Black Sea.

From 1991 to 1999, approximately 32% of the total Romanian sturgeon catch in the Danube River comprised *A. stellatus*. Romanian scientific surveys have concluded that the population of *A. stellatus* is quite well represented in the Danube River (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, 5 September 2000).

Russian Federation: Construction of the Volgograd dam led to a 82% decrease in the range and spawning grounds formerly available in the Volga River. The remaining area can accommodate 280,000 spawning sturgeon, but during the last 30 years their numbers have been estimated to be considerably less than this (Veshecv, 1993; 1995). Dam construction has also restricted the range in the Terek and Sulak rivers, with the spawning ground having been reduced to 132 ha and 202 ha respectively (Vlasenko, 1990). In the Black Sea and Sea of Azov region, the Kuban River flow has been regulated resulting in the loss of 140,000 ha of spawning grounds. The Tsymlyansk reservoir, on the Don River in 1952, removed approximately 68,000 ha of spawning ground (Volovik *et al.*, 1993).

The Volga River hatcheries started to release juveniles in 1957. From the release of half a million juveniles in 1959, numbers increased to around 16 million in 1970. In the late 1980's, approximately 30% of *A. stellatus* caught in the Volga River were from hatcheries. The numbers released did not seem to affect the numbers of wild spawners later entering the rivers (Khodorevskaya *et al.*, 1997).

Shagaeva *et al.* (1993) found that 100% of the eggs taken from *A. stellatus* in the lower Volga (both of wild and captive bred (hatchery) sources) in 1989 had anomalous larvae because of poor environmental conditions. It is thought that these eggs may not have been viable. However, water quality has started to improve in the Volga–Caspian region since the decline of heavy industrial activities in the former USSR.

From 1992 to 1997, approximately 40% of the Russian sturgeon catches in the Caspian Sea were composed of *A. stellatus* (Anon., 2000b)

Catch of *A. stellatus* in the Caspian Sea (tonnes)

	1992	1993	1994	1995	1996	1997
Catch	3,610	1,349	1,467	968	675	449

Source: Caspian Fisheries Research Institute (KaspNIRKh), cited in Anon., 2000b

The level of domestic consumption of sturgeon products in the Russian Federation is unknown, but is said to be the main outlet of illegal sturgeon fisheries (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). Based on the number of sets of illegal fishing equipment recorded by enforcement agencies of the Caspian Sea and the Sea of Azov, the illegal sturgeon catch was evaluated at six to ten times the legal catch (Anon., 2000b). In 1997, the estimated volume of the Moscow sturgeon meat market was three times the annual catch quotas for sturgeon species (Anon., 1998).

Turkey: Spawning grounds have been reduced. From 1969 to 1979, the Turkish annual sturgeon landings ranged from 310 t to 10 t, with an annual average of 112 t (Edwards and Doroshov, 1989). In 1989 in Samsun, a Turkish landing site on the Black Sea, one caviar processor claimed that his production had decreased from 8 t per year of caviar between the 1940s and the 1970s, to 200 kg per year at the end of the 1980s. In 1989, sturgeon were caught as by-catch by trawlers that operated legally along the Turkish coast at a distance greater than three nautical miles, but also illegally within the three nautical mile limit. This represented the main fishing pressure on sturgeon species. In April 1989, *A. stellatus*, *A. gueldenstaedtii*, *A. sturio* and *H. huso* were observed in Samsun (Edwards and Doroshov, 1989). The investigators were told that *A. nudiiventris* was also landed in Samsun.

Ukraine: Pollution and, to a lesser extent, loss of spawning grounds, have affected this species. Overfishing has been a problem in recent years.

Former Yugoslavia: Most of the Danube River spawning grounds are no longer accessible due to the construction of the Djerdap dams.

INTERNATIONAL TRADE

The listing of *A. stellatus* in CITES Appendix II entered into effect on 1 April 1998. Complete data for this species were therefore limited to nine months of trade (April-December 1998). Gross exports of trade in specimens of *A. stellatus* in 1998 are given in the Appendix.

CITES Annual Reports for 1998 suggest that caviar of *A. stellatus*, commercially known as “Sevruga”, represented the largest portion of all sturgeon caviar traded internationally. Range States exported a total of 103,085 kg in 1998; Iran (45,955 kg), the Russian Federation (38,981 kg), Kazakhstan (17,467 kg), Azerbaijan (422 kg) and Romania (260 kg).

Other specimens reported in international trade in 1998 were meat (8,794 kg exported by Iran and imported by France and Italy), live fish (25,000 individuals exported by Poland and imported into Germany (CITES Management Authority of Poland, *in litt.* to TRAFFIC Europe, 31 August 2000)), and eight skins (probably destined to be processed into leather, exported by Iran to Switzerland and subsequently re-exported to Italy). The live juveniles exported by Poland in 1998 were of captive bred source (see “Captive Breeding”).

In addition to caviar, a further 51,500 specimens (fertilised eggs or caviar ; see *Note* at the end of this section) were traded internationally. Of these, 26,500 were declared to be from a wild source and

originated from Hong Kong (non-range State), and 20,400 were declared of pre-Convention stock imported from the Russian Federation into Greece and subsequently re-exported to the USA.

Exports of *A. stellatus* from Russia in 1999

	Quantity (kg)	Description	Importer	No of permits
A. stellatus	2.50	Live eggs	GR DE	2 Export
	32,418.72	Caviar	US, CY, DK, DE, GB, FR, CH, NL, BE, JP	52 Export + 10 Re-export
	5,740.00	Meat	US	2 Export

Source: CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 September 2000

Export quotas established by range States (kilogrammes)

	1998		1999		2000	
	Caviar	Meat	Caviar	Meat	Caviar	Meat
Azerbaijan			3,800	38,000	2,450	24,500
Iran	50,000		40,000		35,000	
Kazakhstan			24,000		22,000	199,000
Romania			2,000	4,000	2,100	23,000
Russian Fed.	85,000		42,000		Caspian Azov	30,350 11,000
Total	135,000		111,800	42,000		102,900 334,500

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Illegal trade: In 1998, a total of 541 kg *A. stellatus* caviar shipments of unknown origin was confiscated by US Customs upon import. This comprised 410 kg re-exported by Turkey and 114 kg re-exported by the United Arab Emirates. Discrepancies between trade statistics of countries importing and exporting caviar can indicate illegal activities. Such discrepancies were found in 1995 and 1996 for instance for Turkey (customs declared caviar exports 21 t and 80 t respectively) and European Union Member States (6 t and 20 t respectively) (Raymakers, 1999). Additionally, Turkish 1996 caviar exports totalled 121 t (Raymakers, 1999), while the country's sturgeon fisheries has been limited to only few incidental catches since the late 1980s (Edwards and Doroshov, 1989).

Poaching and smuggling are closely related and have been intensively reported in the media of both range States and importing countries (Evtouchenko, 1997; McDonald, 2000; Snyder, 2000). Much of the illegally caught sturgeon is destined to the domestic market, particularly meat (Anon., 1998).

Note: The use of the term "eggs" and the unit "kg" or "g" to describe shipments of both caviar and fertilised eggs complicates the analysis of trade data. Similar difficulties arise from the unit used for export quotas of fertilised eggs, which is kilogrammes ("kg") instead of "number of specimens". Kilogrammes do not directly correspond to the number of specimens, and hence does not allow for the effective evaluation of quotas.

CONSERVATION MEASURES

A. stellatus is not fully protected in any range State other than Moldova. However, a license is required for fishing in most of the countries and private sturgeon fisheries are prohibited in Iran.

Regarding juvenile restocking efforts, despite the large-scale aquaculture programmes of the Russian Federation since the 1960s (20 hatcheries were fully operational in the 1980s) and Iran (hatcheries have been operational for 20 years), stocks have continued to decline (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Azerbaijan: The Intergovernmental Commission for Caspian Biological Resources, composed of representatives of all range States, has set annual catch quotas.

Annual catch quotas allocated to Azerbaijan (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
A. stellatus	120	163	163								42.7
Total	200	308	510	480	360	182	160.5	160.5	160.5	108.67	108.35

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Three hatcheries were built in the lower Kura River by the authorities of the former Soviet Union in the 1960s. A World Bank feasibility study reported that the total capacity of the hatcheries was originally 12 million, but that this capacity had dropped significantly over the years (T. Turner, World Bank consultant, *in litt.* to TRAFFIC Europe, 6 October 2000). World Bank funds have been allocated for the construction of a new sturgeon hatchery and farm that will have the capacity of 15 million fry per year; this should be operational in the next few years. Concerns have been raised about the availability of adult sturgeon to constitute the broodstock of the farm.

Estimated number of sturgeon fry released in the Kura River (million fingerlings)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
A. stellatus	3.747	2.870	0.400	0.608	0.392	0.302	1.040	2.100	1.320	8.822	6.869
Total	17.523	9.082	2.980	1.838	1.142	1.242	4.070	6.065	6.220	20.290	16.965

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

The State institution responsible for sturgeon fishery and processing matters is "State Concern Azerbalyg". Commercial sturgeon fisheries are prohibited in the open sea. No fishing license was required until 2000. The only fishing gear permitted for sturgeon catch in the Kura River is "sweep nets" of 35-55 mm mesh size. The minimum size limit set for *A. stellatus* is 90 cm in the Kura River, and 110 cm in coastal waters (Anon., 2000c).

Bulgaria: Measures at the national level as described hereunder are needed, but they must be linked to co-ordinated actions of all countries in the Black Sea region. In Bulgaria, sturgeon fisheries are prohibited for 30 days, from 20 April to 20 May each year, at the peak of the breeding season. The authorities may approve a 15-day extension of the closed season pending the natural conditions of the year. Explosives, poison, bottom trawling and dragging, as well as stunning substances and devices are prohibited fishing methods. The minimum size limit for *A. stellatus* is 75 cm in the Danube River and 100 cm along the Black Sea coast.

Export quotas are set on the basis of catch data, trends and the biological characteristics of the species. A system was established for the distribution of export quotas among export companies that are obliged to restock the Danube River with sturgeon fry before they are authorised to export sturgeon specimens, but until now, no fry of *A. stellatus* have been released. Due to budgetary constraints, Bulgaria has not monitored the wild sturgeon population in recent years, nor the success of its restocking programme (e.g. through tagging of released fingerlings). However, plans have been prepared for these activities (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). A special prohibition was adopted on the use of "unbaited hooks" for sturgeon catch (Anon., 2000c).

Georgia: Until the early 1990s, artificial breeding of sturgeon was carried out in the Varistkhi Sturgeon Plant built in 1983 on the Rioni River with mixed results (Zarkua and Tsuladze, 1999).

Hungary: The species is protected according to *Decree No. 1/1982 (III.15.) OKTH on the Protected and Strictly Protected Species of Flora and Fauna, Value of their Specimens, Determination of the Range of Protected and Strictly Protected Caves and Exemptions from Restrictions and Prohibitions Set for Certain Protected Animal Species*, last amended by the *15/1996 (VII.26.) Decree of the Minister for Environment*. The *Nature Conservation Act No. 53 of 1996, Article 43* orders that for all protected species, prior authorisation from the National Park Directorate is required for the capture, killing, possession, captive breeding, introduction of specimens from exotic population, artificial exchange of genetic matter between populations, exchange and sale or purchase of *A. stellatus* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: At all sturgeon catch stations, biometric measurements are carried out for each species. The age, length, sex and weight of fish landed are recorded. The population structure is estimated on the basis of age groups. As a result of these studies carried out for the past 10 years, the number of catch stations were decreased by 50%, changes were adopted on sturgeon catch methods, and lower export quotas were set for the past five years. Export quotas are established on the basis of results of research programmes and the annual monitoring of stocks. These activities include the following parameters and activities: Maximum Sustainable Yield and total catch, rehabilitation of stocks, management and sustainable exploitation through restriction of illegal catch and overfishing. Monitoring fishing efforts and regulating the time of catch are also carried out on the basis of research results (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Annual release of *A. stellatus* fry (million fish)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<i>A. stellatus</i>	0.36	0.47	0.07	0.30	0.46	0.27	0.22	0.29	0.18	0.13
Total	4.56	6.60	3.45	4.17	5.91	9.13	12.35	21.63	24.56	19.10

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

A labelling system is in place, where each tin of caviar holds a sign or mark that indicates at least the date and place of catch, the name of the catch station, name of processor and the species.

Additional conservation measures have been undertaken for all sturgeon in the southern shores of the Caspian Sea. The fishing permits of more than 6,000 fishermen engaged in fishing using gill nets have been bought, and employment for these fishermen has been created in other fishery operations regarding catch of kilka and beach seine catch. Coastal patrols have also been strengthened (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Commercial sturgeon catch is only allowed in the tributaries of the Caspian Sea, but scientific catch is permitted in the open sea with an annual quota of 20 t. Sturgeon fisheries are regulated by law. There are two fishing seasons, one during spring spawning and the other during the time of winter migration. The exact dates of the fishing seasons vary with the climate and are therefore set annually by authorised scientific organisations. As in most former Soviet Republics, "sweep nets" (bottom nets that are hauled several times a day) with a minimum mesh size and which are marked with the name and address of the owner, are the only legal fishing gear for sturgeon. Each fisherman can only set one net at the time. A fishing license is also required. Sturgeon fishing grounds (called "tonia") are limited: there are 12 on the Ural River and 8 on the Kigach River. The minimum catch size limit for *A. stellatus* is 110 cm. The ratio of males and females caught is not regulated. The purchase price at the landing site is based on a fixed legal rate of caviar weight extracted per female; *A. stellatus* is set at 15.4% of the total body weight (Anon., 2000c). Annual commercial catch quotas are allocated to Kazakhstan by the Intergovernmental Commission for Caspian Biological Resources.

Annual commercial catch quotas allocated to Kazakhstan (tonnes)

	1996	1997	1998	1999	2000
<i>A. stellatus</i>			305	215	215.5
Total	375	480	460	405	371

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Two hatcheries, the Uralo-Atyrau and the Atyyrau, have been operational since 1998.

Annual release of *A. stellatus* fry (number of fingerlings)

	1998	1999	2000
<i>A. stellatus</i>	1,058,993	2,581,005	3,771,060
Total	1,706,193	5,318,471	5,507,160

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Moldova: Fishery of *A. stellatus* is prohibited in Moldova since the species was included in the National Red Data Book (Anon., 2000c) and is protected by law (Anon., 1999). No monitoring of the population of *A. stellatus* has been undertaken recently (Dr V. Lobcenco, *in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, September 2000).

Romania: Bacalbasa-Dobrovici (*in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, September 2000) believes that the fishing pressure should be reduced in the lower Danube. He reports that the existing legislation pertaining to the region is neither adequate nor enforced, particularly in Romania. A fishing license is required in Romania to capture this species (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, 5 September 2000). There is a research programme for artificial spawning and culture of *A. stellatus*, and a restocking programme started in 1993. The total number of juveniles restocked in the Danube River is 44,850 fingerlings (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, 5 September 2000).

The type of sturgeon fishing gear is strictly regulated by Romanian law. The fishing season is set every year by Ministerial Order. A fishing license is required to catch sturgeon in Romanian waters. The minimum size limit for *A. stellatus* is 100 cm. The catch quota for *A. stellatus* for 1998 and 1999 was set at 20 t per year. Catch quotas are calculated by independent scientific institutes, such as the Danube Delta Institute in Tulcea and the Centre for Conservation, Fish Farming, Fishing and Fish Processing in Galati (Anon., 2000c).

Release of sturgeon juveniles in the Danube River (number of fingerlings)

	1993	1994	1995	1996	1997	1998	1999
<i>A. stellatus</i>	1,000	4,000	10,000	11,600	10,000	6,750	

Source: Centre for Conservation, Fish Farming, Fishing and Fish Processing, Galati, December 2000, cited in Anon., 2000c

Russian Federation: For Caspian Sea populations, the former Soviet Republics set annual catch quotas in agreement with Iran. Sturgeon catch quotas have been established by the Intergovernmental Commission for Caspian Biological Resources since the early 1990s. The annual share of sturgeon catches of each former USSR Caspian range State as adopted by the Commission since 1993 are as follows: the Russian Federation 70%, Kazakhstan 17.6%, Azerbaijan 6.1%, and Turkmenistan 6.3%. There are two fishing seasons in the Caspian Sea waters: from ice melting to June 5 and from 1 September until the time of ice formation. A license is required to catch sturgeon in all Russian waters. The minimum size limit for *A. stellatus* caught in Russian Caspian waters is 110 cm along the coast and 90 cm in the northern Caspian Sea (Anon., 2000c). Russian catch quotas are currently declared by the Federal Government following the recommendations given by the State Fisheries Committee in consensus with an independent council of experts (Anon., 2000b).

Release of *A. stellatus* in the Volga River and the Sea of Azov by Russian hatcheries (number of fingerlings)

	1979-1980	1981-1985	1986-1990	1991-1995	1996-1998
Volga River	16,400,000	19,400,000	16,200,000	10,100,000	15,500,000
Sea of Azov	18,950,000	19,120,000	73,800,000	13,040,000	14,293,103

Source: Azov and Caspian Fisheries Research Institute (Az- and KaspNIRKh), cited in Anon., 2000b

From 1979 to 1998, approximately 24.6% and 50.4% of all fry released by Russian hatcheries in the Volga River and the Sea of Azov respectively comprised *A. stellatus* (Anon., 2000b).

Export quotas established by the CITES Management Authority of the Russian Federation consist of 90% of the caviar production, which in turn varies according to the catch quotas set for each species of sturgeon (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). The annual catch quotas or Total Allowable Catches (TAC) allocated by the State Fisheries Committee, is based on the recommendations of scientific agencies such as the Russian Federal Research Institute of Fisheries and Oceanography (VNIRO) and the Pacific Research Institute of Fisheries and Oceanography (TINRO). The TAC includes the commercial catch, the scientific catch, and the catch of mature males and females destined to form hatchery broodstock.

Recent data on the monitoring of catch per species, including the size composition and sex ratio of fish caught are not available. The volume of domestic consumption of sturgeon products in the Russian Federation, mostly meat and caviar, is unknown (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000).

Turkey: In 1979, with the exception of *H. huso* of more than 140 cm long, the fishery for sturgeon was prohibited in Turkish waters (Edwards and Doroshov, 1989). Circular 34/1, effective from 2000-2002, prohibits the collection of Acipenseriformes (CITES Management Authority of Turkey, *in litt.* to TRAFFIC Europe, 20 September 2000).

Turkmenistan: In coastal commercial waters, sturgeon fishery is allowed for scientific research only, with an annual catch limit of 20 t. The Turkmenistan State Committee of Fisheries issues an annual license for sturgeon commercial catches to the State enterprise, Balkanbalyk. No other company, Turkmen or otherwise has the right to engage in commercial sturgeon fisheries in Turkmen inland waters. An annual catch quota for the species is set by the Intergovernmental Commission for Caspian Biological Resources. There is no sex ratio set for sturgeon landings (Anon., 2000c).

Ukraine: Sturgeon fishing in the Danube River is permitted throughout the year, except from 16 March to 15 April. The by-catch of sturgeon is regulated. For example, for sprat trawling, a tolerance of two sturgeon specimens of trade length (see below) may be caught per tonne of sprat. The mesh size of nets used to catch sturgeon should be a minimum of 120 mm and a maximum of 150mm. Only fishermen who have been allocated an annual sturgeon quota by the State Committee for Fishery are allowed to catch sturgeon. The minimum size limits for *A. stellatus* are 100 cm for fish caught in the Black Sea and the Danube River, and 80 cm for those caught in the Sea of Azov (Anon., 2000c).

Annual catch quotas for *A. stellatus* for 1997 to 1999 (tonnes)

		1996	1997	1998	1999
Total		505	424	210	225
A. stellatus	Sea of Azov and adjacent waters		125	45	30
	Black Sea and Danube River			10	3

Source: State Committee for Fishery of Ukraine, cited in Anon., 2000c

There is currently no sturgeon fry restocking programme in Ukraine.

Yugoslavia: The closed fishing season set in the Danube River for *H. huso*, *A. gueldenstaedtii* and *A. stellatus*, is from 15 May to 15 June. Certain types of fishing gear are prohibited to catch sturgeon and an agreement was signed between Romania and Yugoslavia on the prohibition of gear that cause high mortality of untargeted fish. Any fishing activity, including all commercial and sport fisheries, requires a license in Yugoslav waters. Licenses are issued on a per district basis and the sturgeon fishing grounds are limited to 17.8 km downstream of the Djerdap II dam and the mouth of the Timok River. The minimum size limit for *A. stellatus* was set at 75 cm, but catches of the species have been prohibited from 1996 to 2000. A restocking programme is being carried out by the Centre for Fish Reproduction of the "Djerdap Fishing Company" in Malla Vrbica village (Anon., 2000c).

CAPTIVE BREEDING

The reduction in area of natural spawning grounds due to river flow regulation has caused population declines. Fish reared in hatcheries have therefore been introduced to maintain the stock. Approximately 30% of the Caspian stock originates from hatcheries (Barannikova, 1995). In 1993, 60-98% of each generation of *A. stellatus* in the Sea of Azov was derived from aquaculture (Volovik *et al.*, 1993). *A. stellatus* has also been introduced into aquaculture in Belgium and France (Welcomme, 1988).

Overfishing has prevented hatcheries from catching enough mature fish to maintain the necessary broodstock (Artyukhin, 1997). Levin (1995) stated that although aquaculture contributes considerably to the maintenance of the stocks, it cannot compensate for the damage caused to natural reproduction.

Azerbaijan: Three hatcheries were built in the 1960s and have contributed to the former Soviet Union sturgeon restocking programme ever since. Their combined annual productive capacity was 12 million fry, but this has dropped significantly due to rising levels of the Caspian Sea, and the lack of funds allocated to hatchery maintenance (T. Turner, World Bank consultant, *in litt.* to TRAFFIC Europe, 6 October 2000).

Georgia: Until the early 1990s, artificial breeding of sturgeon species was carried out at the Varistkhi Sturgeon Plant. Built in 1983 and located on the Rioni River, the results obtained were not always 'satisfactory' (Zarkua and Tsuladze, 1999).

Hungary: There are no captive breeding facilities for *A. stellatus* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Captive breeding activities have been carried out for *A. stellatus* in the southern shores of the Caspian Sea. The breeding, rearing and release of more than 10 million fingerlings into the sea has been undertaken in recent years. The artificial rearing and production of broodstock consisting of 2,000 breeders in earthen ponds has also been undertaken and *A. stellatus* has also been introduced into permanent inland water bodies (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Two hatcheries exist in Kazakhstan, Uralo-Atyrau and Atyrau, which have been operational since 1998 (CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, September 2000).

Poland: The presence of fish farms and hatcheries of sturgeon in Poland was confirmed by the CITES Management Authority of Poland (*in litt.* to TRAFFIC Europe, 31 August 2000). The following sturgeon species are bred in captivity: *A. baerii*, *A. gueldenstaedtii*, *A. ruthenus*, *A. stellatus*, *H. huso*, *Polyodon spathula* and hybrids of some of these species. In most cases, hybrids are F1 generation live specimens (fertilised eggs) imported in to Poland from range States, particularly the Russian Federation. Some farms keep F2 generation specimens, which are bred in Poland and used as broodstock, partly for the production of juveniles for both the domestic and international market.

Romania: An aquaculture programme exists and has been releasing 10,000 fish of 1-3 g per year (Bacalbasa-Dobrovici, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). The programme is thought to be efficient, but quantitative data are lacking (Suciu *et al.*, 1998). Approximately 30,000 juveniles are retained each year for aquaculture and research purposes (Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme Trade Programme, 5 September 2000). In Romania, hatcheries and captive breeding of *A. stellatus* in earth ponds, with the aim of developing commercial farms, are still at an experimental stage (Dr N. Patriche, pers. comm. to TRAFFIC Europe, September 1999).

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database, but according to aquaculture companies, there is currently no significant production of meat of *A. stellatus* in commercial farms.

REFERENCES

- Anon. (1996). *A Guide to Interpreting Outputs from the WCMC CITES Trade Database*, Produced by the World Conservation Monitoring Centre, Cambridge UK. Pp. 16
- Anon. (1997). Document Doc. 10.89; Prop. 10.65. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Anon. (1998). *Sturgeon catch and trade in the Russian part of the Caspian Sea*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, pp. 21.
- Anon. (1999). The Law of the Republic of Moldova on Fund of Natural Areas Protected by State (1998). In: "*Ekologicheskoe Zakonodatel'stvo Respubliki Moldova (1996-1998)*". Chisinau, BIOTICA Publ., 1999. P. 85-209. ISBN 9975-923-80-1.
- Anon. (2000a). <http://www.cites.org/eng/dbase/fauna->

- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, pp. 23.
- Anon. (2000c). *Sturgeon fisheries management and trade control measures in the Caspian Sea and Black Sea/ Sea of Azov range States*. TRAFFIC Europe field investigations, December 1999-January 2000. Unpublished.
- Artyukin, E.N. (1997). The current status of commercial sturgeon species in the Volga River-Caspian Sea Basin. In: Birstein, V.J., Bauer, A. Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 9-18.
- Bacalbasa-Dobrovici, N. (1997). Endangered migratory sturgeons of the lower Danube River and its delta. *Environmental Biology of Fishes* 48: 201-207.
- Barannikova, I.A., Burtsev, I.A., Vlasenko, A.D. Gershanovich, A.D., Markarov, E.V. and Chebanov, M.S. (1995). Sturgeon Fisheries in Russia. *Proceedings of the Second International Symposium on Sturgeons, September 6-11, 1993*. Moscow-Kostroma-Moscow. VNIRO Publication. Pp.124-130.
- Birstein, V.J. (1993a). Sturgeon and Paddlefishes: Threatened fishes in need of conservation. *Con. Biol.* (4): 773-787.
- Birstein, V.J. (1993b). Draft Application to CITES: Order Acipenseriformes. Unpublished.
- De Meulenaer, T. and Raymakers, C. (1996). *Sturgeons of the Caspian Sea and the international trade in caviar*. TRAFFIC International, iv + 71 Pp.
- Dobbs, M. (1992). The Coming Caviar Crisis: Overfishing, Pollution Endanger Sturgeon. *The Washington Post*, 30 May 1992.
- Doward, J. (2000). Roe rage as piracy rules. *The Guardian*, 21 May 2000.
- Dumont, H. (1995). Ecocide in the Caspian Sea. *Nature* 377:673-674.
- Edwards, D. and Doroshov, S. (1989). *Appraisal of the Sturgeon and Seatrout Fisheries and Proposals for a Rehabilitation Programme*. FAO Field Document FI.TCP/TUR/8853. Pp. 38.
- Evtouchenko, A. (1997). [Au Dagestan une explosion fait 54 morts. Accusés: les gangs du caviar. *Courrier International*, N° 323 (9-15 January 1997)](in French)
- Hensel, K. and Holèik, J. (1997). Past and current status of sturgeon in the upper and middle Danube. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 185-200.
- Holèik, J., (1995): 2. druh *Acipenser* (*Gladostomus*) *stellatus* Pallas, 1771. In: V. Baruš and O. Oliva (eds.) *Mihulovci Petromyzontes a ryby Osteichthyes*. 1. Fauna ČR a SR 28/1. Academia, Praha. Pp. 381-384.
- IUCN. (1996). *1996 Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- Jankoviè, D. (1993). Populations of Acipenseridae prior and after the construction of the HEPS Djerdap I and II. *Acta Biologica Iugoslavica, Seria E. Ichthyologia* 25:29-34
- Jankoviè, D. (1996). Ichthyofauna of the Danube in the Djerdap area after the construction of the Iron Gate I Hydroelectric Power Station. *Acta Universitatis Carolinae, Biologica* 40 (1996): 123-131
- Karapetkova, M., Zhivkov, M. and Pchelarov, T. (1995): *Ribite v B'Igariya*. Geya Libris, Sofia. 247 pp.
- Khodorevskaya, R.P., Dovgopol, G.F., Zhuravleva, O.L and Vlasenko, A.D. (1997). Present status of commercial stocks of sturgeons in the Caspian Sea basin. In: Birstein, V.J., Bauer, A. Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 209-219.
- King, R. (1998). [Por un puñado de caviar. *El Mundo*, 13 December 1998.] (in Spanish)
- Levin, A.V. (1995). Russian Sturgeon *Acipenser gueldenstaedti* Brandt, stocking in the Volga-Caspian Basin. *Proceedings of the Second International Symposium on Sturgeons, September 6-11, 1993*. Moscow-Kostroma-Moscow. VNIRO publication Pp.178-188.
- Levin, A.V. (1997). The Distribution and Migration of Sturgeon in the Caspian Sea. In: Birstein, V.J., Bauer, A. Kaiser-Pohlmann, A. (eds.) *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 13-19.
- McDonald, L. (2000). Fish that laid the golden egg faces extinction. *Daily Express* (UK), 26 July 2000.
- Pavlov, P.J. (1980). *Ribi*. Fauna Ukrainy. 8/1. Naukova Dumka, Kiiiv. 336 pp.
- Pintér, K. (1991). Sturgeon in Hungary, past and present situation. In: P. Williot (Ed.) *Acipenser*. Actes du premier colloque international sur l'esturgeon. Bordeaux, 3-6 October 1989. pp. 173-178.
- Pourkazemi, M.D., Skibinski, O.F. and Beardmore, J.A. (1995). Population structure of Stellate Sturgeon *Acipenser stellatus* Pallas, in southern parts of the Caspian sea studied using mtDNA and allozyme

- analysis. Abstracts, International Conf. Sturgeon Biodiversity and Conservation, NY. *The Sturgeon Quarterly*, 3 (4):10.
- Raymakers, C. (1999). Trade in Sturgeons from the Caspian Sea. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 149-161.
- Romanaov, A. A. and Altuf'ev, Yu.V. (1993). Ectopic Histogenesis of sexual cells of Caspian Se Sturgeon. *J. Ichthyol.* 33 (2): 140-150.
- Sagers, M.J. (1994). The oil industry in the southern-tier former soviet republics. *Post-Soviet Geography*, 35(5):267-298.
- Shagaeva, V.G., Nikol'skaya, M.P., Akimova, K.P., Markov, N.G. and Nikol'skaya, N.G. (1993). A study of the early ontogeny of Volga Sturgeon, (Acipenseridae) subjected to human activity. *J. Ichthyol.* 33(6):23-41.
- Shubina, T.N., Popova, A.A. and Vasil'ev, V.V. (1989). *Acipenser stellatus* Pallas, 1771. In: Holèik J. (ed). *The Freshwater Fishes of Europe. Vol. I/II: General Introduction of Fishes. Acipenseriformes.* Wiesbaden, AULA-Verlag. Pp. 395-443. In: Document Doc. 10.89; Prop. 10.65. 1997. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Simonovic, P.D., and Nikolic, V.P. (1996). Freshwater fish of Serbia: an annotated check list with some faunistic and zoogeographical considerations. *Bios (Macedonia, Greece)* 4: 137-156.
- Snyder, D. (2000). Md. Caviar Company Admits Smuggling. *The Washington Post*, 22 July 2000.
- Suciu, R. Ene, F. and Bacalbasa-Dobrovici, N. (1998). New data on the distribution of sturgeon in the Lower Danube River. Eds. Raute, M., Bacalbasa Dobrovici, N. and Oprea, L. In: *Aquarium 1998*, Galati, Romania. University "Dunarea de Jos".
- Taler, Z. (1953). Rasprostranjenje i popis slatkovodnih riba Jugoslavije. (The distribution and description of the freshwater fishes of Yugoslavia). *Glasnik Prirodnjaèog Muzeja Srpske Zemlje*, ser.B 5/6:425-455.
- Veschev, P.V. (1993). Effect of water level of the Volga River on reproduction of *A. stellatus*. *Vodnye Resoursy*, No.2: 225-228.
- Veschev, P.V. (1995). Natural reproduction of the Volga River Stellate Sturgeon, *Acipenser stellatus*, under new Fishing regulations, *J. Ichthyol.* 35 (9): 281-294.
- Volovik, S.P., Dubinina, V.A. and Semenova, A.D. (1993). Hydrobiology and dynamics of fisheries in the Asov Sea. Studies and reviews. *General Fisheries Council for the Mediterranean. No. 64. FAO, Rome. Pp1-58.*
- Vlasenko, A.D. (1990). Sturgeon Population Size in the Caspian Sea. *Rybnoe Khozyaistvo*, 7:53-56.
- Ward, P. (2000a). Azov Sturgeons Can't be Touched. *Crimean Time*, 18 August 2000.
- Ward, P. (2000b). Makhachkala: the Scale of Poaching in the Caspian has Reached Enormous Proportions. *Regions.Ru/Russia Regions*, 15 August 2000.
- Welcomme R.L. (1988). International introductions of inland aquatic species. *FAO Fisheries Technical Paper*. 294: 318 pp.
- Williot, P. and Bourguignon, G. (1991). Sturgeon and caviar production, current status and perspectives. In: Williot, P. (ed.), *Acipenser*, Cemagref Publications, Pp. 509-513.
- Zarkua, Z. and Tsuladze, V. (1999). Initiative needed to boost sturgeon numbers. *Eurofish* 3/99:40-41.

Gross exports* and export quotas for *Acipenser stellatus* 1998

TAXON	TERM	UNIT	Exporter	1998	Export quotas 1998
<i>Acipenser stellatus</i>	bodies #1	Kg	IR	4,000	
<i>Acipenser stellatus</i>	Eggs		AE	237	
<i>Acipenser stellatus</i>	Eggs		AU	4	
<i>Acipenser stellatus</i>	Eggs		CH	12	
<i>Acipenser stellatus</i>	Eggs		DE	26,538	
<i>Acipenser stellatus</i>	Eggs		DK	128	
<i>Acipenser stellatus</i>	eggs #2		GR	20,400	
<i>Acipenser stellatus</i>	Eggs		NL	713	
<i>Acipenser stellatus</i>	Eggs		SG	1	
<i>Acipenser stellatus</i>	Eggs		US	297	
<i>Acipenser stellatus</i>	Eggs	Kg	AE	12,168	
<i>Acipenser stellatus</i>	Eggs	Kg	AU	214	
<i>Acipenser stellatus</i>	Eggs	Kg	AZ	422	
<i>Acipenser stellatus</i>	Eggs	Kg	BE	221	
<i>Acipenser stellatus</i>	Eggs	Kg	CA	21	
<i>Acipenser stellatus</i>	Eggs	Kg	CH	19,059	
<i>Acipenser stellatus</i>	Eggs	Kg	DE	15,638	
<i>Acipenser stellatus</i>	Eggs	Kg	DK	4,694	
<i>Acipenser stellatus</i>	eggs	Kg	FI	50	
<i>Acipenser stellatus</i>	eggs	Kg	FR	1,398	
<i>Acipenser stellatus</i>	eggs	Kg	GB	1,015	
<i>Acipenser stellatus</i>	eggs	Kg	GR	68	
<i>Acipenser stellatus</i>	eggs	Kg	HK	41	
<i>Acipenser stellatus</i>	eggs	Kg	IR	45,955	50,000
<i>Acipenser stellatus</i>	eggs	Kg	KZ	17,467	
<i>Acipenser stellatus</i>	eggs	Kg	NL	6	
<i>Acipenser stellatus</i>	eggs	Kg	PE	17	
<i>Acipenser stellatus</i>	eggs	Kg	PL	4,820	
<i>Acipenser stellatus</i>	eggs	Kg	RO	260	
<i>Acipenser stellatus</i>	eggs	Kg	RU	38,981	85,000 (incl. quotas for AZ and KZ)
<i>Acipenser stellatus</i>	eggs	Kg	SA	79	
<i>Acipenser stellatus</i>	eggs	Kg	SG	121	
<i>Acipenser stellatus</i>	eggs	Kg	TR	10,994	
<i>Acipenser stellatus</i>	eggs	Kg	US	9,656	
<i>Acipenser stellatus</i>	live		AE	706	
<i>Acipenser stellatus</i>	live		PL	25,000	
<i>Acipenser stellatus</i>	meat	Kg	DE	50	
<i>Acipenser stellatus</i>	meat	Kg	IR	4,735	
<i>Acipenser stellatus</i>	meat	Kg	US	9	
<i>Acipenser stellatus</i>	skins		CH	8	
<i>Acipenser stellatus</i>	skins		IR	8	

* **Definition:** "Gross exports are the sum of all reported exports and re-exports in a particular commodity or species in a particular year or series of years;... Gross trade is thus a simple measure of the total number of items recorded in international trade. However, gross trade may be an overestimate of the total number of actual specimens in trade as re-exports are not deducted from the total." (Anon., 1996). This is particularly applicable to caviar of which shipments are often exported, re-packaged and re-exported. For instance, in the mid-1990s, Germany re-exported as much as 45% of its annual caviar imports (De Meulenaer and Raymakers, 1996).

#1 The term "bodies" should be interpreted as "meat" (Dr P. Bronzi, *in litt.* to IUCN/SSC Wildlife Trade Programme, 8 September 2000).

#2 The import of 20,400 eggs into Greece (re-exported to the US) without a specified unit, almost certainly refers to "fertilised eggs" and not to caviar. The term "eggs" being used for both caviar and fertilised eggs and the absence of a unit (e.g. specimens, grammes, etc.) renders data analyses problematic. For example, in this case, if the quantity is interpreted in kilogrammes (20,400 kg) it would lead to an overestimate of 20.4 t of caviar in trade.

Comparative Tabulation of trade in *Acipenser stellatus* 1998

Year	Imp.	Exp.	Origin	Imports reported			P	S	Exports reported			P	S
				Quantity	Unit	Term			Quantity	Unit	Term		
1998	AU	AE	RU						21 kg	eggs		T	O
1998	CA	AE	IR						24 kg	eggs		T	W
1998	CA	AE	IR						4 kg	eggs		T	O
1998	CH	AE	IR	331 kg	eggs		T	W	99 kg	eggs		T	W
1998	CH	AE	IR	75 kg	eggs		T	O	75 kg	eggs		T	O
1998	CH	AE	RU	507 kg	eggs		T	O	39 kg	eggs		T	O
1998	CH	AE	RU						256 kg	eggs		T	W
1998	DK	AE	IR	34 kg	eggs		T	W	68 kg	eggs		T	W
1998	DK	AE	RU	2 kg	eggs		T	O					
1998	FR	AE	IR	252 kg	eggs		T	W					
1998	FR	AE	RU	741 kg	eggs		T	O	500 kg	eggs		T	O
1998	GB	AE	RU						10 kg	eggs		T	W
1998	HK	AE	IR						209 kg	eggs		T	W
1998	HK	AE	RU						419 kg	eggs		T	W
1998	JP	AE	IR						8 kg	eggs		T	W
1998	JP	AE	RU						139 kg	eggs		T	O
1998	KR	AE	RU						32 kg	eggs		T	O
1998	NO	AE	IR						35 kg	eggs		T	W
1998	NO	AE	IR	22 kg	eggs		T	O	21 kg	eggs		T	O
1998	PR	AE	DE						43 kg	eggs		T	O
1998	SG	AE	RU	4,845 kg	eggs		T	O	807 kg	eggs		T	O
1998	SG	AE	RU						606	live		T	O
1998	TH	AE	RU						200 kg	eggs		T	O
1998	TH	AE	RU						100	live		T	O
1998	TR	AE	RU						22 kg	eggs		T	O
1998	TW	AE	IR						40 kg	eggs		T	W
1998	US	AE		114 kg	eggs		T	I					
1998	US	AE	RU	21.895 kg	eggs		T	I					
1998	US	AE	RU	899 kg	eggs		T	W	191 kg	eggs		T	W
1998	US	AE	RU	772 kg	eggs		T	O	3,931 kg	eggs		T	O
1998	US	AE	RU	237	eggs		T	W					
1998	US	AE	RU	112 kg	eggs		T	I					
1998	US	AE	XX	93 kg	eggs		T	O					
1998	CN	AU	IR						15 kg	eggs		T	O
1998	NZ	AU							3,966 g	eggs		T	W
1998	NZ	AU	IR						22 kg	eggs		T	O
1998	NZ	AU	IR						4	eggs		T	O
1998	SG	AU	IR						98 kg	eggs		T	O
1998	TH	AU	IR						75 kg	eggs		T	O
1998	CA	BE	RU						67 kg	eggs		T	O
1998	TR	BE	RU						360 g	eggs		T	O
1998	US	BE	RU						153 kg	eggs		T	O
1998	US	CA	RU	21 kg	eggs		T	O					
1998	AT	CH	AZ	2 kg	eggs		T	O					
1998	AT	CH	IR						7 kg	eggs			O
1998	AT	CH	IR	5 kg	eggs		T	O					

Year	Imp.	Exp.	Origin	imports reported			exports reported			P	S
				Quantity	Unit	Term	Quantity	Unit	Term		
1998	AT	CH	RU					2 kg	eggs		O
1998	BE	CH	IR					5 kg	eggs		O
1998	BE	CH	IR	5 kg	eggs		T W				
1998	BE	CH	KZ	3,229 kg	eggs		T W				
1998	BE	CH	RU					3,228 kg	eggs		W
1998	CH	CH	IR					54 kg	eggs		O
1998	CH	CH	RU					18 kg	eggs		O
1998	CY	CH	IR					2 kg	eggs		O
1998	CY	CH	RU					1 kg	eggs		O
1998	DE	CH	IR	1,243 kg	eggs		T W				
1998	DE	CH	IR					934 kg	eggs		W
1998	DE	CH	IR					3 kg	eggs		O
1998	DE	CH	KZ	9,844 kg	eggs		T W				
1998	DE	CH	RU					9,855 kg	eggs		W
1998	DE	CH	RU					467 kg	eggs		O
1998	DK	CH	AZ	150 kg	eggs		T O				
1998	DK	CH	IR	289 kg	eggs		T W				
1998	DK	CH	IR					288 kg	eggs		W
1998	DK	CH	IR	203 kg	eggs		T O				
1998	DK	CH	IR					203 kg	eggs		O
1998	DK	CH	RU					150 kg	eggs		O
1998	FR	CH	IR					743 kg	eggs		W
1998	FR	CH	IR					495 kg	eggs		O
1998	FR	CH	IR	493 kg	eggs		T O				
1998	FR	CH	IR	331 kg	eggs		T W				
1998	FR	CH	KZ	154 kg	eggs		T W				
1998	FR	CH	RU					153 kg	eggs		W
1998	GA	CH	IR					36 kg	eggs		O
1998	GA	CH	IR					5 kg	eggs		W
1998	GA	CH	RU					32 kg	eggs		O
1998	GB	CH	AZ	145 kg	eggs		T W				
1998	GB	CH	AZ	97 kg	eggs		T O				
1998	GB	CH	KZ	12	eggs		T W				
1998	GB	CH	RU					679 kg	eggs		O
1998	GR	CH	IR					5 kg	eggs		O
1998	HK	CH	IR					78 kg	eggs		O
1998	HK	CH	IR					4 kg	eggs		W
1998	IT	CH	RU					230 kg	eggs		O
1998	IT	CH	IR					8	skins		W
1998	JP	CH	IR					88 kg	eggs		O
1998	JP	CH	RU					944 kg	eggs		O
1998	KW	CH	IR					22 kg	eggs		O
1998	MU	CH	RU					228 kg	eggs		O
1998	MX	CH	IR					28 kg	eggs		O
1998	PA	CH	IR					6 kg	eggs		O
1998	SA	CH	IR					1 kg	eggs		W
1998	SG	CH	IR	37 kg	eggs		T O				
1998	SG	CH	IR					37 kg	eggs		O
1998	SG	CH	IR					1 kg	eggs		W
1998	SG	CH	IR	1 kg	eggs		T W				
1998	SG	CH	RU					1 kg	eggs		O
1998	SG	CH	RU	1 kg	eggs		T O				
1998	TN	CH	RU					20 kg	eggs		O
1998	US	CH		500 g	eggs		T W				

Year	Imp.	Exp.	Origin	imports reported			exports reported			P	S					
				Quantity	Unit	Term	Quantity	Unit	Term							
1998	US	CH	AZ	400	g	eggs				T	O					
1998	US	CH	AZ	100	g	eggs				T	I					
1998	US	CH	AZ	2	kg	eggs				T	O					
1998	US	CH	IR					0	kg	eggs	O					
1998	US	CH	RU					4	kg	eggs	O					
1998	US	CH	XX	500	g	eggs				T	O					
1998	AD	DE	KZ					7	kg	eggs	T	W				
1998	AD	DE	RU					42	kg	eggs	T	W				
1998	AE	DE	XX					5	kg	eggs	T	O				
1998	AR	DE	KZ					2	kg	eggs	T	O				
1998	AR	DE	RU					20	kg	eggs	T	O				
1998	AU	DE	RU					140	kg	eggs	T	O				
1998	BR	DE	KZ					90	kg	eggs	T	O				
1998	BR	DE	RU					120	kg	eggs	T	O				
1998	CA	DE	KZ					19	kg	eggs	T	W				
1998	CA	DE	KZ					11	kg	eggs	T	O				
1998	CA	DE	RU					106	kg	eggs	T	O				
1998	CA	DE	RU					1	kg	eggs	T	W				
1998	CH	DE	IR	550	kg	eggs				T	W	508	kg	eggs	T	W
1998	CH	DE	IR	482	kg	eggs				T	O	147	kg	eggs	T	O
1998	CH	DE	KZ						97	kg	eggs	T	W			
1998	CH	DE	KZ						30	kg	eggs	T	O			
1998	CH	DE	RU	464	kg	eggs				T	O	65	kg	eggs	T	O
1998	CH	DE	RU	192	kg	eggs				T	W	89	kg	eggs	T	W
1998	CL	DE	IR						87	kg	eggs	T	W			
1998	CL	DE	RU						30	kg	eggs	T	O			
1998	CO	DE	XX						8	kg	eggs	T	O			
1998	CY	DE	KZ						22	kg	eggs	T	W			
1998	CZ	DE	IR						780	g	eggs	T	O			
1998	CZ	DE	IR	1	kg	eggs				T	W					
1998	CZ	DE	RU						672	g	eggs	T	O			
1998	CZ	DE	RU	6	kg	eggs				T	W	1	kg	eggs	T	W
1998	CZ	DE	RU						5	kg	eggs	T	O			
1998	HK	DE	IR						8	kg	eggs	T	O			
1998	HK	DE	KZ						25	kg	eggs	T	O			
1998	HK	DE	RU						157	kg	eggs	T	O			
1998	HR	DE	KZ						6	kg	eggs	T	W			
1998	HR	DE	RU						4	kg	eggs	T	O			
1998	IN	DE	KZ						200	g	eggs	T	O			
1998	IN	DE	XX						10	kg	eggs	T	O			
1998	JO	DE	KZ						5	kg	eggs	T	O			
1998	JP	DE	AZ						25	kg	eggs	T	O			
1998	JP	DE	KZ						268	kg	eggs	T	O			
1998	JP	DE	RU						2,990	kg	eggs	T	O			
1998	JP	DE	RU						727	kg	eggs	T	W			
1998	KR	DE	KZ						25	kg	eggs	T	O			
1998	KR	DE	KZ						20	kg	eggs	T	W			
1998	KR	DE	RU						204	g	eggs	T	O			
1998	KR	DE	RU						68	kg	eggs	T	O			
1998	KR	DE	RU	4	kg	eggs				T	W					
1998	KW	DE	RU						131	kg	eggs	T	O			
1998	MT	DE	RU						336	g	eggs	T	O			
1998	MU	DE	RU						7	kg	eggs	T	O			
1998	MX	DE	XX						10	kg	eggs	T	O			

Year	Imp.	Exp.	Origin	imports reporteda			exports reporteda						
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	MY	DE	AZ						150	kg	eggs	T	O
1998	MY	DE	IR						85	g	eggs	T	O
1998	MY	DE	RU						20	kg	eggs	T	O
1998	NO	DE	IR						21	kg	eggs	T	O
1998	NO	DE	KZ						50	kg	eggs	T	W
1998	NO	DE	KZ						24	kg	eggs	T	O
1998	NO	DE	RU						27	kg	eggs	T	O
1998	NO	DE	RU						9	kg	eggs	T	W
1998	NZ	DE	RU						60	kg	eggs	T	O
1998	PE	DE	KZ						2	kg	eggs	T	O
1998	PH	DE	KZ						100	kg	eggs	T	O
1998	PL	DE	KZ						5	kg	eggs	T	O
1998	PL	DE	RU						56	kg	eggs	T	O
1998	RU	DE	RU						8	kg	eggs	T	O
1998	SA	DE	KZ						115	kg	eggs	T	O
1998	SA	DE	RU						735	kg	eggs	T	O
1998	SG	DE	AZ	25	kg	eggs	T	O	13	kg	eggs	T	O
1998	SG	DE	IR	8	kg	eggs	T	O	8	kg	eggs	T	O
1998	SG	DE	KZ	168	kg	eggs	T	O	169	kg	eggs	T	O
1998	SG	DE	KZ						87	kg	eggs	T	W
1998	SG	DE	RU	1,095	kg	eggs	T	O	66	kg	eggs	T	O
1998	SG	DE	RU	87	kg	eggs	T	W					
1998	SY	DE	KZ						7	kg	eggs	T	O
1998	TH	DE	IR						256	g	eggs	T	O
1998	TH	DE	KZ						363	kg	eggs	T	O
1998	TH	DE	KZ						256	g	eggs	T	O
1998	TR	DE	KZ						18	kg	eggs	T	O
1998	TW	DE	RU						40	kg	eggs	T	O
1998	US	DE	HK	26,008		eggs	T	W					
1998	US	DE	KZ	174	kg	eggs	T	O	708	kg	eggs	T	O
1998	US	DE	KZ	114	kg	eggs	T	W	420	kg	eggs	T	W
1998	US	DE	NZ	300		eggs	T	O					
1998	US	DE	RU	3,716	kg	eggs	T	W					
1998	US	DE	RU	1,053	kg	eggs	T	O	2711	kg	eggs	T	O
1998	US	DE	RU	170		eggs	T	W					
1998	US	DE	RU	100	kg	eggs	P	W					
1998	US	DE	RU	60		eggs	T	O					
1998	US	DE	RU	50	kg	eggs	T	U					
1998	US	DE	XX	246	kg	eggs	T	O	59	kg	eggs	T	O
1998	US	DE	XX	17	kg	eggs	T	I					
1998	US	DE	XX	8	kg	eggs	T	W					
1998	US	DE	RU	50	kg	meat	T	W					
1998	UY	DE	KZ						14	kg	eggs	T	O
1998	ZA	DE	RU						682	g	eggs	T	W
1998	ZA	DE	RU						41	kg	eggs	T	O
1998	ZA	DE	RU						1	kg	eggs	T	W
1998	ZA	DE	XX						17	kg	eggs	T	O
1998	BR	DK	RU						65	kg	eggs	T	W
1998	CH	DK	IR	2,658	kg	eggs	T	W	1,586	kg	eggs	T	W
1998	CH	DK	IR	1,425	kg	eggs	T	O	1,930	kg	eggs	T	O
1998	CH	DK	RU	302	kg	eggs	T	W	262	kg	eggs	T	W
1998	CH	DK	RU	150	kg	eggs	T	O	270	kg	eggs	T	O
1998	JP	DK	IR						2	kg	eggs	T	W
1998	MA	DK	RU						21	kg	eggs	T	O

Year	Imp.	Exp.	Origin	imports reported			exports reported						
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P	S
1998	MU	DK	IR						27	kg	eggs	T	W
1998	NO	DK	IR						3	kg	eggs	T	W
1998	SA	DK	IR						3	kg	eggs	T	W
1998	TH	DK	IR						15	kg	eggs	T	O
1998	US	DK	RU	23	kg	eggs	T	W					
1998	US	DK	RU						2	kg	eggs	T	O
1998	US	DK	XX	128		eggs	T	W					
1998	RU	FI	RU						50	kg	eggs	T	W
1998	AD	FR	IR						16	kg	eggs	T	W
1998	AD	FR	IR						2	kg	eggs	T	O
1998	AN	FR	IR						2	kg	eggs	T	W
1998	AO	FR	IR						1	kg	eggs	T	W
1998	AR	FR	IR						3	kg	eggs	T	O
1998	AR	FR	KZ						4	kg	eggs	T	O
1998	BR	FR	IR						10	kg	eggs	T	O
1998	BR	FR	IR						4	kg	eggs	T	W
1998	BS	FR	IR						85	kg	eggs	T	W
1998	BS	FR	IR						2	kg	eggs	T	O
1998	CA	FR	AZ						9	kg	eggs	T	O
1998	CA	FR	IR						1	kg	eggs	T	O
1998	CA	FR	KZ						10	kg	eggs	T	O
1998	CA	FR	KZ						7	kg	eggs	T	W
1998	CA	FR	RU						10	kg	eggs	T	O
1998	CG	FR	IR						3	kg	eggs	T	W
1998	CH	FR	IR	155	kg	eggs	T	O	155	kg	eggs	T	O
1998	CH	FR	IR	143	kg	eggs	T	W	153	kg	eggs	T	W
1998	CH	FR	KZ	2	kg	eggs	T	W	2	kg	eggs	T	W
1998	CI	FR	IR						252	kg	eggs	T	W
1998	CI	FR	IR						8	kg	eggs	T	O
1998	CM	FR	IR						4	kg	eggs	T	W
1998	CM	FR	KZ						2	kg	eggs	T	O
1998	CY	FR	KZ						2	kg	eggs	T	W
1998	GA	FR	IR						7	kg	eggs	T	W
1998	GA	FR	KZ						1	kg	eggs	T	O
1998	HK	FR	IR						13	kg	eggs	T	W
1998	HK	FR	IR						11	kg	eggs	T	O
1998	HK	FR	KZ						19	kg	eggs	T	O
1998	HK	FR	KZ						5	kg	eggs	T	W
1998	HT	FR	IR						1	kg	eggs	T	W
1998	ID	FR	IR						2	kg	eggs	T	W
1998	JP	FR	IR						49	kg	eggs	T	W
1998	JP	FR	IR						1	kg	eggs	T	O
1998	JP	FR	KZ						4	kg	eggs	T	O
1998	KR	FR	IR						100	kg	eggs	T	O
1998	LB	FR	IR						22	kg	eggs	T	W
1998	LB	FR	IR						4	kg	eggs	T	O
1998	LC	FR	IR						2	kg	eggs	T	W
1998	LI	FR	KZ						1	kg	eggs	T	W
1998	MC	FR	IR						7	kg	eggs	T	W
1998	MR	FR	IR						1	kg	eggs	T	W
1998	MU	FR	IR						3	kg	eggs	T	W
1998	MX	FR	IR						36	kg	eggs	T	W
1998	MX	FR	IR						11	kg	eggs	T	O
1998	MX	FR	KZ						5	kg	eggs	T	W

Year	Imp.	Exp.	Origin	imports reported			exports reported					
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P
1998	MY	FR	IR						1 kg	eggs	T	W
1998	NC	FR	IR						12 kg	eggs	T	W
1998	PF	FR	IR	5 kg	eggs		T	W	3 kg	eggs	T	W
1998	PF	FR	KZ						1 kg	eggs	T	O
1998	PF	FR	RU	1 kg	eggs		T	O	1 kg	eggs	T	O
1998	SA	FR	IR						15 kg	eggs	T	W
1998	SA	FR	KZ						2 kg	eggs	T	O
1998	SG	FR	IR	8 kg	eggs		T	O	9 kg	eggs	T	O
1998	SG	FR	IR	4 kg	eggs		T	W	34 kg	eggs	T	W
1998	SG	FR	KZ						3 kg	eggs	T	O
1998	SG	FR	KZ	1 kg	eggs		T	W				
1998	SN	FR	IR						73 kg	eggs	T	W
1998	SN	FR	IR						8 kg	eggs	T	O
1998	TH	FR	IR						2 kg	eggs	T	W
1998	TN	FR	IR						5 kg	eggs	T	W
1998	TW	FR	IR						17 kg	eggs	T	W
1998	TW	FR	IR						1 kg	eggs	T	O
1998	US	FR	KZ						134 kg	eggs	T	O
1998	US	FR	KZ						20 kg	eggs	T	W
1998	US	FR	RU	154 kg	eggs		T	W				
1998	UY	FR	IR						2 kg	eggs	T	W
1998	ZA	FR	IR						1 kg	eggs	T	O
1998	ZA	FR	KZ						1 kg	eggs	T	O
1998	AR	GB	IR						2 kg	eggs	T	W
1998	BH	GB	AZ						43 kg	eggs	T	W
1998	BH	GB	AZ						11 kg	eggs	T	O
1998	CH	GB	RU	41 kg	eggs		T	O				
1998	HK	GB	IR						1 kg	eggs	T	W
1998	IS	GB	IR						2 kg	eggs	T	W
1998	JP	GB	IR						129 kg	eggs	T	W
1998	JP	GB	XX						575 kg	eggs	T	W
1998	MX	GB	IR						1 kg	eggs	T	W
1998	OM	GB	AZ						20 kg	eggs	T	W
1998	OM	GB	AZ						8 kg	eggs	T	O
1998	PH	GB	XX						59 kg	eggs	T	W
1998	QA	GB	AZ						51 kg	eggs	T	O
1998	QA	GB	AZ						4 kg	eggs	T	W
1998	SA	GB	IR						1 kg	eggs	T	W
1998	XX	GB	AZ						6 kg	eggs	T	W
1998	ZA	GB	KZ						60 kg	eggs	T	W
1998	CH	GR	IR	5 kg	eggs		T	O				
1998	US	GR	RU	20,400	eggs		T	O				
1998	US	GR	RU	63 kg	eggs		T	O				
1998	US	HK	RU	40.8 kg	eggs		T	O				
1998	AT	IR		8,250 kg	eggs		T	O				
1998	BE	IR		80 kg	eggs		T	W				
1998	CH	IR		3,205 kg	eggs		T	O				
1998	CH	IR		540 kg	eggs		T	W				
1998	CH	IR	RU	6,160 kg	eggs		T	W				
1998	CH	IR		20 kg	meat		T	W				
1998	CH	IR	RU	8	skins		T	W				
1998	DE	IR		2,446 kg	eggs		T	O				
1998	DE	IR		1167 kg	eggs		T	W				
1998	DK	IR		1,600 kg	eggs		T	W				

Year	Imp.	Exp.	imports reported			exports reported								
			Origin	Quantity	Unit Term	P	S	Quantity	Unit	Term	P	S		
1998	ES	IR		4430	kg	eggs	T	W						
1998	ES	IR		400		eggs	T	W						
1998	FR	IR		13,651	kg	eggs	T	W						
1998	FR	IR		1,020	kg	eggs	T	O						
1998	FR	IR		4,715	kg	meat	T	W						
1998	GB	IR		1,505	kg	eggs	T	W						
1998	IT	IR		4,000	kg	bodies	T	W						
1998	US	IR		1,500		eggs		W						
1998	US	IR		1,000	g	eggs		W						
1998	ID	NL	IR						11	eggs		T	W	
1998	ID	NL	RU						2	eggs		T	W	
1998	JP	NL	RU						700	eggs		T	O	
1998	JP	NL	RU						5	kg	eggs	T	O	
1998	ZA	NL	IR						1	kg	eggs	T	W	
1998	US	PE	RU	16,800	g	eggs	P	W						
1998	DE	PL	RU						3,050	kg	eggs			W
1998	DE	PL	RU	2,550	kg	eggs	T	O						
1998	DE	PL		25,000		live	T	C	25,000		live			C
1998	GB	PL	RU						340	kg	eggs			W
1998	US	PL	RU						1,430	kg	eggs			W
1998	US	PL	RU	300	kg	eggs	T	W						
1998	US	PL	RU	240	kg	eggs	T	O						
1998	US	PL	RU	40	kg	eggs	T	I						
1998	GR	RO		2	kg	eggs	T	W						
1998	US	RO		200		eggs	T	W						
1998	US	RO		58	kg	eggs	T	O						
1998	AE	RU							255	kg	eggs	T	W	
1998	AE	RU	KZ						263	kg	eggs	T	W	
1998	AZ	RU							106	kg	eggs	T	W	
1998	BE	RU		872	kg	eggs	T	W	870	kg	eggs	T	W	
1998	CH	RU		14,625	kg	eggs	T	W	865	kg	eggs	T	W	
1998	CH	RU		331	kg	eggs	T	O						
1998	CH	RU	AZ						405	kg	eggs	T	W	
1998	CH	RU	KZ						14,625	kg	eggs	T	W	
1998	DE	RU		3,833	kg	eggs	T	W	5,024	kg	eggs	T	W	
1998	DE	RU		455	kg	eggs	T	O	748	kg	eggs	T	O	
1998	DK	RU		395	kg	eggs	T	W	702	kg	eggs	T	W	
1998	FI	RU		50	kg	eggs	T	W	50	kg	eggs	T	W	
1998	FR	RU		627	kg	eggs	T	W	674	kg	eggs	T	W	
1998	FR	RU	KZ	840	kg	eggs	T	W	2,842	kg	eggs	T	W	
1998	JP	RU							3,025	kg	eggs	T	W	
1998	NL	RU		816	kg	eggs	T	W	2,082	kg	eggs	T	W	
1998	RO	RU							4	kg	eggs	T	W	
1998	US	RU		92	kg	eggs	T	I						
1998	US	RU		3,094	kg	eggs	T	W	3,630	kg	eggs	T	W	
1998	US	RU		500	kg	eggs	T	O	2,520	kg	eggs	T	O	
1998	US	RU		263		eggs	T	W						
1998	US	RU							25	kg	eggs	T	C	
1998	DE	SA	KZ	79	kg	eggs	T	O						
1998	CH	SG	IR	15	kg	eggs	T	O	15	kg	eggs	T	O	
1998	JP	SG	RU						53	kg	eggs	T	O	
1998	JP	SG	RU						53	kg	eggs	T	W	
1998	US	SG	RU	1		eggs	T	W						
1998	US	TK	RU	21	kg	eggs	T	W						

Year	Imp.	Exp.	Origin	imports reported			exports reported			P	S		
				Quantity	Unit	Term	Quantity	Unit	Term				
1998	CH	TR	RU	1,517	kg	eggs			1,519	kg	eggs	T	O
1998	JP	TR	XX						25	kg	eggs	T	O
1998	US	TR		390	kg	eggs						T	I
1998	US	TR		2	g	eggs						T	I
1998	US	TR	RU	3,401	kg	eggs						T	W
1998	US	TR	RU	2,098	kg	eggs			6,209	kg	eggs	T	O
1998	US	TR	RU	310		eggs						T	O
1998	US	TR	RU	2		eggs						T	I
1998	US	TR	RU	1	g	eggs						T	W
1998	US	TR	XX						2,909	kg	eggs	T	O
1998	US	TR	XX	18	kg	eggs						T	I
1998	AR	US	RU						53	kg	eggs	T	W
1998	AR	US	RU						13	kg	eggs	T	O
1998	AR	US	RU						4		eggs	T	O
1998	AU	US	RU						42	kg	eggs	T	O
1998	AU	US	RU						42	kg	eggs	T	W
1998	CA	US	CA						35		eggs	T	O
1998	CA	US	RU						1,264	kg	eggs	T	W
1998	CA	US	RU						339	kg	eggs	T	O
1998	CA	US	RU						139		eggs	T	W
1998	CA	US	RU						99		eggs	T	O
1998	CL	US	RU						20	kg	eggs	T	W
1998	DE	US	KZ						3,600	g	eggs	T	W
1998	DE	US	RU						21,770	g	eggs	T	O
1998	DE	US	RU						5,321	kg	eggs	T	W
1998	DE	US	RU	1,064	kg	eggs			410	kg	eggs	T	O
1998	DE	US	RU						616	g	eggs	T	W
1998	DK	US	RU						202	kg	eggs	T	O
1998	DK	US	RU						81	kg	eggs	T	W
1998	ES	US	RU						41	kg	eggs	T	W
1998	ES	US	RU	21	kg	eggs			15	kg	eggs	T	O
1998	ES	US	XX	36	kg	eggs						T	O
1998	FR	US	RU						128	kg	eggs	T	O
1998	FR	US	RU						91	kg	eggs	T	W
1998	GR	US	RU						230.4	kg	eggs	T	O
1998	GR	US	RU						49	kg	eggs	T	W
1998	GR	US	RU						20		eggs	T	O
1998	GR	US	SV						41	kg	eggs	T	W
1998	IT	US	RU						10.5	kg	eggs	T	O
1998	JP	US	RU						1	kg	eggs	T	W
1998	LC	US	RU						1	kg	eggs	T	W
1998	MC	US	RU						38	kg	eggs	T	W
1998	MC	US	RU						21	kg	eggs	T	O
1998	MX	US	RU						4	kg	eggs	T	W
1998	MX	US	RU						9	kg	meat	T	W
1998	NL	US	RU						42	kg	eggs	T	W
1998	NL	US	RU						41	kg	eggs	T	O
1998	NO	US	RU						225	kg	eggs	T	O
1998	PT	US	RU						167.4	kg	eggs	T	W
1998	PT	US	RU						105	kg	eggs	T	O
1998	RU	US	RU						55	kg	eggs	T	O
1998	SE	US	RU						42	kg	eggs	T	O
1998	SG	US	RU	27	kg	eggs			27	kg	eggs	T	W
1998	TR	US	RU						266	kg	eggs	T	W

Year	Imp.	Exp.	imports reported			exports reported				P	S	
			Origin	Quantity	Unit	Term	P	S	Quantity			Unit
1998	TR	US	RU					167	kg	eggs	T	O
1998	UY	US	RU					20	kg	eggs	T	W
1998	ZA	US	RU					14	kg	eggs	T	O
			Total:	121,274.7	kg	eggs		109,501.3	kg	eggs		

Huso dauricus Georgi, 1775

**Kaluga
Great Siberian Sturgeon**

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

Kaluga *Huso dauricus* is one of two species of Acipenseriformes endemic to the Amur River Basin, located in the Russian Federation and in China. It is one of the largest species of sturgeon and has a relatively low reproductive rate, reaching sexual maturity at 14-23 years and spawning every 3-5 years. The species has apparently been declining in numbers for more than a century. The main threat to the species appears to be over exploitation, but the Amur River is now being increasingly degraded by pollution from oil products, mineral fertilisers from agricultural development, and by-products of gold mining operations. Although the Amur River has not yet been affected by the construction of dams, plans for such construction exist and if achieved would render numerous spawning grounds inaccessible.

Historically, the highest sturgeon catches were recorded on the middle reach of the river, where both Chinese and Russian fisheries operate. However, as the two species *H. dauricus* and *Acipenser schrencki* Amur Sturgeon are rarely distinguished in harvest statistics, it is difficult to assess the impact of harvest on each species individually. Official catch records of *H. dauricus* in the former USSR and the Russian Federation dropped gradually from 595 t in 1881 to 61 t in 1948, and increased to 89 t in 1996. In 1998, the annual catch quota for *H. dauricus* was 309 individuals (legal minimum size limits of 200 cm and 65 kg) with a potential caviar production of 4,045 kg. Kaluga caviar reported in international trade in 1998 totalled 7,239 kg (4,481 kg from China and 2,758 kg from the Russian Federation). The Russian Federation decreased the caviar export quota for 1999 (3,500 kg compared to 3,600 kg for 1998), but increased it for 2000 (6,000 kg, plus a quota of 18,000 kg of meat). According to current available data, it appears that the caviar export quota was exceeded in the Russian Federation by 133 kg for 1999. China has set lower caviar export quotas of 3,430 kg per year for both 1999 and 2000. The domestic consumption of caviar in the Russian Federation is unknown and in China it is estimated to be 20% of the total national production. Captive breeding of this species has not yet been successful and *H. dauricus* restocking programmes and commercial captive breeding facilities therefore do not exist.

DISTRIBUTION AND POPULATION

The CITES database lists the distribution of *Huso dauricus* as: China, Russian Federation, former USSR (Anon., 2000a).

H. dauricus is classified as Endangered by IUCN (1996):
EN A1acde+ 2d China, Russia [Amur River].

The Amur River is 4,092 km long if its longest tributary, the Shilka River, is included (Krykhtin and Svirskii, 1997a). The middle reach of the Amur River, where it is bordered by both the Russian Federation and China, is 975 km long. It extends from the city of Blagoveshchensk (Heihe in China) to the mouth of the Ussuri (Wusuli) River between Khabarovsk (Russian Federation) and Fuyan (China), 966 km from the estuary (Krykhtin and Svirskii, 1997a).

H. dauricus is endemic to the Amur River basin, its tributaries and lakes. Young enter the Okhotsk Sea during the summer. The species distribution is fragmented with a population living in the estuary and coastal zones that is distinguishable from populations in the lower Amur, middle Amur and Zeya-Bureya River systems (Krykhtin and Svirskii, 1997b; Svirskii, 1971).

All populations have decreased considerably due to increased fishing since the 19th century when the highest catches were recorded, and the middle Amur population was the largest (Krykhtin and Svirskii, 1997b). At present, the estuary population is the most abundant and had increased by 35% since fishing restrictions were introduced in 1976. However by 1993, this population was estimated to have been reduced by 30-35%, due to illegal fishing (Krykhtin and Svirskii, 1997b).

During the latter part of the 1980s, Krykhtin and Svirskii (1997b) estimated population sizes using mark and recapture methods. The results show that the estuary population is the most abundant with approximately 70,000 individuals aged one year or more. Approximately 14% of this population was thought to be potentially sexually mature individuals, but by the early 1990s this percentage had decreased by about one third due to illegal catches (Krykhtin and Svirskii, 1997a). The lower Amur population was estimated to consist of approximately 40,000 individuals aged two years or older and the mid Amur population to consist of 30,000 such individuals. The Zeya-Bureya population was very small (Krykhtin and Svirskii, 1997a).

Current populations consist predominantly of young fish, with mature fish accounting for only 2-3% of the population (Krykhtin and Svirskii, 1997b). As a result of the species' late maturation and generally low reproductive rate, the population decline is expected to continue, especially in the middle Amur.

China: Two populations of the species occur in China, the largest one in the middle Amur River and the other in the brackish estuary waters of the Sea of Japan (Krykhtin and Svirskii, 1997b). Spawning sites are located at the mouth of the Ahaoxing, Pingyanghe, Quingdeli, Xiabaca, Xueshuiwen and Yadanhe rivers (Wei *et al.*, 1997).

Russian Federation: Four populations of the species occur in the Russian Federation, the two largest ones inhabiting the lower and middle reaches of the Amur River, and the two smallest ones occurring in the Burya and Zeya rivers, and the coastal brackish waters of the Sea of Japan and Sea of Okhotsk. The Zeya-Bureya population is thought to be on the verge of extinction having only single specimens recorded in the upper region of the mid Amur and lower regions of the Zeya, Shilka and Argun Rivers (Krykhtin and Svirskii, 1997b).

HABITAT AND ECOLOGY

H. dauricus is one of the largest freshwater fishes; its total size can exceed 5.6 m in length, 1 t in weight and it can live for over 80 years (Anon., 1997). Juveniles up to one year of age feed on invertebrates; from three to four years of age they switch to feed on adult fish and cannibalism is frequent (Krykhtin and Svirskii, 1997b). According to a survey of the *H. dauricus* spawning population conducted in China, at 220 cm few specimens had reached sexual maturity (13.8% for males and 31.6% for females). In a sample of fish taken in 1978-1979, the spawning population ranged from 12-54 years of age, and age correlated positively with length and weight. Fecundity ranged from 383,400-3,280,000, and the number of eggs per gram of body weight averaged 41.4 (Wei *et al.*, 1997). Krykhtin and Svirskii (1997b) reported that males of the estuary population reach sexual maturity at 14-21 years of age and females at 17-23 years. Males spawn every three to four years and females every four to five years. Many individuals of the middle Amur population reach sexual maturity much earlier than the estuary population.

The estuary population comprises 75-80% of the freshwater form and the remainder are the saltwater form. The latter winters in the freshwater zone, and migrates to the brackish water of the delta to the northern part of the Tatar Strait and the south-western part of the Sakhalin Gulf for feeding in June and July. It returns to the freshwater zone in autumn when the salinity increases. For spawning, most of the population migrates in winter to grounds up to 500 km from the river mouth, whilst others enter the mid-Amur River.

The lower Amur population feeds in the lower river; mature fish migrate and use the same spawning grounds and time frame as the estuary population. The middle Amur population inhabits an area that begins at about 950 km upstream of the river mouth (Krykhtin and Svirskii, 1997a). Spawning migration to the lower regions of the middle Amur take place in May and June.

In May and June, adults of the Zeya-Bureya population migrate to the upper Amur to spawn.

THREATS TO SURVIVAL AND DOMESTIC USE

In contrast to most large rivers, the Amur River has not been dammed. However, the revival of certain dam construction plans would threaten to block access to a great number of spawning sites (Birstein, 1993). Investigations on ovaries by Svirskii (1984, cited in Krykhtin and Svirskii, 1997a) showed that a

parasite *Polypodium hydrotorme* decreased the fecundity of *H. dauricus* by approximately 19%. In addition, water pollution (heavy metals, oil products, phenol, mineral fertilisers and gold mining by-products) in the Amur has increased in recent years from both the Russian and Chinese banks (Matthieson, 1993; Krykhtin and Svirskii, 1997b). However, studies of the effects of pollution on sturgeon have not been undertaken.

Information provided suggests that amendments to previous regulations and/or adoption of new legislation are in effect in both range States, China and the Russian Federation. However, enforcement measures appear to be lacking and numerous experts as well as government officials have reported increasing pressure from illegal fishing practices and criminal activities around sturgeon poaching and black markets that have been reported in a large part of the range (Medetsky, 2000; Winchester, 2000).

Official catch records of *H. dauricus* in the Russian Federation and the former USSR dropped from 595 tonnes in 1881 to 61 t in 1948 (Krykhtin and Svirskii, 1997b), and was 89 t in 1996 (Anon., 2000b). Russian catch data for 1996 suggest that the current proportion of *A. schrencki* versus *H. dauricus* in the Amur River catch is 9% and 91% respectively (Anon., 2000b). However, late 19th century records suggested a 50% composition for each species (Krykhtin and Svirskii, 1997a).

Between 1993 and 1997, meat of *H. dauricus* was observed for sale both in markets and in shops in Khabarovsk, Komsomolsk-on-Amur, Nikolayevsk-on-Amur and Vladivostok (Anon., 2000b; TRAFFIC Europe-Germany, *in litt.* to TRAFFIC Europe, January 1997). Official records in China indicate that the combined annual catches of *A. schrencki* and *H. dauricus* fluctuated inconsistently since the 1950s

Combined annual catch of *A. schrencki* and *H. dauricus* (tonnes)

1959	1969	1979	1987	1989	1997	1998	1999
47	15	130	452	286	136	149	141

Source: TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000

In China, all caviar of a wild source is from *A. schrencki* and *H. dauricus* (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). Heilongjiang Province is the only province in China where caviar is processed from wild specimens (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000). Most caviar produced in China is destined for the export market; domestic consumption represents less than 20% of the total caviar production in the country (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

Domestic consumption of *H. dauricus* caviar in China (kilogrammes)

	1993	1994	1995	1996	1997	Average
<i>H. dauricus</i>	498.0	492.0	878.0	416.0	452.0	547.2

Source: CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000

INTERNATIONAL TRADE

The listing of *H. dauricus* in CITES Appendix II entered into effect on 1 April 1998. Available complete trade data are therefore limited to nine months of trade (1 April to 31 December 1998).

Gross exports and the comparative tabulation of trade in *H. dauricus* for 1998 are given in the Appendix. CITES Annual Reports for 1998 show that China and the Russian Federation exported, respectively, 4,481 kg and 2,758 kg of caviar. The importing countries were Japan (4,995 kg), Germany (1,300 kg), USA (907 kg) and the United Arab Emirates (36 kg). In 1999, Russia exported a total of 3,632.8 kg of *H. dauricus* caviar, entirely destined for the USA (CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 September 2000).

Caviar export quotas established by the Russian Federation and China for *H. dauricus* (kilogrammes)

		1998	pre-Convention	1999	2000
Caviar	Russian Federation	3,600		3,500	6,000
	China	3,487	1,500	3,430	3,430
	Total	7,087		6,930	9,430
Meat	Russian Federation	0		0	18,000

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Only trade in 'caviar' has so far been detailed in CITES data. However, the Russian Federation has established an export quota of 18,000 kg of meat for *H. dauricus* for 2000.

There were no confiscations of *H. dauricus* reported in 1998, however smuggling of caviar on the Sino-Russian border seems to occur frequently during the migration season (Medetsky, 2000).

CONSERVATION MEASURES

The species is not fully protected in the Russian Federation or China.

China: The listing of *A. schrencki* and *H. dauricus* in the State Category of Protected Wildlife is expected to be adopted in the near future (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). The Heilongjiang Province authorities issued protection and management regulations, such as gear restrictions, harvest size, closed seasons and areas, and the requirement of a fishing license in the early 1950s. These were renewed in 1982 through "The Heilongjiang Ordinance on the Protection and Propagation of Fisheries Resources". However, the regulations have not been fully implemented (Wei *et al.*, 1997).

Restriction on fishing size

The Ordinance of 1982 prescribed the minimum size limits for *A. schrencki* at 100 cm or 4 kg, and for *H. dauricus* at 200 cm or 65 kg (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000). Sturgeon under these size and weight specifications must be released back in to the river (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Other restrictions

According to the "Protocol on Fisheries Resources Conservation, Regulation and Multiplication in Border Areas of Heilong (Amur) River and Wusuli River (Ussuri) of The People's Republic of China and the Russian Federation" adopted on 27 May 1994, fishing activities on the Heilong River are prohibited from mid-June to mid-July. The protocol also established areas where fisheries are prohibited; the mainstream of the Heilong River from Dagangzi, Luobei County, to Saniangkou where the Heilong and Songhua Rivers converge, were designated as no-fishing zones by the Government of China. The area where fishing is prohibited stretches over 1,270 km, from Fuyan (downstream) to Heihe (or Aihui; Blagoveshchensk on the Russian side) (upstream). These measures protect natural breeding sites, survival of juvenile fish and propagation of *A. schrencki* and *H. dauricus*.

The catch of any specimen of *H. dauricus* and *A. schrencki* on the Chinese side of the Heilong (Amur) River requires prior acquisition of a fishing license. In 1991, 2,248 sturgeon fishing licenses were issued, and in 2000, the number was reduced to 1,850 (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Restocking of juveniles

Breeding of *H. dauricus* in captivity has not yet been successful in China (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

Determination and allocation of quotas by local governments

Based on the average annual production from 1986 to 1988, the average size of fish landed, the decreasing rate of caviar production and the hydrological resource and migration conditions, Chinese scientists calculated Amur River sturgeon (*A. schrencki* and *H. dauricus*) annual catch quotas for the period 1989-1997. This quota was revised in 1997 and the Heilongjiang Province Fisheries Bureau

adopted a new total catch quota of 105.4 t for 1998. The latter is composed of: 1,888 individuals for *A. schrencki* with an expected caviar production of 3,235 kg; and 309 individuals for *H. dauricus* with an expected caviar production of 4,045 kg (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, Sept. 2000).

Processing and export company management

Caviar processing companies are required to register with the Heilongjiang Provincial Fisheries Bureau. In 1998, six companies were registered as sturgeon products processors, all established in Heilongjiang Province. Six other companies were registered as caviar exporters: four established in Heilongjiang Province, one in Beijing and one in Dalian Free Trade Zone (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000). According to the Ministry of Agriculture's "*Regulation on Special License of Aquatic Wildlife Utilization, the People's Republic of China*", caviar processing companies must acquire special qualification from the local Fisheries Bureau, including a "special (caviar) purchase license" and a "special (caviar) processing license". Caviar purchase and processing are controlled by the local Fisheries Bureau. In principle, the Fisheries Bureau will gradually decrease the number of licenses and also monitor those companies with special purchase and production licenses. According to the "*Law of Wildlife Protection of the People's Republic of China*", the Ministry of Agriculture only issues caviar trade licenses to a limited number of import and export companies. The Ministry of Agriculture strictly controls companies that receive a special license.

Russian Federation: A prohibition on sturgeon catch throughout the USSR was adopted in 1923 and then withdrawn in 1930. A ban on the catch of *A. schrencki* and *H. dauricus* was introduced in 1958. This prohibition is still in effect (Krykhtin and Svirskii, 1997a), however a tolerance called "controlled catch" for incidental and scientific catches is allowed. These catches are the current source of caviar and sturgeon meat from the Amur River. The "controlled catch" is apparently not well defined and difficult to control (Anon., 2000b).

Russian catch quotas for 1999 and 2000 were adopted by the Federal Government following the recommendations set by the State Fisheries Committee in consensus with an independent body of experts, at 15 t per year for *A. schrencki* and 65 t per year for *H. dauricus* (Anon., 2000b).

CAPTIVE BREEDING

Both the Russian Federation and China are constructing sturgeon hatcheries on the Amur River, but aquaculture of *H. dauricus* specifically is not reported (Krykhtin and Svirskii, 1997a). In China, experimental captive breeding is carried out in governmental and private facilities. In 2000, the Heilongjiang Province Fisheries Bureau authorised eight stations to undertake research on the captive breeding of *A. schrenckii* and *H. dauricus* (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database and China has not reported the results of their research on sturgeon captive breeding and the development of commercial farms.

Sturgeon hatcheries and farms are developing rapidly world-wide. These have led to the successful production of juveniles for the rehabilitation of wild populations, as well as significant production of sturgeon meat and caviar. In China, sturgeon aquaculture started in 1957 and has developed quickly since the 1980s; most species bred in captivity in China are sturgeon species occurring in the Russian Federation and *A. schrencki* is recognised as an important species for the sturgeon aquaculture industry (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000). The captive breeding of *H. dauricus* has not yet been successful (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000), but research stations are carrying out experimental breeding of the species in Heilongjiang Province (TRAFFIC East Asia, *in litt.* to TRAFFIC Europe, September 2000).

REFERENCES

Anon. (1997). Document Doc. 10.89; Prop. 10.65. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.

- Anon. (2000a). <http://www.cites.org/eng/dbase/fauna->
- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia, Field investigations. Unpublished report, Pp. 23.
- IUCN. (1996). *1996 Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- Krykhtin, M.L. and Svirskii, V.G. (1997a) Sturgeon of the Amur River, Kaluga *Huso dauricus* Georgy and Amur sturgeon *Acipenser schrenckii* Brandt. In: Birstein, V.J., Waldman, J.R., and Bemis, W.E. 1997. Sturgeon biodiversity and conservation. Kluwer Academic Publishers, Dordrecht. *Environment Biology of Fishes* 48:231-239.
- Krykhtin, M.L. and Svirskii, V.G. (1997b). Sturgeon catch and the current status of sturgeon stocks in the Amur River. In: Birstein, V.J., Bauer, A. and Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. pp. 29-34.
- Matthieson, P. (1993). The last cranes of Siberia. *The New Yorker*, May 3 1993:76-86.
- Medetsky, A. (2000). Black caviar feeds thousands on Amur. *Vladivostok News*, 15 September 2000.
- Svirskii, V.G. (1971). The Amur River Sturgeon and Kaluga. *Uchenye Zapiski dalnevostochnogo Gosudarstvennogo Universiteta*, 15:19-33.
- Wei, Q., Ke, F., Zhang, J., Zhuang, Luo, J., Zhou, R. and Yang, W. (1997). Biology, fisheries and conservation of sturgeons and paddlefish in China. In: Birstein, V.J., Waldman, J.R., and Bemis, W.E. 1997. Sturgeon biodiversity and conservation. Kluwer Academic Publishers, Dordrecht. *Environment Biology of Fishes*. 48:241-255.
- Winchester, S. (2000). *National Geographic*, February 2000. Pp. 7-14.

Gross exports of, and export quotas for *Huso dauricus* 1998

TAXON	TERM	UNIT	Exporter	1998	Quotas 1998 (kg)
<i>Huso dauricus</i>	eggs	kg	CN	4,481	3,487 + 1,500 pre-Convention
<i>Huso dauricus</i>	eggs	kg	DE	0	
<i>Huso dauricus</i>	eggs	kg	RU	2,758	3,600

Comparative Tabulation of trade in *Huso dauricus* 1998

Year	Imp.	Exp.	Origin	Imports reported				Exports reported					
				Quantit y	Unit	Term	P	S	Quantit y	Unit	Term	P	S
1998	AE	CN						36 kg	eggs	T	W		
1998	DE	CN		1,100 kg	Eggs	T	O	1,100 kg	eggs	T	O		
1998	DE	CN						200 kg	eggs	T	W		
1998	JP	CN						2,629 kg	eggs	T	W		
1998	US	CN		516 kg	eggs	T	W	516 kg	eggs	T	W		
1998	JP	RU						2,366 kg	eggs	T	W		
1998	US	RU		392 kg	eggs	T	W	391 kg	eggs	T	W		
TOTAL				2,008 kg				7,238 kg					

Huso huso Gray, 1827

Beluga
Giant Sturgeon

Order: ACIPENSERIFORMES

Family: ACIPENSERIDAE

SUMMARY

Beluga *Huso huso* has a wide distribution. It occurs in the Caspian Sea, Black Sea and the Sea of Azov and many of the tributaries of these seas including the Volga and Danube rivers. It is the largest species of Acipenseriformes reaching a length of six metres and a weight of more than one tonne. The Volga River is the major spawning river for the species. However the construction of dams on virtually all these rivers has significantly reduced the spawning area. Currently, 96% of *H. huso* in the Volga River consist of hatchery-reared fish. Captive breeding does not compensate for the loss of natural reproduction and *H. huso* continues to decline. In addition to loss of habitat, the species has been subject to overfishing and egg-production has been disrupted due to environmental pollution.

Historically, *H. huso* were harvested more intensively than Russian Sturgeon *Acipenser gueldenstaedtii* or Stellate Sturgeon *A. stellatus* (Khodorevskaya *et al.*, 1997). The catch of *H. huso* in the northern Caspian Sea dropped from approximately 2,800 tonnes (t) in 1970 to less than 300 t in 1994. Available data suggest that the Caspian Sea catch of *H. huso* in 1997 was 339 t. The species was always abundant in the Danube River, with an average of 23 t caught annually in the middle Danube River from 1972 to 1976. This catch decreased to 7.5 tonnes per year from 1985 to 1989 after the construction of Djerdap dams I and II. More recently, recorded landings from the Lower Danube River in Romania and Bulgaria decreased from 62.1 t in 1997 to 42.7 t in 1999. The importance of domestic markets of caviar and sturgeon meat is unknown in the Russian Federation. In Iran, for the past five years, the domestic consumption has been estimated at 5% and 65% of the country's respective production of caviar and meat. In 1998, exports of *H. huso* totalled 21,267 kg of caviar (8,035 kg from the Russian Federation, 7,033 kg from Kazakhstan and 4,013 kg from Iran) and 15,755 kg of meat and 22 skins from Iran, destined for Italy and Switzerland, respectively. Assuming that 6-7 kg of caviar is obtained per 100 kg of sturgeon caught (both male and female), total caviar exports for 1998, i.e. 21.3 t, represents a catch of 304-354 t. Caviar export quotas for 1998 were set at 5 t by Iran, 2.5 t by Bulgaria and 5 t by the Russian Federation. For 1999, the Russian Federation's caviar export quota was set at 3 t; export quotas were exceeded in 1998 and 1999 by 3 t and 2.7 t respectively. The combined export quotas set by range States have increased from 16.6 t to 17.2 t of caviar, and from 15.5 t to 111 t of meat for 1999 and 2000, respectively.

The species is recommended under Decision 11.106 for inclusion in category 1 (i).

DISTRIBUTION AND POPULATION

The CITES database lists the distribution of *H. huso* as: Azerbaijan, Bulgaria, Croatia, Czech Republic? (ex), Georgia, Hungary, Iran (Islamic Republic of), Italy (ex), Kazakhstan, Moldova, Republic of, Romania, Russian Federation, Slovenia, Turkey, Turkmenistan, Ukraine, Yugoslavia (Anon., 2000a).

H. huso is classified as Endangered by IUCN (1996):

EN A2d Azerbaijan, Bulgaria, Hungary, Iran, Italy, Kazakhstan, Moldova, Romania, Russian Federation, Turkmenistan, Ukraine, Yugoslavia

Sea of Azov stock: CR A1acde+ 2d Russia [Mediterranean and Black Sea]

Black Sea stock: EN A1acde+ 2d Hungary, Romania, Russian Federation, Ukraine, Yugoslavia [Mediterranean and Black Sea]

Caspian Sea stock: EN A1acde+ 2d in Azerbaijan, Iran, Kazakhstan, Russian Federation, Turkmenistan

Adriatic Sea stock: EX Italy [Mediterranean and Black Sea].

H. huso is historically widespread, inhabiting the Caspian and Black seas and the Sea of Azov. The most recent record of the species in the Sea of Azov is from the mid 1980s (Volovik *et al.*, 1993), when *H. huso* was believed to make up 3% of the total sturgeon biomass. It is now believed extinct in the Adriatic where historically, solitary individuals occurred. The last report of the species in the Po River Basin is from 1972 (Rossi *et al.*, 1991).

A seasonal migration occurs in the Caspian; in the summer the main feeding grounds are in the north and in the winter in the south (Barannikova *et al.*, 1995). There are no estimates of the total population size. The main spawning population enters the Volga; the other spawning populations of the Caspian are thought to be minimal (Khodorevskaya and Novikova, 1995; Levin, 1997; Vlasenko, 1990).

Azerbaijan: Historically the species spawned in the Kura River, but more recent information suggests that *H. huso* no longer use the Kura River (Khodorevskaya *et al.*, 1997; Levin, 1997). The coastal waters of Azerbaijan are important feeding grounds for *H. huso*, which feeds primarily on fish such as sprat and *kilka* (Clupeidae) (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Bulgaria: Part of the Black Sea population, *H. huso* in the Danube River breed in spring (early April - when the first catch of the year are recorded) and in autumn (end-September) (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Czech Republic: *H. huso* is extinct, with the last specimen being caught in the Morava River at Lanžhot at the beginning of 20th century (Holcík, 1995).

Georgia: The southeast coast of the Black Sea is an important feeding and wintering area for sturgeon (*H. huso*, *Acipenser stellatus*, *A. nudiiventris*, *A. persicus colchicus* and *A. sturio*) that migrate upstream in a number of rivers for spawning. The main rivers used are the Supsa, Inguri, Chorokhi, and particularly the Rioni (Zarkua and Tsuladze, 1999).

Hungary: Historically, Hungary formed the main spawning ground for this species in the Danube, but catches began to decline from the 16th century onwards. Very few were taken during the early 20th century. The last specimen caught in 1987 was an exceptional rarity and the species is regarded as extinct in Hungary (Pintér, 1991).

Iran: Razavi (1988) notes that *H. huso* enters the Sefidrud, Tajen and Gorganrud rivers in the southern shores of the Caspian Sea for spawning. This species is considered one of the most valuable sturgeon species of Iran. It migrates during February to April, but sometimes is also seen in these rivers during the autumn months. However, Dr J. Holcík (*in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000) maintains that the Mangil Dam on the Sefidrud River prevents sturgeon from entering this river where all spawning grounds have been destroyed due to heavy pollution and water extraction (pumping). Hensel and Holcík (1997) suggest that *H. huso* no longer use the Sulak River for spawning. According to Dr M. Pourkazemi (*in litt.* to IUCN/SSC Wildlife Trade Programme, 25 September 2000), among the Mangil Dam and other dams constructed on the Sefidrud River, there is a remaining distance of 45 km from the dam closest to the estuary, and fish are still observed entering the river to spawn.

Kazakhstan: *H. huso* spawns naturally in the Ural River and since 1979, the numbers entering the Ural have exceeded those entering the Volga (Khodorevskaya *et al.*, 1997).

Moldova: The Moldovan range on the Danube River is 811 m. Sturgeon also occur in the Prut River, a tributary of the Danube River (junction at km 137th) (Anon., 2000c). In the early 1990s, *H. huso* had not been recorded for two decades (Chepurnova, 1991).

Romania: The Danube River has the only current *H. huso* spawning population in the Black Sea Basin. However, the population has been extirpated in the upper part of the river and is critically endangered in the central region because of overfishing, pollution and damming (Bacalbasa-Dobrovici, 1997; Hensel and Holcík, 1997; Suciú *et al.*, 1998).

Russian Federation: The largest population is believed to occur in the Volga River (Vlasenko, 1990). Levin (1997) estimated that the spawning population entering the Volga in recent years is approximately 8,000-9,000 individuals, up to 25% of which are female. The decline in commercial catches from this region (northern Caspian Sea) from 2,650 t in the 1970s to 460 t in 1993, reflect a decline in the population. Recent information suggests that *H. huso* no longer use the Terek River (Khodorevskaya *et al.*, 1997). In the Sea of Azov, some natural spawning grounds are no longer accessible due to the construction of dams on

the Don and Kuban rivers (Anon., 2000b). The biomass of Beluga in the Sea of Azov, was estimated to be around 1,770 t in the mid-1980s, before a mass die-off of various sturgeon species (Volovik *et al.*, 1993).

Slovakia: The last *H. huso* recorded in the Slovakian-Hungarian portion of the Danube was in 1925 (Hensel and Holcák, 1997).

Slovenia: There are no records of *H. huso* occurring in Slovenia (Povž and Sket, 1990).

Turkey: The Coruh River is used for spawning by *H. huso*, and unidentified sturgeon species have been reported in small numbers in the Sakarya and Mert Rivers (Edwards and Doroshov, 1989). In the late 1980s, local fishermen estimated that only five to six gravid female sturgeon could be taken from the Yesilirmak to supply a hatchery and up to 20 from the Kizilirmak (Edwards and Doroshov, 1989).

Turkmenistan: No information provided.

Ukraine: *H. huso* is known from the east and west coasts of the Crimea. It is also known from the north-western part of the Black Sea, from where it enters the Danube, Dniester, Dnieper, and occasionally the Bug rivers. Its distribution was limited in the last century by pollution and the damming of these rivers (Pavlov, 1980). The biomass of *H. huso* in the Sea of Azov was estimated to be around 1,770 t in the mid 1980s, before a mass die off of sturgeon (Volovik *et al.*, 1993).

Former Yugoslavia: Extinct in Croatia, *H. huso* was formerly recorded in the Sava River at Zagreb and also in its tributary, the Kupa River (Glowacki, 1896; Munda, 1926; Taler 1954). It is Endangered in Serbia. Access to spawning grounds has been prevented by the Djerdap I and Djerdap II dams, and catch has dramatically decreased (Jankoviæ, 1993; 1996; Simonoviæ and Nikoljæ, 1996). There have been no records from Slovenia since at least 1990 (Povž and Sket, 1990).

Introduced populations: In the Baltic Sea river estuaries east of Rostock (Germany), and further upstream, hybrid sturgeon specimens have been caught. They probably originated from upstream aquaculture farms and/or aquaria (i.e. specimens that had grown too large to be kept in aquaria) (Jörn Gessner, Scientist at the Institute of Freshwater Ecology and Inland Fisheries, pers. comm. to TRAFFIC Europe, 15 March 2000).

HABITAT AND ECOLOGY

H. huso is the largest of all sturgeon species. It can reach a total length of six metres and more than one tonne in weight (Berg, 1948). Khodorevskaya and Novikova (1995) reported that the mean length of immature males is 2.25 m and females 2.6 m. The maximum reported age is 118 years (Babushkin *et al.*, 1964, cited in Pirogovskii *et al.*, 1989). Sexual maturity is reached at 10-16 years for males and 14-20 years for females; the average size at first spawning is 2 m and 50 kg (Hochleithner and Gessner, 1999).

H. huso is piscivorous and begins to prey upon other fish in an early life stage. The species is anadromous and enters rivers flowing into the Black, Caspian and Sea of Azov to spawn. There are two forms; one of which spawns in winter and one in spring. The spring form migration peaks in summer and ends in late autumn. Fish migrating in winter are the largest of the two forms. They hibernate and reproduce the following year whereas the spring form migrates and reproduces during the same year. Unlike *A. gueldenstaedtii* and *A. stellatus*, *H. huso* do not congregate and remain dispersed in winter and summer (Levin, 1997).

Based on catches and caviar production in the late-1970s, the rate of caviar extracted per fish caught was estimated at 7 kg per 100 kg of the total catch (including males and females) for the three main commercial species of the northern Caspian Sea, *H. huso*, *A. gueldenstaedtii* and *A. stellatus* (Doroshov and Binkowski, 1985, cited in Williot and Bourguignon, 1991).

Iran: On the Iranian coast of the Caspian Sea, *H. huso* is primarily found in the eastern regions and in the Turkmen and Gomeeshan ports (Ghazel, 1993, cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC, Wildlife Trade Programme, September 2000). Sexual maturity in the southern shores of the Caspian Sea is attained

at an average age of 21 years in females (min. 10 years and max. 44 years) and at an average age of 14 years in males (min. 10 and max. 23 years) (International Sturgeon Research Institute, 1998; cited in Dr M. Pourkazemi, *in litt.* to IUCN/SSC, Wildlife Trade Programme, September 2000).

Kazakhstan: Until 1993, the mouth of the Ural River was dredged annually to allow vessels to travel upstream to Atyrau (former Gur'yev) harbour. Since 1993, this annual management of the river channel has not been carried out and is an obstacle to sturgeon migration (Anon., 2000c).

THREATS TO SURVIVAL AND DOMESTIC USE

Since the 1950s, all northern and western Caspian Sea tributaries, with the exception of the Ural River, have been dammed for hydroelectric power production, hence virtually all spawning grounds of the Caspian, Black and Sea of Azov regions have been lost. Approximately 90% of spawning grounds in the Caspian Sea have been lost and 91% of each generation now originates from hatcheries (Barannikova *et al.*, 1995). In the Sea of Azov, 100% of each generation is thought to consist of farm grown fish (Volovik *et al.*, 1993). There are also high levels of pollution in most rivers within its range. *H. huso* are particularly threatened by overfishing since Beluga caviar is highly prized and the species is relatively slow to reach maturity. In the mid 1990s, a period of open sea fishing which caught young and immature sturgeon as by-catch is likely to have destroyed a major component of future sturgeon stocks (Anon., 1997).

The commercial sturgeon catch in the Caspian Sea Basin peaked at 27,300 t in 1977, with *A. gueldenstaedtii* being the most abundant species (79%, 21,550 t), followed by *A. stellatus* (17%, 4,650 t). The remaining 4% (1,100 kg) was shared between various sturgeon species including *H. huso* (Levin, 1997). The catch of *H. huso* in the northern Caspian Sea dropped from approximately 2,800 t in 1970 to less than 300 t in 1994 (Khodorevskaya *et al.*, 1997). The Caspian Sea catch of *H. huso* was 339 t in 1997. The species was formerly abundant in the Danube River, with an average annual catch of 23 t in the middle Danube River from 1972 to 1976. This decreased to an average of 7.5 t/year from 1985 to 1989 after the construction of Djerdap dams I and II (Hensel and Holcik, 1997). More recently, recorded catch from the Lower Danube River in Romania and Bulgaria decreased from 62.1 t in 1997 to 42.7 t in 1999.

In the former Soviet Union and Iran, strict sturgeon fishery legislation had been in effect for decades until the emergence of the three new independent States in the Caspian Sea Basin in 1992 (Azerbaijan, Kazakhstan and Turkmenistan) (De Meulenaer and Raymakers, 1996). Current information suggests that amendments of previous regulations and/or adoption of new legislation are now in effect in most new range States in the Caspian Sea, as well as in the Black Sea (including the Danube River) and the Sea of Azov basins. In the latter, Bulgaria, Romania and Ukraine are the most important sturgeon fishing nations. However, enforcement measures appear to be lacking and numerous experts as well as government officials have reported an increasing pressure of illegal fishing practices and criminal activities surrounding the caviar trade in much of the range (Anon., 2000b; Dobbs, 1992; Doward, 2000; Evtouchenko, 1997; King, 1998; Ward, 2000a; 2000b).

Azerbaijan: The Vavarin reservoir on the Kura River prevents sturgeon from accessing spawning grounds. Lakes of deposited oil occur in industrial regions; the number of these is expected to increase with the rising sea level (Dumont, 1995). The waters of this country are particularly important winter feeding grounds. Along the northern and southern coast of Azerbaijan, within the 200-nautical mile Exclusive Economic Zone, *H. huso* is caught as by-catch in stationary nets and *kilka* (Clupeidae) fishing devices. Young and adult fish are caught by coastal fisheries during migration. Juvenile fish comprise up to 17.7% of the by-catch (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Species composition of the total catch in the 1990s

	<i>A. persicus</i> & <i>A. gueldenstaedtii</i>	<i>A. stellatus</i>	<i>Huso huso</i>	<i>A. nudiiventris</i>
Catch	63.47%	30.77%		5.76%
By-catch	78.50%	16.80%	3.70%	1%

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Adult *H. huso* are caught in spring and autumn. This fishery is carried out in low salinity waters.

Annual recorded landings of *H. huso* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
H. huso	27.0	21.0	49.00	74.60	33.30	29.50	24.10	22.30	13.1	20.00
	0	0							0	
Total	74.0	88.2	111.0	240.1	201.2	162.5	120.3	128.6	82.4	108.9
	0	0	0	0	0	0	0	0	0	0

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

Bulgaria: Wild sturgeon populations are threatened by a decrease in the number of spawning grounds on the Danube River as a result of human development, particularly hydroelectric facilities. The long lasting unregulated fisheries have disrupted the age composition of the populations due to selective catch of sexually mature fish, and to incidental catch of young migrating sturgeon. Water pollution has also influenced the metabolism of fish (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

The largest part of the Bulgarian sturgeon catch is composed of *H. huso*. This share has decreased progressively from 92% in 1995 to 73% in 1999.

Annual sturgeon catch in Bulgaria (tonnes)

	1995	1996	1997	1998	1999
H. huso	13.6	28.8	42.2	43.5	37.0
Total sturgeon catch	14.7	32.5	49.8	55.9	50.5
n° of licenses issued	0	0	122	155	169

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Domestic consumption of caviar in Bulgaria is limited to 7-10% of the total caviar production of Bulgaria. The entire sturgeon meat production is consumed within Bulgaria.

Georgia: During the fishing seasons, particularly coastal anchovy fisheries, numerous sturgeon juveniles are caught as by-catch. The estimated number of sturgeon caught (no species specific data available) declined from 77,000 individuals in 1977, to 57,000 in 1980 and to 27,000 in 1986. During the 1990s, illegal fishing in Georgia expanded rapidly as a result of poor control (Zarkua and Tsuladze, 1999).

Iran: For the past 5 years, the Iranian domestic market of caviar and sturgeon meat is estimated at 5% and 65% of the country's total production, respectively (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Annual catch of *H. huso* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
H. huso	257.4	249.4	198.4	137.8	140.1	137.6	167.1	135.4	139.6	117.6	14.1
Total	2,296.3	2,315.1	2,058.0	1,462.1	1,380.9	1,216.3	1,310.0	1,043.5	1,127.9	923.3	182.4

* Catch in spring 2000 only.

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

The annual production of *H. huso* meat and caviar in Iran has decreased significantly (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Sturgeon meat and caviar production for *H. huso* in 1992 and 1999 (tonnes)

	1992				1999			
	Meat	% *	Caviar	% *	Meat	% *	Caviar	% *
H. huso	162.0	10.1	6.2	2.4	96.0	13.1	4.2	4.2
Total	1,604.2		262.3		730.2		99.3	

* Percentages relate to the share of *H. huso* products of the total Iranian sturgeon fisheries.

Source: Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000

The above tables indicate that the total tonnage of meat and caviar of *H. huso* produced in Iran comprised 85% of the country's annual catch in weight in 1992 and 1999.

Kazakhstan: Oil fields are developing rapidly in the northern Caspian Sea, especially the Tengiz oil field in Kazakhstan (Sagers, 1994). Pollution from this industry affects spawning and feeding grounds. Radioactive contamination from a nuclear reactor also poses a threat (Dumont, 1995).

Annual catch of *H. huso* (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>H. huso</i>	330.0	288.0	280.0	389.0	209.0	321.5	124.6	95.3	80.0	53.0	54.7*
Total	1,963.0	1,767.0	1,437.3	1,109.0	556.0	573.0	367.8	389.5	490.0	279.0	230.5

* Spring catch.

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 September 2000

Moldova: Poaching is the main threat (Vedrashko *et al.*, 1998).

Romania: The Danube River population is threatened by the Djerdap I and II dams which prevent access to spawning grounds. This population is also under threat by other anthropogenic factors which have caused dramatic environmental changes in the Black Sea and the Danube River (Bacalbasa-Dobrovici, 1997).

Annual catch of *H. huso*, including the Danube River, Danube Delta and the Black Sea (kilogrammes)

	1991	1992	1993	1994	1995	1996	1997	1998	1999
	2,017	1,138	810	6,787	7,051	10,724	19,903	7,736	5,701

Source: Dr N. Patriche, *in litt.* to IUCN/SSC Wildlife Trade Programme, 5 September 2000

Scientific surveys concluded that the population of *H. huso* is quite well represented in Danube River. From 1991 to 1999, about 51% of the total Romanian sturgeon catch in the Danube River were composed of *H. huso*.

Russian Federation: The construction of the Volgograd Dam decreased the range area and spawning ground to only 12% of the former area available in the Volga River. Similarly, available spawning grounds in the Terek and Sulak rivers have been reduced to 132 ha and 202 ha respectively. Flow regulation of the Kuban River has caused the loss of approximately 140,000 ha of breeding ground and the Tsymlyansk Reservoir construction on the Don River in 1952 has removed approximately 68,000 ha of spawning ground (Volovik *et al.*, 1993).

The current spawning population consists mainly of fish that hatched after the alteration of the Volga River flow. Fish growth here appears to be retarded and the weight has almost halved in the 20 years from the early 1970s to the early 1990s (Khodoresvskaya *et al.*, 1993). Pollution has caused anomalies and may have rendered 100% of *A. stellatus* larvae in the Volga River non-viable (Shagaeva *et al.*, 1993). This effect may also be applicable to *H. huso*.

Catch of *H. huso* in the Caspian Sea (tonnes)

	1992	1993	1994	1995	1996	1997
Catch	746	247	157	94	109	86

Source: Caspian Fisheries Research Institute (KaspNIRKh), cited in Anon., 2000b

From 1992 to 1997, about 50% of the Russian catches of sturgeon in the Caspian Sea was composed of *H. huso* (Anon., 2000b).

The domestic consumption of sturgeon products in the Russian Federation is unknown, but is said to be the main form of consumption of illegal sturgeon fisheries (Mr Roman V. Ivakhnenko, State Fisheries Committee, pers. comm. to TRAFFIC Europe-Russia, September 2000). Based on the number of sets of illegal fishing

equipment recorded by enforcement agencies of the Caspian Sea and the Sea of Azov, the illegal sturgeon catch was evaluated to be six to ten times the legal catch (Anon., 2000b). In 1997, the estimated volume of the Moscow sturgeon meat market was three times the annual catch quotas for sturgeon species (Anon., 1998).

Turkey: The Yesilirmak River changed dramatically due to the construction of two dams in 1979, the lowest of which limits upstream migration. The Kizilirmak was dammed in 1998 and is heavily polluted. Habitat loss was predicted due to canalisation, and the spawning ground area has been drastically reduced. From 1969 to 1979, the Turkish annual sturgeon landings ranged from 310 t to 10 t, with an annual average of 112 t (Edwards and Doroshov, 1989). In Samsun, a Turkish landing site on the Black Sea, a caviar processor claimed in 1989 that his business had decreased from 8 t/year of caviar between the 1940s and the 1970s to 0.2 t/year at the end of the 1980s. In 1989, sturgeon were caught as by-catch by illegally operating trawlers along the Turkish coast at a distance greater than 3 nautical miles from the shore, but also illegally within the 3 nautical mile limit. This represented the main fishing pressure on sturgeon. In April 1989, *A. stellatus*, *A. gueldenstaedtii*, *A. sturio* and *H. huso* were observed in Samsun. The investigators were told that *A. nudiventris* was also landed in Samsun (Edwards and Doroshov, 1989).

Ukraine: The population is threatened by pollution in the Dniester River. Catch of *H. huso* in Ukrainian waters is prohibited since the species was listed in the Red Data Book of Ukraine in 1992 (Anon., 2000c).

Yugoslavia: The Danube River spawning grounds are no longer accessible due to the construction of the Djerdap dams. The catch of *H. huso* in the region of Djerdap, on both the Yugoslav and Romanian sides of the Danube River, dropped from 8,596 kg in 1989 to 3,994 kg in 1994 (Jankoviæ, 1999). Sturgeon fishing is now limited to 17.8 km of fishing ground from the Djerdap II dam to the border, close to the Timok River mouth (Anon., 2000c) (see "Conservation Measures").

INTERNATIONAL TRADE

The listing of *Huso huso* in CITES Appendix II entered into effect on 1 April 1998. Complete data for this species are therefore limited to nine months of trade (April-December 1998).

Gross exports of *H. huso* are given in the Appendix. According to 1998 CITES data, gross exports of *H. huso* caviar (Beluga) from range States totalled 21.3 t. The main exporters were Russia (8 t), Kazakhstan (7 t), Bulgaria (1.7 t) and Iran (1.3 t). The main importers of Beluga caviar in 1998 were: Switzerland (9 t), USA (8.5 t), France (4 t) and Germany (3.5 t).

In 1998, Iran also exported a variety of other Beluga products: 15.8 t of meat and 22 skins (probably processed into leather). Most of the meat was imported into European countries (Italy, Switzerland and France in decreasing order of importance). In 1999, Iran's exports included meat of wild origin for a total of 49.6 t. However, not all Annual Reports for 1999 have been submitted and hence this figure may under-reflect the actual volume of its trade in meat. Other *H. huso* products exported from Iran in 1999 included 30 kg of swim bladders imported by Germany to be processed into isinglass (fish glue used to clarify beer, wine and champagne, and for use in antique restoration). Iran did not report exports of any sturgeon specimens in its 1999 Annual Report.

Exports of *H. huso* from Russia in 1999

	Quantity (kg)	Description	Importer	No of permits
<i>H. huso</i>	5,738.27	Caviar	US, DK, BE, FR, DE, CH	11 Ex. + 7Re-ex.
	617.00	Meat	US	2 Export
Hybrids:				
<i>H. huso</i> x <i>A. ruthenus</i>	62.80	Live eggs	PL, DE, JP, CN, LV, US, HU, IL, GR	16 Export
	300.00	Live fish	KR	1 Export

Source: CITES Management Authority of the Russian Federation, *in litt.* to TRAFFIC Europe, 18 September 2000

Exports of *H. huso* caviar from Bulgaria in 1998 and 1999 (kilogrammes)

	1998	1999	Jan.-Sept. 2000
Caviar of <i>H. huso</i>	1,392	2,213	901

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Export quotas for *H. huso* and its hybrids (kilogrammes)

	1998		1999		2000		
	Caviar	Meat	Caviar	Meat	Caviar	Meat	Fert. Eggs
Azerbaijan			434	13,000	700	7,000	
Bulgaria	2,500		2,400		2,500		
Iran	5,000		3,000		3,000		
Kazakhstan			6,000		3,600	56,000	
Romania			1,750	2,500	3,200	35,000	
Russian Federation	5,000		3,000		3,500	13,000	10
<i>H. huso</i> x <i>A. ruthenus</i>							90
Turkmenistan					700		
Annual total	12,500	0	16,584	15,500	17,200	111,000	100

Source: CITES Notification Nos. 1998/35-36-61, 1999/21-47-53-68 and 2000/053-056-061

Illegal trade: In 1998, confiscated caviar shipments of unknown origin totalling 87 kg have been reported by USA upon import. This was composed of 18 kg re-exported by Turkey and 69 kg re-exported by the United Arab Emirates. Discrepancies between trade statistics of countries importing and exporting caviar can indicate illegal activities. Such discrepancies were found in 1995 and 1996 for instance for Turkey (customs declared caviar exports 21 t and 80 t respectively) and European Union Member States (6 t and 20 t respectively) (Raymakers, 1999). Additionally, Turkish 1996 caviar exports totalled 121 t (Raymakers, 1999), while the country's sturgeon fisheries has been limited to only few incidental catches since the late 1980s (Edwards and Doroshov, 1989).

Poaching and smuggling are closely related and have been intensively reported in the media of both range States and importing countries (Evtouchenko, 1997; McDonald, 2000; Snyder, 2000). However, much of the illegally caught sturgeon is destined to the domestic market, particularly meat (Anon., 1998).

CONSERVATION MEASURES

H. huso is not fully protected in any range State, with the exception of Moldova. However, a license is required for fishing in most of the range States and in Iran private sturgeon fisheries are prohibited. In 1996, an agreement between countries bordering the Caspian Sea prohibited fishing in the open sea. Fish lifts and artificial spawning grounds have been introduced in some Caspian regions.

Azerbaijan: The Intergovernmental Commission for Caspian Biological Resources, composed of representatives of all range States, sets annual catch quotas (CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000).

Catch quotas for *H. huso* allocated to Azerbaijan (tonnes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>H. huso</i>	10	122	122								10.85
Total	200	308	510	480	360	182	160.5	160.5	160.5	108.67	108.35

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

The restocking programme initiated in the 1960s by the former Soviet Union has been particularly important for *H. huso*. Three hatcheries were built in the lower Kura River. Two million *H. huso* fry were released annually in the Kura River from 1967 to 1980. A feasibility study of the World Bank reports that the total capacity of the hatcheries was originally 12 million, but that this capacity had dropped significantly over the years (T. Turner, *in litt.* to TRAFFIC Europe, 6 October 2000). World Bank funds have been allocated for the construction of a new sturgeon hatchery and farm that will have the capacity to produce 15 million fry per annum. This should be operational in the next couple of years, late 2002.

However, the major concern for *H. huso* is the availability of adult fish to constitute the broodstock of the farm.

Number of sturgeon fry released into the lower Kura River (million fingerlings)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<i>H. huso</i>	0.406	0.144	0.450	0	0	0	0	0.072	0	0.162	0.357
Total	17.523	9.082	2.980	1.838	1.142	1.242	4.070	6.065	6.220	20.290	16.965

Source: CITES Management Authority of Azerbaijan, *in litt.* to TRAFFIC Europe, 18 September 2000

In Azerbaijan, the State institution responsible for sturgeon fishery and processing matters is the "State Concern Azerbalyg". Commercial sturgeon fisheries are prohibited in the open sea. No fishing licenses were required before 2000. The only fishing gear permitted for sturgeon catch in the Kura River is "sweep nets" of 35-55 mm mesh size. The minimum size limit set for *H. huso* catch is 165 cm in the Kura River, and 180 cm in coastal waters (Anon., 2000c).

Bulgaria: Measures at national level as described hereunder are needed, but they must be linked to co-ordinated actions of all countries in the Black Sea region. In Bulgaria, sturgeon fisheries are prohibited for 30 days, from 20 April to 20 May each year, at the peak of the breeding season. The authorities may approve a 15-day extension of the closed season pending the natural conditions of the year. Explosives, poison, bottom trawling and dragging, as well as stunning substances and devices are prohibited fishing methods. The minimum size limit for *H. huso* is 140 cm. Export quotas are set on the basis of catch data, trends and the biological characteristics of the species. A restocking programme exists and is controlled by the relevant government agencies. A system has been established for the distribution of export quotas among export companies that are obliged to restock the Danube River with sturgeon fry before they are authorised to export sturgeon specimens (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000). For example, for a given year, for each kilogramme of caviar exported, the company must release 30 juvenile sturgeon (Anon., 2000c).

Number of *H. huso* released through the restocking programme (number of fish)

	1998	Size	1999	Size	2000	Size
<i>H. huso</i>	200	20 g each	2,700	60 g each	750	550 g each

Source: CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000

Due to budgetary constraints, Bulgaria has not recently monitored the wild sturgeon population, nor the success of its restocking programme (e.g. through tagging of released fingerlings). However, plans have been prepared for these activities. A special prohibition was adopted on the use of "unbaited hooks" for sturgeon catch (Anon., 2000c).

Hungary: The species is protected under *Decree No. 1/1982 (III.15.) OKTH on the Protected and Strictly Protected Species of Flora and Fauna, Value of their Specimens, Determination of the Range of Protected and Strictly Protected Caves and Exemptions from Restrictions and Prohibitions Set for Certain Protected Animal Species*, last amended by the *15/1996 (VII.26.) Decree of the Minister for Environment*. The *Nature Conservation Act No. 53 of 1996, Article 43* orders that for all protected species, prior authorisation from the National Park Directorate is required for the: capture, killing, possession, captive breeding, introduction of specimens from exotic population, artificial exchange of genetic matter between populations, exchange, sale and purchase of *H. huso* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Biometric measurements are carried out for each species at all sturgeon catch stations. The age, length, sex and weight of fish landed are recorded. The population structure is estimated on the basis of age groups. As a result of these studies carried out for the past 10 years, the number of catch stations were decreased by 50%, changes were adopted on sturgeon catch methods and lower export quotas were set for the past five years. Export quotas are established on the basis of results of research programmes and annual monitoring of stocks. These activities include the following parameters and activities: Maximum Sustainable Yield and total catch, rehabilitation of stocks, management and sustainable exploitation through restricting illegal catch and overfishing. Monitoring fishing efforts and regulating the time of catch are also carried out

on the basis of research results (CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000).

Annual release of sturgeon fry in the 1990s (million fingerlings)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
H. huso	0.14	0.17	0.45	0.30	0.49	0.29	0.34	1.44	0.69	0.41
Total	4.56	6.60	3.45	4.17	5.91	9.13	12.35	21.63	24.56	19.10

Source: CITES Management Authority of Iran, *in litt.* to TRAFFIC Europe, 24 September 2000

A labelling system is in place, where each tin of caviar holds a sign or mark that indicates at least the date and place of catch, the name of the catch station, name of processor and the species.

Considering the biological and commercial importance of this species, catch of *H. huso* has been banned in some of the provinces for more than five years. Additional conservation measures have been taken for all sturgeon species in the southern shores of the Caspian Sea; banning the use of gillnets in the Caspian Sea has been effective in conserving the stocks of *H. huso* (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: Commercial sturgeon catch is only allowed in the tributaries of the Caspian Sea. However, scientific catch is permitted in the open sea with an annual quota of 20 t (Anon., 2000c).

Sturgeon fisheries are regulated by law. There are two fishing seasons, one during spring spawning and the other during winter migration. The exact dates of the fishing seasons vary with the climate and are therefore set annually by the authorised scientific research organisations. As in most former Soviet Republics, the only legal fishing gear for sturgeon are "sweep nets" (bottom nets that are hauled several times a day) with a minimum mesh size and each being marked with the name and address of the owner. Each fisherman can only set one net at the time. A fishing license is required. Sturgeon fishing grounds ("tonia") are limited: there are twelve on the Ural River and eight on the Kigach River. The minimum size limit for *H. huso* is 180 cm. The ratio of males and females caught is not regulated. The purchase price at the landing site is based on a fixed legal rate of caviar weight extracted per female; *H. huso* is set at 19.4% of the total body weight.

Annual commercial catch quotas are allocated to Kazakhstan by the Intergovernmental Commission for Caspian Biological Resources which meets annually in Astrakhan.

Annual catch quotas for Kazakhstan (tonnes)

	1996	1997	1998	1999	2000
H. huso			80	60	58.5
Total	375	480	460	405	371

Source: CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, 21 Sept. 2000

Two hatcheries, Uralo-Atyrau and Atyrau, have been operational since 1998.

Annual release of *H. huso* fry (number of fingerlings)

	1998	1999	2000
H. huso		515,666	192,000
Total	1,706,193	5,318,471	5,507,160

Source: CITES Management Authority, *in litt.* to TRAFFIC Europe, 21 September 2000

Moldova: *H. huso* fishery is prohibited by law (Anon., 1999).

Romania: The type of sturgeon fishing gear is strictly regulated by Romanian law. The fishing season is set each year by Ministerial Order. A fishing license is required to catch sturgeon in Romanian waters. The minimum size limit for *H. huso* is 170 cm. Catch quotas are calculated by independent scientific institutes, such as the Danube Delta Institute in Tulcea and the "Centre for Conservation, Fish Farming, Fishing and Fish Industrialisation" in Galati. The 1998 and 1999 catch quotas for *H. huso* were set at 17.5 t/year. The Romanian restocking programme has not released juvenile Beluga in the Danube River (Anon., 2000c).

Russian Federation: For Caspian Sea populations, the former Soviet Republics set annual catch quotas in agreement with Iran. Sturgeon catch quotas have been established by the Intergovernmental Commission for Caspian Biological Resources for many years. Since 1993, the annual share of sturgeon catches of each former USSR Caspian range State are as follows: the Russian Federation 70%, Kazakhstan 17.6%, Azerbaijan 6.1% and Turkmenistan 6.3%. There are two fishing seasons in the Caspian Sea waters: from ice melting to 5 June, and from 1 September until the time of ice formation. A license is required to catch sturgeon in Russian waters. The minimum size limit for *H. huso* caught in Caspian Russian waters is 180 cm along the coast, and 165 cm in the northern Caspian Sea. Russian catch quotas are declared by the Federal Government following recommendations given by the State Fisheries Committee in consensus with an independent council of experts (Anon., 2000b).

Release of *H. huso* in the Volga River and the Sea of Azov by Russian hatcheries (number of fish)

	1979-1980	1981-1985	1986-1990	1991-1995	1996-1998
Volga River	15,000,000	18,300,000	13,000,000	8,500,000	12,000,000
Sea of Azov	2,000,000	222,800	377,000	27,200	404,218

Source: Azov and Caspian Fisheries Research Institute (Az- and KaspNIRKh), cited in Anon., 2000b

From 1979 to 1998, *H. huso* comprised approximately 21.2% and 4.2% of all fry released by Russian hatcheries into the Volga River and the Sea of Azov respectively (Anon., 2000b).

Turkey: In 1979, with the exception of *H. huso* of more than 140 cm long, the fishery for sturgeon was prohibited in Turkish waters (Edwards and Doroshov, 1989). Circular 34/1, effective from 2000-2002, prohibits the collection of Acipenseriformes (CITES Management Authority of Turkey, *in litt.* to TRAFFIC Europe, 20 September 2000).

Turkmenistan: In the country's coastal waters, sturgeon fishery is allowed for scientific research only, with an annual catch limit of 20 t. The Turkmenistan State Committee of Fisheries issues an annual license for sturgeon commercial catches to the State enterprise, Balkanbalyk. No other company, Turkmen or otherwise is permitted to commercially fish sturgeon in Turkmen inland waters. An annual catch quota for the species in Turkmenistan is set by the Intergovernmental Commission for Caspian Biological Resources. There is no sex ratio set for sturgeon landings. Information on the minimum size limit per species and data on the annual catch quotas set for Turkmenistan were not provided (Anon., 2000c).

Ukraine: Catch of *H. huso* in Ukrainian waters is prohibited since the species was listed in the Red Data Book of Ukraine in 1992 (Anon., 2000c).

Yugoslavia: The closed fishing season set for *H. huso*, *A. gueldenstaedtii* and *A. stellatus* is from 15 May to 15 June. An agreement was signed between Romania and Yugoslavia on the prohibition of fishing gear causing high mortality of untargeted fish. Any fishing activity, including all commercial and sport fisheries, requires a license in Yugoslav waters. The licenses are issued on a per district basis and the sturgeon fishing grounds are limited to 17.8 km downstream of the Djerdap II dam and the mouth of the Timok River. The minimum size limit for *H. huso* is 140 cm and the annual catch quota for the species from 1996 to 2000 was set at 4,500 kg per year. A restocking programme is being carried out by the Centre for fish reproduction of the "Djerdap Fishing Company" in Malla Vrbica village. Data on the release of juvenile sturgeon have not been provided (Anon., 2000c)

CAPTIVE BREEDING

The reduction in the area of natural spawning grounds due to river flow regulation has caused population declines. Fish reared in hatcheries have therefore been produced to maintain the stock. In the former Soviet Union *H. huso* has been bred in captivity since the 1960s in order to maintain the spawning stocks for commercial harvests of the Sea of Azov, the Caspian and Black seas. An average of 0.8 million fry were released annually into the southern Caspian Sea at the end of the 1980s. Most hatcheries of Azerbaijan, Kazakhstan and the Russian Federation, ceased operation in the early to mid-1990s following shortage of government funds (De Meulenaer and Raymakers, 1996).

Azerbaijan: Three hatcheries were built in the 1960s and have contributed to the former Soviet Union sturgeon restocking programme ever since. Their combined annual productive capacity was 12 million fry, but this has dropped significantly due to rising levels of the Caspian Sea and the lack of funds allocated to hatchery maintenance (T. Turner, World Bank consultant, *in litt.* to TRAFFIC Europe, 6 October 2000).

Bulgaria: A fish farm was established in 1997 in Boliartzi village (Plovdiv region). It is the only existing hatchery. All juveniles released in the Danube River in Bulgaria from 1998 to 2000 were produced there. The hatchery has a total capacity of 60,000 juveniles and the farm is currently stocked with 3.5 year old *H. huso* and *A. gueldenstaedtii* that weigh 11-14 kg each. Since 1999, attempts have been made to breed sturgeon in artificial water basins near Sofia (CITES Management Authority of Bulgaria, *in litt.* to TRAFFIC Europe, 15 September 2000).

Georgia: Until the early 1990s, artificial breeding of sturgeon species was carried out at the Varistkhi Sturgeon Plant. Built in 1983 and located on the Rioni River, the results were not always considered satisfactory (Zarkua and Tsuladze, 1999).

Hungary: There are no captive breeding facilities for *H. huso* (CITES Management Authority of Hungary, *in litt.* to TRAFFIC Europe, 13 September 2000).

Iran: Five hatcheries are in operation in Iran (De Meulenaer and Raymakers, 1996). In the mid-1990s, more than five million *H. huso* fingerlings weighing 3-5 g were released into the Caspian Sea by Iran. In 2000, more than 1.5 million *H. huso* weighing 30-50 g were released into the Caspian Sea (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000). In Iran, this species has been reared in captivity for more than ten years. Apart from the governmental sector, the non-governmental sector has also invested in the captive breeding of this species. At present, more than 30,000 *H. huso* individuals between one and ten years of age have been introduced into rearing ponds. Iran intends to culture 1,000 t of *H. huso* for meat and caviar production (Dr M. Pourkazemi, *in litt.* to IUCN/SSC Wildlife Trade Programme, September 2000).

Kazakhstan: There are two hatcheries in Kazakhstan, the Uralo-Atyrau and the Atyrau. These have been operational since 1998 (CITES Management Authority of Kazakhstan, *in litt.* to TRAFFIC Europe, September 2000).

Russian Federation: In the mid-1990s, it was estimated that approximately 91% of the Caspian Sea stock of *H. huso* originated from hatcheries (Barannikova, 1995), whilst in 1993, 100% of each generation of *H. huso* in the Sea of Azov derived from captive-bred released fry (Volovik *et al.*, 1993). However, high levels of poaching have reduced natural populations to such low levels that most hatcheries have not been able to catch enough mature fish to produce fry for the restocking programme (Artyukhin, 1997). A hatchery on the Kuban River (Russian Federation) was still operating in 1997 (Chebanov and Savelyeva, 1999). In European countries, a hybrid of *H. huso* and *A. ruthenus* called 'bester' is subject to aquaculture. It is fast growing and produces fertile and early maturing offspring (Burtsev, 1997).

United States of America: Sturgeon species are in preliminary stage of captive breeding in Hawaii (CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000).

REFERENCES

- Anon. (1996). *A Guide to Interpreting Outputs from the WCMC CITES Trade Database*, Produced by the World Conservation Monitoring Centre, Cambridge UK. Pp. 16.
- Anon. (1997). Document Doc. 10.89; Prop. 10.65. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Anon. (1998). *Sturgeon catch and trade in the Russian part of the Caspian Sea*. TRAFFIC Europe-Russia field investigations. Unpublished report, Pp. 21.
- Anon. (1999). The Law of the Republic of Moldova on Fund of Natural Areas Protected by State (1998). In: "*Ekologicheskoe Zakonodatel'stvo Respubliki Moldova (1996-1998)*". Chisinau, BIOTICA Publications, 1999. Pp. 85-209.
- Anon. (2000a). <http://www.cites.org/eng/dbase/fauna->

- Anon. (2000b). *Estimation of the Sturgeon Stocks in the Russian Federation and Monitoring of Domestic Trade in Sturgeon Products*. TRAFFIC Europe-Russia field investigations. Unpublished report, Pp. 23.
- Anon. (2000c). *Sturgeon fisheries management and trade control measures in the Caspian Sea and Black Sea/ Sea of Azov range States*. TRAFFIC Europe field investigations, December 1999-January 2000. Unpublished.
- Artyukin, E.N. (1997). The current status of commercial sturgeon species in the Volga River-Caspian Sea Basin. In: Birstein, V.J., Bauer, A. Kaiser-Pohlmann, A. (eds.) *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 9-18.
- Bacalbasa-Dobrovici, N. (1997). Endangered migratory sturgeons of the lower Danube River and its delta. *Environmental Biology of Fishes*, 48:201-207.
- Barannikova, I.A., Burtsev, I.A., Vlasenko, A.D. Gershanovich, A.D., Markarov, E.V. and Chebanov, M.S. (1995). Sturgeon Fisheries in Russia. Proceedings of the Second International Symposium on Sturgeons, September 6-11, 1993. Moscow-Kostroma-Moscow. VNIRO Publication. Pp. 124-130.
- Berg, L.S. (1948). [O. polozhenii Acipenseriformes v sisteme ryb. Trudy Zoological Institute, 7: 7-57.] (in Russian)
- Burtsev, I.A. (1997). Bester in aquaculture. In: Birstein, V.J., Bauer, A. and Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 35-43.
- Chebanov, M.S. and E.A. Savalyeva. (1999). New strategies for broodstock management of sturgeon in the Sea of Azov basin in response to changes in patterns of spawning migration. *J. Appl. Ichthyol.* 15:183-190.
- Chepurnova L.V. (1991). Zakonomernosti Funktsii Gonad, Razmnozhenia I Sostoiania Populiatsii Ryb Basseina Dnestra v Usloviah Hidrostroytel'stva. Kishinev, Shtiintsa Publ. House, Pp. 164.
- De Meulenaer, T. and Raymakers, C. (1996). *Sturgeons of the Caspian Sea and the international trade in caviar*. TRAFFIC International, iv+ 71 Pp.
- Dobbs, M. (1992). The Coming Caviar Crisis: Overfishing, Pollution Endanger Sturgeon. *The Washington Post*, 30 May 1992.
- Doward, J. (2000). Roe rage as piracy rules. *The Guardian*, 21 May 2000.
- Dumont, H. (1995). Ecocide in the Caspian Sea. *Nature* 377pp. 673-674.
- Edwards, D. and Doroshov, S. (1989). Appraisal of the Sturgeon and Seatrout Fisheries and Proposals for a Rehabilitation Programme. FAO Field Document FI.TCP/TUR/8853. Pp. 38.
- Evtouchenko, A. (1997). Au Dagestan une explosion fait 54 morts. Accusés: les gangs du caviar. *Courrier International*, N° 323 (9-15 January 1997)[in French].
- Glowacki, J., (1896). Die Fischfauna der Save und der Isonzo. I.- Jahresbericht d. k.k. Staats-Untergymnasium in Cilli, 1895-1896:1-37.
- Hensel, K. and Holcik, J. (1997). Past and current status of sturgeon in the upper and middle Danube. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 185-200.
- Hochleithner, M. and Gessner, J. (1999). *The Sturgeon and Paddlefishes (Acipenseriformes) of the World: Biology and Aquaculture*. AquaTech Publications, Kitzbühl. Pp. 165.
- Holcik, J., 1995: 1.druh Huso huso (Linnaeus, 1758). p.374-378. In: V.Baruš and O.Oliva (eds.) *Mihulovci Petromyzontesa ryby Osteichthyes. 1. Fauna ĚR a SR 28/1*. Academia, Praha.
- IUCN, (1996). *IUCN Red List of Threatened Animals*. IUCN, Gland, Switzerland
- Jankoviæ, D. (1999). The Ecological and Geographical Fish Distribution in the Yugoslav Section of the Danube. Contribution to the Zoogeography and Ecoogy of the Eastern Mediterranean Region. Vol. I (1999), Pp. 321-326.
- Jankoviæ, D. (1993). Populations of Acipenseridae prior and after the construction of the HEPS Djerdap I and II. *Acta Biologica iugoslavica, Seria E. Ichthyologia* 25:29-34
- Jankoviæ, D. (1996). Ichthyofauna of the Danube in the Djerdap area after the construction of the Iron Gate I Hydroelectric Power Station. *Acta Universitatis Carolinae, Biologica*, 40:123-131
- Karapetkova, M., M. Zhivkov, and T. Pchelarov, 1995: Ribite v B'lgariya. Geya Libris, Sofia. 247 pp.
- Khodorevskaya, R.P., Dovgopol, G.F., Zhuraleva, O.L., and Vlasenko, A.D. (1997). Present status of commercial stocks of sturgeons in the Caspian Sea Basin. In: Birstein, V.J., Waldman, J.R. and Bemis, W.E. (eds) (1997). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publications, Dordrecht. Pp. 209-219.
- Khodorevskaya, R.P. and Novikova, A.S. (1995). Status of Beluga Sturgeon, *Huso Huso*, in the Caspian Sea. *J. Ichthyol.* 35 (9):59-68.

- King, R. (1998). Por un punado de caviar. *El Mundo*, 13 December 1998. [In Spanish].
- Levin, A.V. (1997). The Distribution and Migration of Sturgeon in the Caspian Sea. In: Birstein, V.J., Bauer, A. Kaiser-Pohlmann, A. (eds.) (1997). *Sturgeon Stocks and Caviar Trade Workshop*. IUCN, Gland, Switzerland and Cambridge, UK. Pp. 13-19.
- Ljiljana, B., B.Cececic, N.Sekulic and D.Brancovic, (1998). Protection of the Yugoslav section of the Danube downstream from "HE Djerdap II". p.70-71. In M.Rauta, N.Bacalbasa-Dobrovici, G.Vasilescu, and L.Oprea (eds.) *Fisheries Management in the Danube River Basin*. Extended abstracts of contributions presented at the international symposium AQUAROM '98, Galati-Romania 18-22 May 1998. Universitatea "Dunarea de Jos", Galati, Asociatia Piscicultilor si Pescailor din Romania.
- McDonald, L. (2000). Fish that laid the golden egg faces extinction. *Daily Express* (UK), 26 July 2000.
- Medetski, A. (2000). Black caviar feeds thousands on Amur. *Vladivostok News*, 15 September 2000.
- Munda, A. (1926). *Ribe v Slovenskih vodah*. Izd. Slovensko Ribarsko Društvo v Ljubljani, Ljubljana. Pp. 63
- Pavlov, P.J., (1980): *Ribi*. Fauna Ukrainy. 8/1. Naukova Dumka, Kiiiv. 336 pp.
- Pintér, K. (1991). Sturgeon in Hungary, past and present situation. In: P. Williot (Ed.) *Acipenser*. Actes du premier colloque international sur l'esturgeon. Bordeaux, 3-6 October 1989. pp. 173-178.
- Pirogovskii, M.I., Sokolov, L.I. and Vasil'ev, V.P. (1989). *Huso huso* (Linnaeus, 1758). In: Holcík J. (ed). *The Freshwater Fishes of Europe*. Vol. I/II: General Introduction of Fishes. Acipenseriformes. Wiesbaden, AULA-Verlag. Pp. 295-344. In: Document Doc. 10.89; Prop. 10.65. 1997. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Povž, M. and Sket, B. (1990). *Naše sladkovodne ribe*. Založba Mladinska Kniga, Ljubljana. 374 pp.
- Razavi, S.B. (1988). Analysis on great sturgeon stocks in the South Caspian Sea. Iranian Fisheries Research and Training Organisation Publications.
- Raymakers, C. (1999). Trade in Sturgeons from the Caspian Sea. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 149-161.
- Rossi, R., Grandi, G., Trisolini, P., Franzoi, P., Carrieri, a., Dezfuli, B. S. and Vecchietti, E. (1991). Osservazioni sulla biologica e la pesca dello storione cobice (*Acipenser naccarii*, Bonaparte) nelle parte terminale de fiume Po. *Atti Soc.Ital.Sci.Natur.Mus.Civ.Stor. Natu. Milano*, 132:121-142.
- Sagers, M.J. (1994). The oil industry in the southern-tier former soviet republics. *Post-Soviet Geography*, 35(5):267-298.
- Shagaeva, V.G., Nikol'skaya, M.P., Akimova, K.P., Markov, N.G. and Nikol'skaya, N.G. (1993). [A study of the early ontogeny of Volga Sturgeon, (Acipenseridae) subjected to human activity. *J. Ichthyol.* 33(6):23-41.] (in Russian)
- Suciu, R., Ene, F. and Bacalbasa Dobrovici, N. (1998). New data on the distribution of you-of-the-year sturgeon in the Lower Danube River. Eds. Raute, M., Bacalbasa Dobrovici, N. and Oprea, L. In: *Aquarum 1998*, Galati, Romania. University "Dunarea de Jos".
- Simonovic, P. D. and Nikolic, V.P. (1996). Freshwater fish of Serbia: an annotated check list with some faunistic and zoogeographical considerations. *Bios* (Macedonia, Greece) 4:137-156.
- Snyder, D. (2000). Md. Caviar Company Admits Smuggling. *The Washington Post*, 22 July 2000.
- Taler, Z. (1954). Rasprostranjenje i popis slatkovodnih riba Jugoslavije. *Glasnik Prirodnaèog Muzeja Srpske Zemlje*, ser.B 5/6:425-455.
- Vedrashko A., Lobchenko V., Pirtsu, I. (1998). [The Problem of Conservation of the Acipenserids in the Lower Dniester. In: Problemele Conservarii Biodiversitatii din Cursul Medial si Inferior al Fluviului Nistru. Abstr. Int. Conf. Chisinau, 6-7 Nov. 1998. Chisinau, BIOTICA Publ.,1998. Pp. 35-36.] (In Russian)
- Vlasenko, A.D. (1990). Sturgeon Population Size in the Caspian Sea. *Rybnoe Khozyaistvo*, 7:53-56.
- Vlasenko, A.D., Pavlov, A.V., Sokolov, L.I. and Vasil'eva, V.P. (1989). *Acipenser gueldenstaedti* Brandt, 1833. In: Holcík J. (ed). *The Freshwater Fishes of Europe*. Vol. I/II: General Introduction of Fishes. Acipenseriformes. Wiesbaden, AULA-Verlag. Pp. 295-344. In: Document Doc. 10.89; Prop. 10.65. 1997. Proposal to list all Acipenseriformes in Appendix II. Submitted by Germany and The United States of America.
- Volovik, S.P., Dubinina, V.A., and Semenova, A.D. (1993). Hydrobiology and dynamics of fisheries in the Asov Sea. Studies and reviews. *General Fisheries Council for the Mediterranean*. No. 64. FAO, Rome. Pp. 1-58.
- Ward, P. (translation) (2000a). Azov Sturgeons Can't be Touched. *Crimean Time*, 18 August 2000.

- Ward, P. (translation) (2000b). Makhachkala: the Scale of Poaching in the Caspian has Reached Enormous Proportions. *Regions.Ru/Russia Regions*, 15 August 2000.
- Williot, P. and Bourguignon, G. (1991). Sturgeon and caviar production, current status and perspectives. In: Williot, P. (ed.), *Acipenser*, Cemagref Publications, Pp. 509-513.
- Zarkua, Z. and Tsuladze, V. (1999). Initiative needed to boost sturgeon numbers. *Eurofish* 3/99:40-41.

APPENDIX

Gross exports of, and export quotas for *Huso huso* 1998

TAXON	TERM	UNI T	Export er	1998	Export Quotas 1998
<i>Huso huso</i>	bodies		PG	8	
<i>Huso huso</i>	bodies #	kg	IR	12,750	
<i>Huso huso</i>	eggs		AE	267	
<i>Huso huso</i>	eggs		DE	88	
<i>Huso huso</i>	eggs		FR	350	
<i>Huso huso</i>	eggs		NL	16	
<i>Huso huso</i>	eggs		RO	90	
<i>Huso huso</i>	eggs		SG	1	
<i>Huso huso</i>	eggs		TR	69	
<i>Huso huso</i>	eggs		US	2	
<i>Huso huso</i>	eggs	kg	AE	3,001	
<i>Huso huso</i>	eggs	kg	AZ	114	
<i>Huso huso</i>	eggs	kg	BE	28	
<i>Huso huso</i>	eggs	kg	BG	1,717	2,500
<i>Huso huso</i>	eggs	kg	CH	6,467	
<i>Huso huso</i>	eggs	kg	CZ	8	
<i>Huso huso</i>	eggs	kg	DE	1,281	
<i>Huso huso</i>	eggs	kg	DK	226	
<i>Huso huso</i>	eggs	kg	ES	0	
<i>Huso huso</i>	eggs	kg	FI	183	
<i>Huso huso</i>	eggs	kg	FR	789	
<i>Huso huso</i>	eggs	kg	GB	18	
<i>Huso huso</i>	eggs	kg	HK	32	
<i>Huso huso</i>	eggs	kg	IR	4,013	5,000
<i>Huso huso</i>	eggs	kg	KZ	7,033	
<i>Huso huso</i>	eggs	kg	NL	1,418	
<i>Huso huso</i>	eggs	kg	PL	4,365	
<i>Huso huso</i>	eggs	kg	RO	303	
<i>Huso huso</i>	eggs	kg	RU	8,035	5,000 (including AZ & KZ)
<i>Huso huso</i>	eggs	kg	SG	8	
<i>Huso huso</i>	eggs	kg	TR	1,075	
<i>Huso huso</i>	eggs	kg	UA	52	
<i>Huso huso</i>	eggs	kg	US	150	
<i>Huso huso</i>	meat	kg	IR	3,005	
<i>Huso huso</i>	meat	kg	US	18	
<i>Huso huso</i>	skins		CH	22	
<i>Huso huso</i>	skins		IR	22	
<i>Huso huso</i>	specimens		DE	0	

The term "bodies" should be interpreted as "meat" (Dr P. Bronzi, *in litt.* to IUCN/SSC Wildlife Trade Programme, 8 September 2000).

* **Definition:** "Gross exports are the sum of all reported exports and re-exports in a particular commodity or species in a particular year or series of years;... Gross trade is thus a simple measure of the total number of items recorded in international trade. However, gross trade may be an overestimate of the total number of actual specimens in trade as re-exports are not deducted from the total." (Anon., 1996).

This is particularly applicable to caviar of which shipments are often exported, re-packaged and re-exported. For instance, in the mid-1990s, Germany re-exported as much as 45% of its annual caviar imports (De Meulenaer and Raymakers, 1996).

Comparative Tabulation of trade in *Huso huso* 1998

Year	Imp.	Exp.	Origin	Imports reported		Term	P	S	Exports reported		P	S
				Quantity	Unit				Quantity	Unit		
1998	AU	AE	RU						5 kg	eggs	T	O
1998	CA	AE	IR						4 kg	eggs	T	W
1998	CH	AE	IR	26 kg	eggs	T	W		8 kg	eggs	T	W
1998	CH	AE	IR	10 kg	eggs	T	O		10 kg	eggs	T	O
1998	CH	AE	RU						58 kg	eggs	T	W
1998	CH	AE	RU	16 kg	eggs	T	O					
1998	DK	AE	IR	4 kg	eggs	T	W		6 kg	eggs	T	W
1998	FR	AE	IR	18 kg	eggs	T	W					
1998	FR	AE	RU	20 kg	eggs	T	O					
1998	GB	AE	RU						25 kg	eggs	P	W
1998	GB	AE	RU						4 kg	eggs	T	W
1998	HK	AE	IR						54 kg	eggs	T	W
1998	HK	AE	IR						10 kg	eggs	T	O
1998	HK	AE	RU						15 kg	eggs	T	W
1998	JP	AE	RU						223 kg	eggs	T	O
1998	KR	AE	RU						4 kg	eggs	T	O
1998	NO	AE	IR	600 kg	eggs	T	O		600 kg	eggs	T	O
1998	NO	AE	RU						122 kg	eggs	T	O
1998	TW	AE	IR						10 kg	eggs	T	W
1998	US	AE		69 kg	eggs	T	I					
1998	US	AE	RU	19,625 g	eggs	T	I					
1998	US	AE	RU	651 kg	eggs	T	O		1,630 kg	eggs	T	O
1998	US	AE	RU	361 kg	eggs	T	W		155 kg	eggs	T	W
1998	US	AE	RU	190	eggs	T	O					
1998	US	AE	RU	77	eggs	T	W					
1998	US	AE	RU	30 kg	eggs	T	I					
1998	US	AE	XX	127 kg	eggs	T	O					
1998	UZ	AE	RU						20 kg	eggs	T	O
1998	CA	BE	RU						18 kg	eggs	T	O
1998	CA	BE	TR						9 kg	eggs	T	O
1998	TR	BE	RU						360 g	eggs	T	O
1998	CH	BG		1,056 kg	eggs	T	W					
1998	CH	BG		211 kg	eggs	T	O					
1998	FR	BG		450 kg	eggs	T	W					
1998	AT	CH	AZ	1 kg	eggs	T	O					
1998	AT	CH	IR						61 kg	eggs		O
1998	AT	CH	IR	5 kg	eggs	T	O					
1998	AT	CH	RU						5 kg	eggs		O
1998	AT	CH	RU	4 kg	eggs	T	O					
1998	BE	CH	KZ	87 kg	eggs	T	W		87 kg	eggs		W
1998	CH	CH	IR						28 kg	eggs		O
1998	CH	CH	RU						300 kg	eggs		O
1998	CY	CH	IR						1 kg	eggs		O
1998	DE	CH	AZ	212 kg	eggs	T	O					
1998	DE	CH	IR	61 kg	eggs	T	W					
1998	DE	CH	IR						60 kg	eggs		W
1998	DE	CH	IR						2 kg	eggs		O
1998	DE	CH	KZ	2,927 kg	eggs	T	W					
1998	DE	CH	RU						2,926 kg	eggs		W
1998	DE	CH	RU						158 kg	eggs		O
1998	DK	CH	RU						50 kg	eggs		O
1998	FR	CH	IR	291 kg	eggs	T	O					
1998	FR	CH	IR	103 kg	eggs	T	W					

Year	Imp.	Exp.	Origin	Imports reported		Term	P	S	Exports reported		Term	P	S
				Quantity	Unit				Quantity	Unit			
1998	FR	CH	IR						102 kg	eggs			W
1998	FR	CH	IR						1 kg	eggs			O
1998	FR	CH	KZ	1,253 kg	eggs		T	W					
1998	FR	CH	RU						1,251 kg	eggs			W
1998	GB	CH	AZ	47 kg	eggs		T	W					
1998	GB	CH	KZ	19 kg	eggs		T	W					
1998	GB	CH	RU						412 kg	eggs			W
1998	GB	CH	RU						21 kg	eggs			O
1998	HK	CH	IR						12 kg	eggs			O
1998	HK	CH	IR						5 kg	eggs			W
1998	HK	CH	RU						38 kg	eggs			O
1998	HK	CH	RU						21 kg	eggs			W
1998	IT	CH	IR						22	skins			W
1998	JP	CH	RU						12 kg	eggs			O
1998	KW	CH	IR						10 kg	eggs			O
1998	MU	CH	IR						0 kg	eggs			O
1998	MX	CH	IR						9 kg	eggs			O
1998	PA	CH	IR						1 kg	eggs			O
1998	SG	CH	IR	13 kg	eggs		T	O					
1998	SG	CH	IR						13 kg	eggs			O
1998	SG	CH	IR						5 kg	eggs			W
1998	SG	CH	IR	5 kg	eggs		T	W					
1998	US	CH	BG	432 kg	eggs		T	I					
1998	US	CH	BG						240 kg	eggs			W
1998	US	CH	BG						192 kg	eggs			O
1998	US	CH	RU						4 kg	eggs			O
1998	US	CH	YU	94 kg	eggs		T	I					
1998	ZA	CH	IR						2 kg	eggs			O
1998	CA	CZ	AZ						8 kg	eggs		T	W
1998	AD	DE	RU						6 kg	eggs		T	W
1998	CA	DE	RU						336 g	eggs		T	O
1998	CA	DE	RU						9 kg	eggs		T	O
1998	CH	DE	AZ						106 kg	eggs		T	O
1998	CH	DE	IR	29 kg	eggs		T	O					
1998	CH	DE	IR	27 kg	eggs		T	W	27 kg	eggs		T	W
1998	CH	DE	KZ						27 kg	eggs		T	W
1998	CH	DE	RU	177 kg	eggs		T	O					
1998	CH	DE	RU	29 kg	eggs		T	W	3 kg	eggs		T	W
1998	CZ	DE	IR						300 g	eggs		T	O
1998	CZ	DE	RU						5 kg	eggs		T	O
1998	CZ	DE	RU	5 kg	eggs		T	W					
1998	HK	DE	IR						2 kg	eggs		T	O
1998	HK	DE	KZ						12 kg	eggs		T	O
1998	HK	DE	RU						11 kg	eggs		T	O
1998	HR	DE	IR						600 g	eggs		T	O
1998	HR	DE	RU						2 kg	eggs		T	O
1998	JP	DE	KZ						13 kg	eggs		T	W
1998	JP	DE	KZ						4 kg	eggs		T	O
1998	JP	DE	RU						34 kg	eggs		T	O
1998	KR	DE	KZ						7 kg	eggs		T	W
1998	KR	DE	KZ						3 kg	eggs		T	O
1998	KR	DE	RU						15 kg	eggs		T	O
1998	KR	DE	RU	1 kg	eggs		T	W					
1998	MT	DE	RU						336 g	eggs		T	O

Year	Imp.	Exp.	Origin	Imports reported		Term	P	S	Exports reported		Term	P	S
				Quantity	Unit				Quantity	Unit			
1998	NO	DE	KZ						2 kg	eggs		T	W
1998	NO	DE	RU						6 kg	eggs		T	O
1998	SG	DE	KZ	10 kg	eggs		T	O	10 kg	eggs		T	O
1998	US	DE	KZ	218 kg	eggs		T	W	602 kg	eggs		T	W
1998	US	DE	KZ	214 kg	eggs		T	O	110 kg	eggs		T	O
1998	US	DE	RU	133 kg	eggs		T	O	137 kg	eggs		T	O
1998	US	DE	RU	89 kg	eggs		T	W	3 kg	eggs		T	W
1998	US	DE	RU	88	eggs		T	O					
1998	US	DE	RU	4 kg	eggs		T	F					
1998	ZA	DE	RU						23 kg	eggs		T	O
1998	BR	DK	RU						21 kg	eggs		T	O
1998	BR	DK	RU						10 kg	eggs		T	W
1998	CH	DK	RU	50 kg	eggs		T	O					
1998	SA	DK	IR						12 kg	eggs		T	W
1998	SA	DK	RU						30 kg	eggs		T	O
1998	TW	DK	IR						104 kg	eggs		T	W
1998	RU	FI	RU						183 kg	eggs		T	W
1998	AN	FR	IR						2 kg	eggs		T	W
1998	BR	FR	IR						1 kg	eggs		T	O
1998	BS	FR	IR						6 kg	eggs		T	W
1998	BS	FR	KZ						1 kg	eggs		T	O
1998	CA	FR	BG						19 kg	eggs		T	O
1998	CA	FR	IR						3 kg	eggs		T	O
1998	CA	FR	KZ						4 kg	eggs		T	O
1998	CA	FR	KZ						2 kg	eggs		T	W
1998	CH	FR	IR	21 kg	eggs		T	W	21 kg	eggs		T	W
1998	CI	FR	BG						4 kg	eggs		T	O
1998	CI	FR	IR						3 kg	eggs		T	W
1998	HK	FR	IR						9 kg	eggs		T	W
1998	HK	FR	IR						5 kg	eggs		T	O
1998	HK	FR	KZ						11 kg	eggs		T	O
1998	HK	FR	KZ						8 kg	eggs		T	W
1998	KR	FR	IR	0 kg	eggs		T	W	1 kg	eggs		T	W
1998	LB	FR	KZ						2 kg	eggs		T	W
1998	MX	FR	IR						13 kg	eggs		T	W
1998	MX	FR	IR						6 kg	eggs		T	O
1998	MX	FR	KZ						3 kg	eggs		T	O
1998	MX	FR	KZ						1 kg	eggs		T	W
1998	NC	FR	KZ						2 kg	eggs		T	W
1998	OM	FR	IR						28 kg	eggs		T	W
1998	OM	FR	KZ						42 kg	eggs		T	W
1998	OM	FR	KZ						13 kg	eggs		T	O
1998	PF	FR	KZ						1 kg	eggs		T	W
1998	PF	FR	KZ						1 kg	eggs		T	O
1998	PH	FR	IR						1 kg	eggs		T	W
1998	SA	FR	IR						3 kg	eggs		T	O
1998	SA	FR	KZ						8 kg	eggs		T	W
1998	SG	FR	IR	1 kg	eggs		T	W	2 kg	eggs		T	W
1998	SG	FR	KZ	2 kg	eggs		T	O	2 kg	eggs		T	O
1998	TH	FR	IR						1 kg	eggs		T	W
1998	TW	FR	IR						3 kg	eggs		T	W
1998	US	FR	KZ						554 kg	eggs		T	W
1998	US	FR	KZ	100	eggs		T	W					
1998	US	FR	RU	250	eggs			W					

Year	Imp.	Exp.	Origin	Imports reported		Term	P	S	Exports reported		Term	P	S
				Quantity	Unit				Quantity	Unit			
1998	US	FR	RU	239 kg	eggs	T	W						
1998	ZA	FR	IR						1 kg	eggs		T	W
1998	ZA	FR	KZ						1 kg	eggs		T	O
1998	CH	GB	RU	7 kg	eggs	T	O						
1998	EG	GB	IR						1 kg	eggs		T	W
1998	HK	GB	IR						2 kg	eggs		T	W
1998	IS	GB	IR						2 kg	eggs		T	W
1998	JO	GB	IR						1 kg	eggs		T	W
1998	JP	GB	IR						2 kg	eggs		T	W
1998	QA	GB	IR						1 kg	eggs		T	W
1998	SA	GB	IR						1 kg	eggs		T	W
1998	CH	HK	IR	1 kg	eggs	T	O						
1998	CH	HK	RU	31 kg	eggs	T	O						
1998	BE	IR		20 kg	eggs	T	W						
1998	BE	IR		5 kg	meat	T	W						
1998	CH	IR		180 kg	eggs	T	W						
1998	CH	IR	RU	319 kg	eggs	T	O						
1998	CH	IR	RU	310 kg	eggs	T	W						
1998	CH	IR		1,361 kg	meat	T	O						
1998	CH	IR		1,020 kg	meat	T	W						
1998	CH	IR		22	skins	T	W						
1998	DE	IR		127 kg	eggs	T	O						
1998	DE	IR		77 kg	eggs	T	W						
1998	ES	IR		710 kg	eggs	T	W						
1998	FR	IR		1,600 kg	eggs	T	W						
1998	FR	IR		160 kg	eggs	T	O						
1998	FR	IR		585 kg	meat	T	W						
1998	FR	IR		34 kg	meat	E	W						
1998	GB	IR		310 kg	eggs	T	W						
1998	IT	IR		12,750 kg	bodies	T	W						
1998	LU	IR		200 kg	eggs	T	W						
1998	BN	NL	IR						1 kg	eggs		T	O
1998	CH	NL	XX	1417 kg	eggs	T	O						
1998	ID	NL	IR						16	eggs		T	W
1998	ES	PG		8	bodies	T	R						
1998	GB	PL	RU						100 kg	eggs			W
1998	US	PL	RU	174.01 kg	eggs	T	O						
1998	US	PL	RU	77.5 kg	eggs	T	I						
1998	US	PL	RU						4,265 kg	eggs			W
1998	US	PL	RU	1,197 kg	eggs	T	O						
1998	US	PL	RU	800 kg	eggs	T	W						
1998	GR	RO		15 kg	eggs	T	W						
1998	US	RO		51,200 g	eggs	T	W						
1998	US	RO		237 kg	eggs	T	W						
1998	US	RO		90	eggs	T	O						
1998	AE	RU	KZ						648 kg	eggs		T	W
1998	AU	RU							2 kg	eggs		P	W
1998	AZ	RU							37 kg	eggs		T	W
1998	BE	RU		45 kg	eggs	T	W		45 kg	eggs		T	W
1998	CH	RU		4,812 kg	eggs	T	W		37 kg	eggs		T	W
1998	CH	RU		520 kg	eggs	T	O						
1998	CH	RU	KZ						4,812 kg	eggs		T	W
1998	DE	RU		137 kg	eggs	T	W		462 kg	eggs		T	W
1998	DE	RU		109 kg	eggs	T	O		211 kg	eggs		T	O

Year	Imp.	Exp.	Origin	Imports reported		Term	P	S	Exports reported		Term	P	S
				Quantity	Unit				Quantity	Unit			
1998	FI	RU		183 kg	eggs	T	W		183 kg	eggs	T	W	
1998	FR	RU	KZ	40 kg	eggs	T	W		40 kg	eggs	T	W	
1998	US	RU		106 kg	eggs	T	I						
1998	US	RU		753 kg	eggs	T	O						
1998	US	RU		94 kg	eggs	T	W		1,074 kg	eggs	T	W	
1998	CH	SG	IR	8 kg	eggs	T	O		8 kg	eggs	T	O	
1998	US	SG	RU	1	eggs	T	W						
1998	CH	TR	RU	96 kg	eggs	T	O		98 kg	eggs	T	O	
1998	US	TR							34 kg	eggs	T	O	
1998	US	TR	RU	416 kg	eggs	T	W						
1998	US	TR	RU	75 kg	eggs	T	O		621 kg	eggs	T	O	
1998	US	TR	RU	69	eggs	T	O						
1998	US	TR	RU	2 g	eggs	T	I						
1998	US	TR	RU	2 g	eggs	T	W						
1998	US	TR	RU	1 kg	eggs	T	I						
1998	US	TR	XX						322 kg	eggs	T	O	
1998	US	TR	XX	18 kg	eggs	T	I						
1998	US	UA	RU	52 kg	eggs	T	O						
1998	AR	US	RU						77 kg	eggs	T	O	
1998	AR	US	RU						2	eggs	T	O	
1998	AR	US	RU						1 kg	eggs	T	W	
1998	BM	US	RU						1 kg	eggs	T	W	
1998	CA	US	RU						35 kg	eggs	T	W	
1998	DE	US	RU	300 g	eggs	T	O		2500 g	eggs	T	W	
1998	GR	US	RU						300 g	eggs	T	O	
1998	JP	US	RU						1 kg	eggs	T	W	
1998	LC	US	RU						1 kg	eggs	T	W	
1998	MX	US	RU						7 kg	eggs	T	W	
1998	MX	US	RU						18 kg	meat	T	W	
1998	PT	US	RU						1 kg	eggs	T	W	
1998	SG	US	RU	3 kg	eggs	T	W		3 kg	eggs	T	W	
1998	TR	US	RU						18 kg	eggs	T	W	
1998	ZA	US	RU						2 kg	eggs	T	W	
1998	CH	XX		120 kg	other	T	W						
Total:				25,889.51 kg	eggs				24,511 kg	eggs			

Polyodon spathula Walbaum, 1792

**American Paddlefish
Poisson spatule**

Order: ACIPENSERIFORMES

Family: POLYODONTIDAE

SUMMARY

American Paddlefish *Polyodon spathula* is endemic to the continent and is the only paddlefish species present in the wild in North America. This species is widely distributed in the large river drainages of the eastern United States of America (USA). During the last century it has been extirpated from four states on the periphery of its range and even within states considered as strongholds for the species, portions of the historical range have diminished. The species reaches sexual maturity between 5-12 years of age and may live up to 40-50 years attaining a weight of up to 80 kg. Southern populations apparently grow more quickly than those in colder northern waters. Following a review of the status of the species in 1983, a number of significant changes in the management of the species were adopted, due to recognition of the threats posed by continued habitat degradation, pollution and the demand for paddlefish caviar.

Commercial catch of *P. spathula* is permitted in six states only: Arkansas, Illinois, Kentucky, Mississippi, Missouri and Tennessee. In the absence of past and present catch data from Mississippi, it is difficult to determine the trends in annual catches of *P. spathula*. A similar problem exists for Arkansas where reporting of paddlefish commercial catch is not compulsory. In 1992, the estimated commercial paddlefish fishery, excluding Mississippi, totalled approximately 242 t. The largest volumes were recorded in Arkansas (65%; 136,000-181,000 kg – 158,500 kg mean) and Tennessee (25%; 60,328 kg), with the remaining 10% shared by Illinois (21,145 kg), Kentucky (3,906 kg) and Missouri (2,188 kg).

The international trade in specimens of *P. spathula* since the species was listed in Appendix II in 1992 has consisted principally of captive-bred live fish and fertilised eggs. Exports from the USA increased during 1993 to 1997, with 9,644 to 265,300 live fish and 60,000-187,000 fertilised eggs respectively, and then decreased in 1998 to 70,000 live fish and 53,440 fertilised eggs. Results of artificial breeding for the release of *P. spathula* fry in the wild are available, but there seems to be no development of commercial farming for production of meat and caviar of *P. spathula* at present.

In 1992, the USA proposed the species for listing in CITES Appendix II in accordance with Article II, paragraph 2(a). Since Resolution Conf. 10.12 *Conservation of Sturgeon* entered into effect in 1998, recommending that range States "consider the feasibility of establishing annual export quotas for sturgeon specimens", the USA did not establish export quotas for *P. spathula*.

This species is recommended under Decision 11.106 for inclusion in category 2 (ii).

DISTRIBUTION AND POPULATION

The CITES database lists the distribution of *Polyodon spathula* as: Canada (ex), United States (Anon., 2000).

P. spathula is classified as Vulnerable by IUCN (1996):
VU C2b USA [Mississippi River].

P. spathula was previously abundant throughout the Mississippi River Basin and some adjacent estuaries of the Gulf of Mexico, and has been observed in the Great Lakes including Canadian waters. *P. spathula* is also found throughout the Missouri River basins and many of their tributaries (Graham, 1997). There has been no recent verification of the occurrence of *P. spathula* in Canada. The species is found in 22 states in the USA. However, whilst several paddlefish populations appear to be stable or increasing, the species is in decline in much of its current range due to continued habitat modification and degradation, increased water contamination and overfishing (Graham, 1997). Information presented in Graham (1997), was compiled in 1994.

In 1994, the population was reported to be stable in Arkansas, Indiana, Iowa, Kentucky, Louisiana, Minnesota, Mississippi, Missouri, Montana, Oklahoma, and South Dakota (Graham, 1997). Graham (1997) also reported that wild populations are considered to be increasing in Texas, West Virginia and Wisconsin, whilst they have been declining in Illinois and North Dakota. The species has been extirpated from Maryland, New York, North Carolina and Pennsylvania (Gengerke, 1986, cited in Todd, 1998).

Alabama: *P. spathula* is the only species of Acipenseriformes confirmed to exist in this state (US Fish and Wildlife Service (USFWS), CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000).

Kansas: *P. spathula* is present in the Arkansas, Kansas, Marais des Cygnes, Missouri, Neosho and Verdigris rivers. In all these rivers, restocking efforts have been undertaken and the population is considered to be stable and probably increasing (Graham, 1997).

Nebraska: The population is considered stable to declining (Graham, 1997) (see South Dakota, Missouri River common boundary: below Gavins Point Dam, near Yankton in South Dakota (USFWS, *in litt.* to TRAFFIC Europe, September 2000)).

Ohio: According to Graham (1997), the status of the *P. spathula* population is unknown (Graham, 1997). However, state authorities consider the species to be limited in distribution and relatively rare (USFWS, *in litt.* to TRAFFIC Europe, September 2000).

South Dakota: Graham (1997) considers the population to be stable. Currently, the paddlefish population below Gavins Point Dam contains too few mature individuals to be attractive to the caviar trade and the remaining paddlefish populations (in the Missouri River above Gavins Point Dam) are not abundant enough to attract interest (USFWS, *in litt.* to TRAFFIC Europe, September 2000).

Virginia: The status of the population is unknown (Graham, 1997).

HABITAT AND ECOLOGY

This species prefers large rivers and lakes. Paddlefish normally inhabit deep water, but during the summer can be found near the surface and at variable depths. *P. spathula* grows to 0.4-0.7 kg in the first year and the largest specimens caught measured 2.3 m and weighed 80 kg. Sexual maturity occurs at 5-9 years for males and 8-12 years for females with 1-1.2 m total body length (Hochleithner and Gessner, 1999). Southern stocks generally grow more quickly than their northern counterparts and rarely exceed 15 years of age, however, northern stocks have been recorded as old as 40-50 years (Scarnecchia and Graham, 1999). Spawning occurs in early spring but adults are unlikely to spawn every year (Hochleithner and Gessner, 1999). Eggs hatch in 12-14 days.

The paddlefish's large mouth strains small crustaceans, algae, ephemeral larvae and mayflies. They are also occasional bottom feeders. During their first year, they are generally particle feeders, after which they begin to filter-feed. It has been suggested that their paddle-like bill senses plankton concentration levels (Hochleithner and Gessner, 1999).

THREATS TO SURVIVAL AND DOMESTIC USE

A major threat to paddlefish is the damming of large rivers. This has eliminated much of their spawning habitat by controlling spring flooding and keeping water within riverbanks. Intentional and unintentional snagging by anglers also results in mortality. Whilst the paddlefish is heavily parasitized by lampreys *Ichthyomyzon* spp., the impact of this is unknown.

An assessment of the impact of fisheries on wild populations of *P. spathula* has not been possible due to the lack of data. Many states, particularly Arkansas and Mississippi, do not require commercial fishermen to report their catches. According to surveys and voluntary reporting, in 1992, 65% of the catch recorded in five of the six states that still allow commercial fishing of *P. spathula* came from Arkansas (Graham, 1997). In Mississippi, no surveys were conducted and no catches were voluntarily reported.

Unless otherwise stated, information hereunder has been provided by USFWS (CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000).

Of all states in the USA where *P. spathula* occurs (Graham, 1997),

- some allow commercial fishery: Arkansas, Illinois, Kentucky, Mississippi, Missouri and Tennessee;
- some prohibit the commercial fishery, but allow sport fishing: Indiana, Iowa, Kansas, Montana, Nebraska, North Dakota, Oklahoma and South Dakota; and
- others do not have large enough populations for a viable commercial fishery and therefore prohibit both commercial and sport fishing: Alabama, Louisiana, Maryland, Minnesota, New York, North Carolina, Ohio, Pennsylvania, Texas, Virginia, West Virginia and Wisconsin.

Reported catch of *P. spathula* in commercial fishing states (kilogrammes)

	1989	1990	1992	1996	1997
Arkansas	?	?	158,500	?	?
Illinois	73,473	?	21,150	?	22,289
Kentucky	?	?	3,910	?	?
Missouri	25,850	?	2,500	4,411	?
Tennessee	?	220,610	60,330	29,411	?
Total	?	?	242,000	?	?

Source: Graham, 1997

The 1992 official reported catches suggest that the largest volumes were caught in Arkansas (65%) and Tennessee (25%) (Graham, 1997).

The average combined annual catch of *P. spathula* for the late-1990s in Illinois, Kentucky, Missouri and Tennessee was estimated at 111.8 t. Compared to data collected by Graham (1997), a sharp drop was recorded in Tennessee (24.9 t in 1999) while other states reported increased catches, such as 62.1 t in Kentucky in 1999 and 2.5 t in Missouri in 1998.

States allowing commercial fishing

Arkansas: Commercial harvest of *P. spathula* in 1992 was estimated at 136-181 t (Graham, 1997). From 1990 to 1999, the Arkansas Game and Fish Commission collected USD150,000-300,000 annually in revenues for the sale of commercial and paddlefish sport fishing licenses, permits and tags. There is no compulsory reporting system for reporting commercial catch data to the authorities in Arkansas. Based on voluntary reporting of commercial fisheries, 69 kg of paddlefish roe (caviar) were harvested in 1998 (total catch of paddlefish estimated at 410 kg), while 230 kg average annual harvest of caviar (worth USD16,256) was reported for 1987 and 1988.

Illinois: The reported commercial catch of *P. spathula* was 73,473 kg in 1989 and 22,289 kg in 1997 (with 33 kg of roe –caviar) (Todd, 1998).

Kentucky: *P. spathula* are caught for commercial purposes in Kentucky.

Catch statistics for *P. spathula* (kilogrammes)

	1999 Jan.-June 2000	
Meat	59,766	84,254
Caviar	2,375	5,645
Total	62,141	89,899

Source: CITES Management Authority of the USA, *in litt.* to TRAFFIC Europe, September 2000

Mississippi: No catch data were reported. Historically, Mississippi has not collected data on *P. spathula* catch. There is no compulsory reporting system to collect either commercial or sport fishing data. No scientific publication has reported past or present paddlefish catches in this State, but Graham (1997) notes that high commercial harvest may have occurred in the mid-1980s. Law enforcement intelligence

suggests that *P. spathula* are being illegally caught by commercial fishermen for their roe (caviar). Poaching is an ongoing problem, consisting not of isolated infrequent occurrences, but rather being organised and secretive operations (D. Ruche, Mississippi Department of Wildlife, Fisheries and Parks, *in litt.* to the USFWS, 16 August 2000).

Missouri: The 1989 estimated catch of *P. spathula* was 25,850 kg.

Annual catch of *P. spathula* in the Mississippi River (section in the State of Missouri) (kilogrammes)

1992	1993	1994	1995	1996	1997	1998
2,344	3,758	2,033	2,911	4,411	1,054	2,504

Source: Todd, 1998

Tennessee: Commercial harvest of *P. spathula* is allowed. The sport fishery harvest is not significant in volume. In 1975, the commercial paddlefish catch reached 197,768 kg.

Commercial catch of *P. spathula* (kilogrammes)

	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Catch	220,651	47,627	60,444	53,885	59,218	57,100	29,417	40,503	24,234	24,899
Roe (caviar)			84.8	724.4	565.6	683.1	63.5	283.9	855.5	1,404.3

Source: CITES Management Authority of the USA, *in litt.* to TRAFFIC Europe, September 2000

Paddlefish roe is an important source of income for commercial fishermen. The flesh (meat) is of less importance and is sold both within and outside Tennessee. Poaching has been reported to the authorities, but the state authorities claimed that it is difficult to evaluate during the fishing seasons because enforcement officers are too busy administering the permits and tags.

States allowing sport fishing only

Indiana: The state's estimated annual harvest was 10,000 kg for 1996 (Hesse and Carreiro, 1997, cited in Todd, 1998).

Iowa: Only sport harvest (snagging) of paddlefish and sturgeon is allowed. Two fish are permitted daily, with a possession limit of four. Sport fishing (snagging) has been prohibited in the Big Sioux and Missouri rivers since 1993. Data on sport fishing were not provided by the state authorities. Some illegal harvest of paddlefish is suspected by the authorities, but has not been confirmed.

Kansas: *P. spathula* sport snagging fisheries are limited to specific locations. They are significant local events attracting anglers from other neighbouring towns that contribute to the economy of relatively small Kansas towns. There are no official reports of poaching, but it is suspected that illegal activities occur.

Number of *P. spathula* harvested by anglers at sport fishery locations (number of fish caught)

	1992	1993	1994	1995	1996	1997	1998	1999	2000
Marais des Cygnes	4	87	19	12	11	75	58	457	5
Neosho River	71	541	325	769	479	394	800	2,010	98
Total	75	628	344	781	490	469	858	2,467	103

Source: USFWS, CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000

Montana: In the late 1980s-early 1990s, an estimated 1,360-4,536 kg of *P. spathula* roe has been collected each spring during the sport snagging (Graham, 1997). From 1990 to 1994, a total of 2,270 kg of *P. spathula* roe was collected annually from a total of 7,000 fish caught at the Intake Diversion Dam east of Glendive (Wilkinson, 1997). Several fishing grounds are on the Montana-North Dakota border. The annual average number of *P. spathula* harvested in Montana from 1995 to 1999 was 1,207 (ranging from 717 in 1998 to 1,706 in 1999). With an average body weight of 19 kg (Scarnecchia *et al.*, 1996), the annual catch would represent 23 t.

Nebraska: There is no commercial fishery of *P. spathula*. Sport fisheries of the species are regulated. Annual catch quotas of 275 fish for bow and arrow and 1,400 fish for snag fishing are allocated. Quotas

are enforced by distributing no more than two tags per fisher. Poaching occurs on the Missouri River. Investigations following on from a 'sting operation' are on-going. (see South Dakota for combined harvest estimates for *P. spathula* snagging fishery only).

North Dakota: During 1993 and 1994, 2,268 kg and 1,726 kg of *P. spathula* roe were respectively collected (Graham, 1997). The annual average number of *P. spathula* harvested in North Dakota from 1995 to 1999 was 1,356 (ranging from 800 in 1997 to 1,970 in 1998). With an average body weight of 19 kg (Scarnecchia *et al.*, 1996), the annual catch would represent 25.8 t. The average annual volume of caviar produced from 1993-2000 by a company collecting the sport fishery catch was 2,705 kg/year, ranging from 4,127 kg in 1998 to 1,537 kg in 1997.

Oklahoma: No catch data were reported by the state authorities.

South Dakota: Only harvest of *P. spathula* by sport anglers during a limited season and on a limited ground (common boundary with Nebraska) is allowed.

Number of *P. spathula* caught by sport fishers along the common South Dakota and Nebraska boundary during the snagging season (number of fish)

	1992	1993	1994	1995*	1996*	1997	1998	1999
No of fish	1,000	1,529	1,568	2,200	1,828	948	1,125	1,334

* quota exceeded due to delay in closing the season caused by the administrative procedure required.

Source: USFWS, CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000

The harvest was managed with an annual 1,600 fish quota system from 1992-1996, and by a tag/permit system from 1997-1999. There are no data for 2000 because the snagging season starts on 1 October. An estimated additional 100-150 paddlefish are harvested annually during the July archery season.

States prohibiting commercial and sport fisheries

Alabama: The species is protected by law but no poaching activity is known or suspected.

Louisiana: There has been no legal commercial harvest of *P. spathula* in Louisiana since 1986. There have been complaints to the state authorities about illegal catch and possession. Recently, a request for a permit to transport and export 5,000 lbs (2,268 kg) of paddlefish roe per month was submitted to the state authorities.

Ohio: No catch data were reported by the state authorities. No poaching has been recorded by the authorities.

Texas: No information provided.

Virginia: Only historic anecdotal reports of fishing.

West Virginia: No information provided.

Wisconsin: *P. spathula* harvest has been prohibited since 1992. No poaching has been recorded, except for a recent case of more than 60 paddlefish incidentally caught in a large seine net and intentionally killed by the commercial fisher.

Introduced populations:

China: Significant imports (see "International Trade" and "Captive Breeding") of live *P. spathula* were introduced in captive breeding facilities in China. There is no documentation on the potential damage of the introduction of exotic Acipenseriformes on local species. If specimens of *P. spathula* escape in to the Yangtze River, this may threaten the Chinese paddlefish *Psephurus gladius*, which is on the brink of extinction. Experts have agreed that the potential threats that must be taken into consideration are: the

introduction of new pathological germs in wild populations, food competition in case of acclimatisation of the exotic specimens and hybridisation with the local species (Cemagref, *in litt.*, press release, 26 January 2000).

INTERNATIONAL TRADE

P. spathula is one of four Acipenseriformes species listed in the CITES Appendices before 1997. The listing was proposed by the USA at the 8th meeting of the Conference of the Parties in 1992.

Gross exports and comparative tabulation of trade in *P. spathula* from 1992 to 1998 are given in the Appendix. Annual Reports from 1992 to 1998 suggest that most specimens of *P. spathula* traded internationally are fertilised eggs and live fish. Trade volumes increased progressively from 1993 (30,000 fertilised eggs and 9,644 live fish) to 1997 (187,000 and 238,700 respectively), before decreasing in 1998 to 70,000 fertilised eggs (as confirmed by China, the only importer in 1998 (CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000)) and 23,440 live fish. China is the main destination for both fertilised eggs (54%, 60,600 eggs/year in average from 1994-1998) and live specimens (31%, 42,000 fingerlings/year in average from 1994-1997). Other major importers (in decreasing order of importance) are: Romania, Hungary, France, Germany, Taiwan (Province of China) and Japan. These specimens are supplying the demand of captive breeding facilities, in Moldova for instance (see "Captive Breeding").

A market for meat seems to exist since small volumes of meat were exported from the USA to Canada in 1995 and 1996 (46 kg and 381 kg respectively).

Only one export quota was established since 1998. It was set by the Russian Federation at 5 kg of fertilised eggs for 2000. Three live specimens (possibly adults) were imported into the Russian Federation from the USA in 1998.

CONSERVATION MEASURES

Both commercial and sport fishing are prohibited in Alabama, Louisiana, Maryland, Minnesota, New York, North Carolina, Ohio, Pennsylvania, Texas, Virginia, West Virginia and Wisconsin (Graham, 1997). Unless otherwise stated, information for this section was provided by the USFWS (CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000).

States allowing commercial fishing

Arkansas: For commercial fisheries, seasonal restrictions apply in some areas. A 76 cm eye-to-fork-of-tail minimum length limit from fall until spring is set as additional protection for females (Graham, 1997). For sport fishing, a daily catch limit of two paddlefish is set.

Illinois: No information provided.

Kentucky: The catch is regulated under the commercial fishing and snagging legislation which controls the type of gear, the fishing requirements (e.g. number and size of fish) and the waters open to fishing.

Mississippi: Sport and commercial fisheries are permitted. However, during the period 1 November to 30 April, it is illegal to fish for, take or possess paddlefish or its parts, including roe (caviar). A list of areas closed to the use of commercial fishing gear exists.

Missouri: *P. spathula* is classified as game, so sport and commercial fisheries are permitted.

Tennessee: Strict regulations exist for both commercial and sport harvest of *P. spathula*. Only one area permits sport fishing, with a limit of one fish per day. Paddlefish and its parts caught by sport fishing methods may not be sold.

States allowing sport fishing only

Indiana: Sport fishery is limited to hook-and-line (Graham, 1997).

Iowa: The commercial fishery was closed in 1986 (Graham, 1997). There is no size limit for sport fisheries. Together with neighbouring states, Iowa closed *P. spathula* sport fisheries on the Missouri and Big Sioux rivers in 1992.

Kansas: State regulation for sport fishing of *P. spathula* is strict. It includes, for instance, reporting of the catch at a check station where a numbered tag is attached to each fish caught. Restocking efforts have been undertaken in all rivers. The fry are reared in a state-run hatchery, using seedlings (eggs or fry) from Missouri or Oklahoma. The policy is to restock the reservoirs of rivers with fish that were obtained from the nearest existing stock. All stock must be tagged with coded wire tags.

Number of *P. spathula* released through restocking by the Kansas Department of Wildlife and Parks in Kansas and Oklahoma (number of fry)

	1992	1993	1994	1995	1996	1997	1998	1999	2000
Number of fry	22,267	11,685	16,918	6,196	0	0	0	0	7,100

Source: USFWS, CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000

Montana: Sport fishing only is allowed. The main fishing area is common to Montana and North Dakota. Regulations of both states are therefore linked. A tag limitation programme has been installed and the catch is allowed only from 1 May to 15 June in a limited area of the Missouri River. The average number of tags distributed from 1995 to 1999 is 6,467/year.

Nebraska: The state shares its *P. spathula* fishery with South Dakota along the Missouri River common boundary. Quotas of 275 and 1,400 fish (individuals) have been adopted for bow and arrow and snag fishing respectively. As in South Dakota, a slot length limit of 88-114 cm and fishing seasons were established for the two types of sport fisheries (see South Dakota).

North Dakota: Only sport fishing is permitted. The main fishing area is common to Montana and North Dakota. Regulations of both states are therefore linked. A tag limitation programme has been installed and catch is allowed only from 1 May to 15 June in a limited area of the Missouri River. The average number of tags distributed from 1992 to 2000 is 5,703/year. A research programme on *P. spathula* began in 1990. This includes research on reproduction, age structure and catch rates by the use of 10,000 micro-tags. Eight years of data suggest that the catch rate is closely correlated to the population constraints. The caviar of fish caught by sport fishermen is collected by an organisation, North Star Caviar, Joint Venture, and sold on the local and international market. The provisions of the sale of paddlefish products in North Dakota have been defined through a memorandum of understanding. This was signed in June 2000, between North Star Caviar, Joint Venture and the North Dakota Game and Fish Department. The benefit of the trade in caviar must be invested in "resource-related" projects.

Oklahoma: The sport fishery of *P. spathula* is strictly regulated, including fishing seasons, daily catch limits, restricted fishing gears and the obligation to tag every fish caught. Some *P. spathula* are being released in Oklahoma rivers by the state of Kansas (see above). The Oklahoma restocking programme is monitored.

Number of *P. spathula* fry released through the Oklahoma restocking programme (number of fry)

Reservoir	1992	1993	1994	1995	1996	1997	1998	1999	2000
Kaw	18,890	25,185	16,750	2,013	?	?	?	?	?
Oologah	?	?	?	5,974	112	10,719	2,037	8,837	3,450
Texhoma	?	?	?	?	?	?	?	5,757	20,846
Total	18,890	25,185	16,750	7,987	112	10,719	2,037	14,594	24,296

Source: USFWS, CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000

South Dakota: *P. spathula* catch is managed jointly with Nebraska, due to the common boundary on the Missouri River. The fishing seasons are limited to 16 days in the middle of July for archery fishermen, and to 30 days in October for the snagging fishery. A catch quota also applies to the latter. From 1992 to 1996, the catch quota was reached in six days or less. As of 1997, the management of the paddlefish snagging fishery was switched to a tag/permit system and a total of 2,250 tag/permits were issued annually from 1997 to 1999 by South Dakota and Nebraska. Several other restrictions have been adopted for the snag fishery. *P. spathula* measuring 35-45 inches (88-114 cm) must be released, and snagging is allowed only from 7:00 to 19:00 daily. Archery anglers can fish from sunrise to sunset. Graham (1997) noted that all anglers are prohibited from selling or trading any paddlefish or sturgeon flesh (meat) and/or body parts. Since 1990, the state has been restocking juveniles.

States prohibiting commercial and sport fisheries

Alabama: *P. spathula* is protected by State Regulation 220-2-94: "It is illegal to take or attempt to take paddlefish from all public waters of Alabama by any method or to process paddlefish or any part of a paddlefish. Any paddlefish accidentally captured shall be immediately returned to the waters from where it came. This regulation does not apply to commercially packaged paddlefish products imported from out-of-state, or to paddlefish cultured at hatchery operations or grow out ponds permitted through the Commissioner, Department of Conservation and Natural Resources".

Louisiana: *P. spathula* has been protected since 1986. No sport or commercial fisheries are allowed. It is illegal to possess paddlefish, sturgeon and their body parts, including roe (caviar). Violations carry a maximum fine of USD2,500/fish. There is a management plan for each species of paddlefish and sturgeon occurring in the state, including population monitoring, habitat restoration and stock enhancement. Captive breeding and restocking efforts of *P. spathula* have been carried out in Louisiana for 10 years.

Minnesota: *P. spathula* fishing has been closed statewide, but there is no regulation, nor restocking programme.

New York: Since the population is too low to be of interest either to commercial or sport fisheries, *P. spathula* has no official status or special protection. Fingerling restocking was initiated in 1998, and if there is significant survival of these, prohibition of harvest will be adopted by the fall of 2002.

Ohio: *P. spathula* was listed as an Ohio Endangered Species from 1974 until 1987 when it was re-listed as Threatened. However, this does not afford any legal or regulatory status. Ohio does not have a commercial or sport catch/quota for the species. *P. spathula* cannot, however be taken within 1,000 feet downstream of four large dams in the state. Ohio is currently participating in research and outreach programs for *P. spathula*.

Pennsylvania: Commercial and sport fisheries are prohibited. The state is engaged in a long-term restoration effort for paddlefish in the major rivers of western Pennsylvania. Individuals released through the programme should have reached sexual maturity.

Texas: The state is actively restocking and there are indications that the populations is increasing (Graham, 1997).

Virginia: Classified as Threatened (Graham, 1997).

West Virginia: The state began stocking juveniles in the Ohio and Kanawha rivers in 1992 (Graham, 1997).

Wisconsin: *P. spathula* has been assigned the status of Threatened in 1989; possession is illegal.

CAPTIVE BREEDING

An aggressive restocking programme for *P. spathula* has been undertaken in many states in the USA (Graham, 1997). The success of experimental releases of *P. spathula* has been investigated (Beamesderfer and Farr, 1997).

According to FAO Fishstat, the world annual production of farmed sturgeon and paddlefish (mostly for the meat market) has increased rapidly, from 160 t in 1987 to 2,576 t in 1998. Species specific data are not available in the FAO database, but according to available information, there seems to be no current significant production of caviar and/or meat of *P. spathula* in commercial farms (USFWS, CITES Management Authority of USA, *in litt.* to TRAFFIC Europe, September 2000).

China: In 1988, the Fishery Institute of Xiantao City in central China's Hubei Province started to introduce a small quantity of fertilised eggs of *P. spathula* from the USA to conduct experimental grow-out in ponds. The institute did not succeed with hatching and rearing until 1991. Since 1991, China imported an annual average of 100,000 fertilised eggs from the USA. However, no individual has been raised to adult age until now (Dr Meng Xianlin, CITES Management Authority of China, *in litt.* to TRAFFIC Europe, 28 August 2000).

Moldova: *P. spathula* was introduced in the country to study multi-species captive breeding, with Stellate Sturgeon *Acipenser stellatus* in grow-out ponds (A. Vedrashco, pers. comm. to TRAFFIC Europe, 14 March 2000).

Poland: The CITES Management Authority of Poland has confirmed that six sturgeon species and their hybrids, including *Polyodon spathula*, are bred in captivity. In most cases, hybrids are F1 generation live specimens (fertilised eggs) imported into Poland from range States, particularly from the Russian Federation. Some farms keep F2 generation specimens, bred in Poland and used as broodstock, partly for the production of juveniles to be sold on the domestic market and exported (Dr Z. Krzeminski, Deputy Director, *in litt.* to TRAFFIC Europe, 31 August 2000).

Russian Federation: At least one large facility is rearing *P. spathula* (Anon., 1999).

USA: Restocking programmes have started in Louisiana, Kansas, New York, Oklahoma, Pennsylvania, South Dakota, Texas and West Virginia. There are no records of commercial captive breeding or aquaculture activities within Arizona, Delaware, Florida, Louisiana, Minnesota, New York, Ohio, Oklahoma, South Dakota, Tennessee, Utah, Wisconsin and Wyoming. Farms and hatcheries have been developed under state or commercial initiatives to produce fingerlings for restocking (Missouri), conduct short term grow-out for research (Wisconsin), refine captive propagation techniques (Georgia) and develop intensive grow-out culture (Texas) (Andreasen, 1999).

REFERENCES

- Andreasen, L. (1999). Captive propagation as a recovery tool for North American sturgeon. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington DC, USA. Pp. 121-129.
- Anon. (1996). *A Guide to Interpreting Outputs from the WCMC CITES Trade Database*, Produced by the World Conservation Monitoring Centre, Cambridge UK. Pp. 16
- Anon. (1999). *Research Center for Commercial Sturgeon Production (BIOS), Russia. Leaflet. 17pp.*
- Anon. (2000). <http://www.cites.org/eng/dbase/fauna->
- Beamesderfer R.C. and Farr R.A. (1997). Alternatives for the protection, restoration of sturgeons, their habitat. *Env. Biol. Fish.*, 48 (1-4): 407-417.
- Graham, L.K. (1997). Contemporary status of the North American Paddlefish, *Polyodon spathula*. In: V. Birstein, J.R. Waldman, and W.E. Bemis (ed). *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publishers, Dordrecht. Environment Biology of Fishes 48: 279-289.
- Hochleithner, M. and Gessner, J. (1999). *The Sturgeon and Paddlefishes (Acipenseriformes) of the World: Biology and Aquaculture*. AquaTech Publications, Kitzbühl. 165 pp.
- IUCN, 1996. *The 1996 Red List of Threatened Animals*. IUCN, Gland, Switzerland.
- Scarnecchia, D.L., Stewart, P.A., and Power, G.J. (1996). Age Structure of the Yellowstone-Sakakawea Paddlefish Stock, 1963-1993, in Relation to Reservoir History. *Transactions of the American Fisheries Society* 125:291-299.
- Scarnecchia, D.L and Graham, L.K. (1999). Paddlefish life history and challenges facing Paddlefish in North America. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on*

the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 40.

Todd, R.M. (1998). Sturgeon and Paddlefish commercial fishery in North America. In: Williamson, D.F., Benz, G.W. and Hoover, C. (eds.) *Proceedings of the Symposium on the Harvest, Trade and Conservation of North American Paddlefish and Sturgeon*, May 7-8, 1998, Chattanooga, TN. TRAFFIC North America/WWF, Washington, DC, USA. Pp. 42-48.

Wilkinson, T. (1997). Plight of the Paddlefish. *Wildlife Conservation*. March/April 1997. Pp. 52-55.

APPENDIX

Gross exports* of *Polyodon spathula* 1992-1998

TAXON	TERM	UNIT	Exp	199	1993	1994	1995	1996	1997	1998
				2						
<i>Polyodon spathula</i>	bodies	US	0	0	0	0	0	0	3,500	0
<i>Polyodon spathula</i>	eggs	US	0	60,000	150,305	88,000	160,000	187,000	180,000	
<i>Polyodon spathula</i>	eggs	kg	US	0	0	0	0	0	0	7
<i>Polyodon spathula</i>	live	GB	0	0	90	600	500	400	0	0
<i>Polyodon spathula</i>	live	HU	0	0	0	31,300	1,000	3,000	0	0
<i>Polyodon spathula</i>	live	US	30	9,644	212,240	57,104	90,004	265,300	53,440	
<i>Polyodon spathula</i>	meat	kg	US	0	0	0	46	381	0	0
<i>Polyodon spathula</i>	specimens	US	0	0	0	0	0	0	0	9

* **Definition:** "Gross exports are the sum of all reported exports and re-exports in a particular commodity or species in a particular year or series of years; Gross trade is thus a simple measure of the total number of items recorded in international trade. However, gross trade may be an overestimate of the total number of actual specimens in trade as re-exports are not deducted from the total." (Anon, 1996). This is particularly applicable to caviar of which shipments are often exported, re-packaged and re-exported. For instance, in the mid-1990s, Germany re-exported as much as 45% of its annual caviar imports (De Meulenaer and Raymakers, 1996).

Comparative Tabulation of trade in *Polyodon spathula* 1992-1998

Year	Imp	Exp	Origin	Imports reported			Exports reported				
				Quantity	Unit	Term	P	S	Quantity	Unit	Term
Eggs											
1993	DE	US		60,000	eggs	S	C	30,000	Eggs	T	C
1994	AT	US		10,000	eggs	T	C				
1994	CN	US						63,000	Eggs	T	C
1994	FR	US						2,000	Eggs	T	C
1994	GB	US		305	live	T	W	305	Eggs	T	C
1994	HU	US						15,000	Eggs	T	C
				Total 1994:	10,305			80,305			
1995	CN	US						40,000	Eggs	T	U
1995	CN	US						10,000	Eggs	T	C
1995	FR	US		10,000	eggs	T	W	10,000	Eggs	T	C
1995	HU	US						25,000	Eggs	T	C
				Total 1995:	10,000			85,000			
1996	CZ	US		20,000	eggs	T	F	20,000	Eggs	T	W
1996	HU	US						100,000	Eggs	T	W
1996	IL	US						20,000	Eggs	T	W
				Total 1996:	20,000			140,000			
1997	CN	US						120,000	Eggs	T	O
1997	GR	US						10,000	Eggs	T	
1997	HU	US						7,000	Eggs	T	
1997	IL	US						50,000	Eggs	T	
				Total 1997:				187,000			
1998	CN	US						150,000	Eggs	T	
1998	CN	US		70,000	eggs	B	C	30,000	Eggs	T	
				Total 1998:	70,000			180,000			
1998	CZ	US	RU	3 kg	eggs	T	F				
1998	SG	US		4 kg	eggs	T	W				
				Total 1998:	7 kg						

Comparative tabulation of trade in *Polyodon spathula* 1992-1998 (continued)

Year	Imp	Exp	Origin	Imports reported				Exports reported				
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P
Live												
1992	JP	US						30		Live	T	C
1993	DE	US						1,500		Live		C
1993	HK	US						644		Live	T	C
1993	JP	US						3,000		Live	T	C
1993	TW	US						4,500		Live	T	C
Total 1993:								9,644				
1994	JP	GB	US	90	live	T	C	90		Live	T	C
1994	CN	US						20,000		Live	T	C
1994	DE	US		60,000	eggs	T	C	60,000		Live		C "eggs" or "live"?
1994	FR	US		80,000	live		C	20,000		Live	T	C i.e. fertilised eggs?
1994	HU	US						40,000		Live	T	C
1994	IT	US		5	live	S	C	5,000		Live		U
1994	JP	US		1,060	live	T	C	3,060		Live	T	C
1994	TW	US						3,875		Live	T	C
Total 1994: 141,155								152,02				
								5				
1995	JP	GB	US	600	live	T	C	600		Live	T	C
1995	AT	HU						800		Live		C
1995	BE	HU	US					2,000		Live		C
1995	DE	HU						5,000		Live	T	C
1995	FR	HU	US					1,500		Live		C
1995	GB	HU	US	1,000	live	T	C	1,000		Live		C
1995	GR	HU	US					21,000		Live		C
1995	DE	US		25,000	live	T	C					
1995	DE	US		2,500	live	T		2,500		Live	T	U
1995	FR	US						10,000		Live	T	C
1995	GB	US		600	live	T	C	600		Live	T	C
1995	GB	US		4	live	B	C	4		Live		W
1995	JP	US		3,000	eggs	T	F	4,000		Live	T	C "eggs" or "live"?
1995	MX	US						5,000		Live	T	U i.e. fertilised eggs?
1995	RO	US						10,000		Live	T	U
Total 1995: 32,704								64,004				
1996	JP	GB	US					500		Live	T	C
1996	AT	HU	US					600		Live		F
1996	BE	HU	US					400		Live		F
1996	CN	US						35,000		Live	T	C
1996	DE	US		20,000	eggs	T	C	20,000		Live	T	C "eggs" or "live"?
1996	DE	US		4	live	Z	C	4		Live		W i.e. fertilised eggs?
1996	FR	US		10,000	live		W	960		Live	T	C
1996	GB	US		1,500	live	T	C	1,000		Live	T	W
1996	JP	US		2,000	live	T	F	1,500		Live	T	C
1996	JP	US						1,500		Live	T	W
1996	RO	US						10,000		Live	T	C
1996	TW	US						8,500		Live	T	C
1996	TW	US						1,000		Live		C
1996	TW	US						1,000		Live	T	W
Total 1996: 33,504								81,964				

Comparative Tabulation of trade in *Polyodon spathula* 1992-1998 (continued)

Year	Imp	Exp	Origin	Imports reported			Exports reported					
				Quantity	Unit	Term	P	S	Quantity	Unit	Term	P
Live (continued)												
1997	MY	GB	US						400	Live	T	C
1997	PL	HU	US						3,000	Live	T	F
1997	AT	US		15,000	live		T	F				
1997	CN	US							113,000	Live	T	C
1997	FR	US		30,000	live			F	15,000	Live	T	C
1997	GB	US		400	live		T	C	400	Live	T	O
1997	JP	US		3,500	bodies		T	F	3,500	Live	T	O
1997	RO	US							100,000	Live	T	C
1997	TW	US							2,400	Live	T	C
1997	TW	US							1,000	Live	T	O
Total 1997:				48,900					238,700			
1998	FR	US		45,000	live		T	F	15,000	Live	T	W
1998	JP	US							2,100	Live	T	O
1998	JP	US							340	Live	T	C
1998	RO	US							6,000	Live	T	W
Total 1998:				45,000					23,440			
Specimens												
1998	DE	US		6	specimens		S	O	6	Specimens		O
1998	RU	US							3	Specimens		O
Meat												
1995	CA	US							46 kg	Meat	T	W
1996	CA	US							381 kg	Meat	T	W